eyes, is scarcely less distressing, being closely applied to the face, and cannot render that soft shade which European hats and bonnets afford.*

Pterigium.—This disease of the eye is very common.

Cataract.—I have operated on eight persons with various degrees of success, but such operations are performed under great disadvantages without an hospital.

Intermittent Fever.—It is difficult to assign an evident cause for the prevalence of this complaint. In the neighbourhood of Beyrout, at least, where the land is parched with drought one half of the year, and where, at no period, marshes or stagnant pools are to be found to any extent, marsh miasma would seem to be out of the question; however, if we omit the association of *marsh*, which is but a casual circumstance, the theory of decomposed vegetable and animal matter will remain as probable as any other, and as applicable. These effluvia are often more offensive from the absence of water.† Intermittent fever is here found among all classes and ages, and frequently recurs several times when the patient is apparently quite well, and it hangs about him for eight or ten weeks together.

Continued Fever is generally of the common inflammatory form. I have only met with two or three cases which assumed a typhoid character.

Rheumatism.—The prevalence of rheumatism manifestly arises from sleeping on the ground in the open air. I seldom find it in

* There is much truth in these observations; but I think we may justly ascribe this "scourge of oriental climates" to the sultry heat of the vallies, exposure to the mid-day sun, check of perspiration caused by the heavy dew of the night, and other circumstances, which, producing irregular determination of blood, favour congestion of the liver and spleen; and this seems to be borne out by the fact that persons of a bilious temperament, who have swarthy complexions, black eyes, and black hair, and generally relaxed bowels, suffer less from ophthalmia, but are the frequent subjects of dysentery and diarrhoea; whereas those of a sanguine temperament, who have light eyes and light hair, a deficiency of bile, and a clearer skin, often escape the latter, but are very prone to constipation and ophthalmia.—*Wm. Holt Yates, M.D.*

† Having been myself a sufferer at Beyrout from this complaint, and witnessed its effects in others, I took some pains to investigate the subject. With your permission, therefore, I will avail myself of an early opportunity of sending you a few remarks on the climate of Syria, as connected with the points referred to in this communication.—*Wm. Holt Yates.*

those who have not thus exposed themselves.

Pulmonary Consumption.—So far as I have been able to judge, this is a rare disease. Bronchitis is the most frequent pulmonary complaint in Syria.

Urinary Diseases.—These are chiefly chronic affections of the mucous membrane, and irritation arising from the habitual use of stimulants.

Ulcers.—These arise from trivial causes, and are generally most obstinate to heal.

Fungous Tumours.—One of these occupied the entire place of the cornea, hung over the cheek, and presented a bleeding surface. I passed a hook through it, and removed it, together with a portion of the globe. It appeared to have originated in a thickened conjunctiva.

Club-Foot.—I operated in two cases with success, which gave great pleasure to the country-people who saw them.

N.B.—Those diseases which might be influenced by season I have marked in months, as they occurred, so that at the end of the year, some useful calculations may be made on the subject.

THOS. KERNS.

NOTE.—*To the Editor*.—Sir: There is nothing extraordinary in coincidences in discoveries, which, however, lose somewhat the semblance of originality when improvements or suggestions having been recorded in them, they are, after the lapse of years, again announced as new. I am led to these remarks by observing two propositions, originally made by me, which have been treated in a similar manner. The first was my project for the employment of internal pressure on the abdominal aorta in uterine hæmorrhage, commented on in *THE LANCET* for Feb. 23, 1828 (No. 234), and published in the "London Medical Repository" three years previously (1825), with cases, and subsequently followed by a notice in Oct., 1828, of this novel practice by Dr. Eichelberger and Von Siebold, at Berlin. The second is the description of a "New Lithotomy Staff" by Mr. Bryan, of London, with remarks on the operation, published in *THE LANCET*, Feb. 11, 1843, which, in many respects, resembles one proposed by me in the same work, for July 12, 1834, especially in cutting on the *concavity* of the staff, as also in the parallelism and angle of the incisions. This proposition, with the observations thereon, have been honoured with the approval of some eminent operators; but at present I content myself with merely noticing the facts, and referring the profession to the communications above-mentioned. I am, Sir, yours very obediently,

G. W. JAMES, M.R.C.S.

March 6, 1843.