safely and easily do the major operation later by the abdominal route and save many lives in that way.

Dr. D. H. CRAIG, Boston—I think cutting through a wall 8 to 10 cm. in thickness should be kept distinctly out of the field of vaginal opening. When we have pus so firmly walled off as that it is perfectly safe to go through the abdominal wall, then we can clean it out thoroughly and make a complete cure instead of leaving the patient with a lot of material to absorb. As regards the occasional operator, I think the vaginal route is better for him, providing he is sure of his technic. But no man can do more harm than the occasional operator going through Douglas' cul-de-sac into the abdomen. The posterior or anterior cul-de-sac operations are not, as a rule, curative. They are palliative, and if the occasional operator will take that route and do these operations it is incumbent on him to advise his patient of the fact that he is doing a palliative operation and that other work may be required by some one who is not an occasional operator.

Dr. J. H. CARSTENS, Detroit—I wish the essayist would say "opening" instead of "puncture." A "puncture" means a little bit of an opening, and to my mind the opening can not be too big. A little one closes up right away, and you must do your work again. Make a good, big opening, and give Nature a chance.

Dr. C. O. THIENHAUS—When I have a hard mass lying in the cul-de-sac of Douglas, and the diagnosis, whether or not this hard mass contains a focus of pus, can not be ascertained by other means, I advocated in my paper exploratory vaginal puncture for this purpose, and if I find pus I drain by way of the vagina. Dr. Craig says it is better to open the abdomen in such cases. Clinical experience has shown evidently that it is not. When you can not before operating determine the virulence of the bacteria present in the pus, you are liable, when a little of this pus contains highly-virulent infectious material, to see your patients die promptly. When bacteriology is so far advanced that we can determine with absolute certainty the grade of virulence of the bacteria in the pus focus, then we can differentiate and state in what cases abdominal section may be done without fear of infecting the general peritoneal cavity. But up to this time it is always better to be on the safe side and to open intraperitoneal abscesses, where there exists a possibility, by an extraperitoneal route. When the speaker says that cases of double pyosalpinx of gonorrheic origin do not heal by simple vaginal drainage, I would like to cite the statistics of Treub, who states that from 79 cases 44 were radically cured by simple vaginal drainage without further interference. I am sorry that Dr. Carstens misunderstood me. Naturally, when the puncturing needle has demonstrated the presence of pus, the vaginal incision must be as large as possible to establish free drainage.

CARCINOMA AND TUBERCULOSIS TREATED BY THE ROENTGEN RAYS.

G. E. PFAHLER, M.D.

Formerly Assistant Chief Resident Physician in the Philadelphia Hospital, Assistant Physician to the Medico-Chirurgical Hospital, Skiagrapher to the Medico-Chirurgical Hospital, Lecturer on Terminology and Symptomatology, and Demonstrator of Physical Diagnosis in the Medico-Chirurgical College.

PHILADELPHIA.

As a basis for the conclusions to be drawn in this paper, I have selected nine cases, each illustrating some special point of interest. I have omitted many of the details which would be of little interest to the general practitioner.

CASE 1.—Mrs. K. E., aged 84 years. Referred by Dr. Ulrich of Chester, Pa. The diagnosis was carcinoma of the right breast. Operation was contraindicated on account of the patient's age and on account of the condition of her heart. The growth was of one year's duration. There was a distinct retraction of the breast below the nipple, making a groove about two and a half inches in length. In connection with this there

was a hard mass about the size of a hen's egg. Surrounding this central mass there were other masses of lesser size and of less density. The overlying skin was healthy. There was no axillary or supraclavicular involvement. Treatment was begun April 20, 1903, and exposures were given daily for two weeks. Each exposure was of ten minutes' duration, at a distance of 15 inches, with a high vacuum tube and 3 to 4 ampères' current. The treatments were then given three times a week for six weeks, then twice a week for ten weeks, and once a week for 2 weeks. She had 57 treatments in all. The length of each exposure was gradually increased from 10 to 25 minutes. After the first week there was improvement in that the semi-solid masses disappeared. At the end of two weeks the central mass seemed to be a little softer. From this time there was

Fig. 1.—Case 4. Before treatment by x-ray, May 25, 1903. Sarcoma of orbit. Scar due to an operation.

constant improvement, the mass gradually became softer and smaller until it disappeared.

The points of special interest in this case are the early and constant improvement, and the absence of any dermatitis in spite of the unusually long exposures. In no other case have I been able to give nearly so long an exposure without producing a marked dermatitis. As a whole, it is one of the most remarkable cases that I have seen.

CASE 2.—Mrs. K. J., aged 40 years. Diagnosis, recurrent carcinoma of the left breast. She had been operated on twice by Dr. Sanford of Memphis, Tenn. The first operation was performed one month after the growth was first noticed, May 7, 1900. A growth reappeared one year later and was removed December, 1901. A recurrence developed in July, 1902. X-ray treatment was begun in October, 1902, at which time there was

an induration involving the entire scar and about one inch on either side, with nodules in the axilla and the left supraclavicular region. There was also an induration in the right breast. Both mammary, axillary and supraclavicular regions were treated. Seventy-five treatments in all were given, between October 11 and May 31, when the carcinomatous tissue seemed to have disappeared. Treatments have been given since that time at intervals of two to four weeks for the purpose of preventing any recurrence or any involvement of the mediastinal glands. The patient's general health before treatment was failing and she was not expected to live six months. Her general health improved while under treatment. A dermatitis was produced on two occasions, each of which seemed to be followed by distinct improvement.

CASE 3.—Mrs. J. W., aged 48 years. Recurrent carcinoma of the left breast. She was operated on by Dr. Wharton in

reapplied the rays, which produced another burn that developed after leaving the hospital. I advised her to go to the seashore and allow the burn to heal. While there a recurrence developed on the site of the burn without the burn healing.

This case to all appearances was similar to Case 2, but the results have been very different. It is an instructive case, because of the idiosyncrasy against the rays, since less rays were used than on either of the two previous patients. It also shows that a severe burn is no proof against carcinoma; also, that it is wrong not to keep a case under observation while a burn is healing, and that even when a good result is obtained the case should be watched.

CASE 4.—Miss D. B., aged 16 years. Referred by Dr. L. Webster Fox. This was a case of retrobulbar sarcoma, the diagnosis being confirmed by microscopic examination. The antrum of Highmore had been previously curetted by Dr. E. B.

Fig. 2.—Case 4. After thirty-four treatments. July 16, 1903.

June, 1902, one month after the growth was first noticed. The whole breast and axillary glands were removed. A recurrence was first noticed in November, 1902. X-ray treatment was begun Jan. 22, 1903. At this time nodules from the size of a pea to that of half a walnut were found scattered through the field of operation. A small indurated gland could be felt in the left supraclavicular region. Twenty-six treatments were given between Jan. 22 and April 7, 1903, when a burn of the second degree had been produced. The growth had lessened very much, both in size and induration. Treatment was discontinued. The thyroid gland, which was large at the beginning, now grew rapidly and the patient developed the symptoms of exophthalmic goiter. She was taken to the Presbyterian Hospital, where the burn healed, the thyroid gland decreased rapidly in size, and the general health improved. The patient then developed pain in the axilla, and Dr. Newcomet

Fig. 3.—Case 4. After forty treatments. Aug. 27, 1903. Scar due to an operation. Sight not disturbed.

Gleason. The growth again rapidly infiltrated the sinuses. The patient had been examined by Dr. Walter Freeman. She had also been examined by Dr. W. L. Rodman, who refused to operate and advised x-ray treatment. No one expected the patient to live. At the time of beginning treatment the patient was too weak to stand, complained of severe headache and pain in the eye. There was a marked bulging of the eye from the orbit and a protrusion of the conjunctiva. The case was treated daily with five-minute exposures, with a high vacuum tube, and 3 to 4 ampères of current, at a distance of 12 inches. At the end of a month the swelling had almost disappeared, the pain was less, and the patient was well able to walk around and left the hospital. There had been an erythema present for two weeks. She was then treated twice a week for a month, at the end of which time she could open her eye slightly, and felt fairly well. She was then treated once a week for two

months, when she was apparently well, except for occasional pain in the region of the inner canthus. She is still being kept under observation and occasional treatment. There was no loss of sight.

This case is important (1) because it was a sarcoma; (2) because of its extensive infiltration; (3) because of its prompt and constant improvement; (4) because there was no injury to the organ of sight as a result of the treatment.

CASE 5.—Mrs. A. S., aged 69 years. Epithelioma of the face. Referred by Dr. J. M. Anders, July 3, 1903. During the previous two months an epithelioma had been developing on the right side of the nose, and extended on the bridge of the nose. It was completely removed by 20 exposures, covering a period of two months. The cosmetic result is perfect. There is no scar, and the skin is smooth and of a pink color.

CASE 6.—Mr. S. G., aged 60 years. Epithelioma of the mucous membrane. Referred for x-ray treatment by Dr. Hollopeter. Ten years ago a warty growth developed on the lower lip. He began treatment at a sanatorium near Boston, July, 1900, where caustics were used. He remained there 18 months, at the end of which time he came to Dr. Hollopeter and was referred to me. His mouth could then not be opened except to admit the little finger, and this gave him severe

Fig. 4.—Case 8. Tubercular ulcers of the leg, March 3, 1903, before x-ray treatment.

pain. The growth involved the right angle of the mouth and the greater part of the cheek. There was an enlarged gland in the right submaxillary region. The gland disappeared in about a month under treatment, but the growth itself, being mostly in the mouth, slowly but gradually increased in size and the amount of pain increased. The case was inoperable from the first, but the rays had little influence except perhaps to slightly retard the growth. Ninety treatments were given.

Cases 5 and 6 are important because of their differences. In Case 5 the epithelioma was entirely superficial, involved the skin, had never been tampered with by caustics, was treated early with the x-rays and yielded promptly with perfect results. In Case 6 the epithelioma was not entirely superficial, involved the mucous membrane, had been tampered with by caustics for eighteen months, x-ray treatment was begun late, after the growth had started to spread, and a long course of treatment was given with practically no results.

CASE 7.—Miss A. N., aged 20 years. Tuberculous adenitis. Referred by Dr. Ernest La Place. She had had tubercular adenitis for nine years and had been operated on by Dr. La Place three years ago, and again one year ago. She was referred for x-ray treatment Feb. 9, 1903, on account of swell-

ing of the glands on both sides of the neck and because of an indurated scar on the right side of the neck. After 10 treatments the scar and the glands were softer. After six weeks and 15 treatments the scar had almost faded and the glands were almost imperceptible. After 5 months and 34 treatments the glands were imperceptible, and the scar was soft, pink and barely noticeable.

CASE 8.—Miss J. F., aged 16 years. Referred by Dr. W. L. Rodman, March 3, 1903. Diagnosis, tuberculous adenitis and tuberculous ulcers of the legs. She was operated on by Dr. Rodman one year ago, when the glands were completely removed from the left side of the neck, from the lower portion of the right side of the neck and from the right axilla. At time of beginning treatment there was enlargement of the postcervical glands and a distinct enlargement just below the right ear. There were also three ulcers on the right leg, varying in size from one inch to two inches in diameter. She has been given 40 treatments covering a period of eighteen weeks. The two smaller ulcers on the leg healed within a month. In six weeks the largest gland had ruptured and healed. At the end of four months the enlargement of the glands had disappeared. The largest ulcer on the leg had

Fig. 5.—Case 8. Oct. 12, 1903. Ulcer healed, scar still red.

been reduced to one-third its original size, and had a healthy appearance, but needed support of the circulation. Dr. Burns then applied an elastic cast and the ulcer healed in three weeks.

The above two cases are important because they show undoubtedly a good influence of the rays on the tuberculous glands and ulcers.

CASE 9.—Miss K. B., aged 30 years. Referred by Dr. La Place, under whose care she had been for fifteen years. Diagnosis, lupus vulgaris. At the time of beginning treatment, the diseased area covered the greater part of the anterior portion of the face and the entire nose. A distinct reaction was produced in two weeks by 7 exposures. Following this reaction the tubercles were leveled to the skin and desquamation took place. She has been given 18 treatments in four months, and is practically well.

I report this case because of a very interesting and remarkable coincidence. For eight years previous to beginning treatment, the patient had been very deaf, so that it was difficult to make her hear. At the end of the course of treatment, she could hear ordinary conversation. This remarkable result was unthought of and, therefore, no accurate observation had been made before treatment; but eight years ago a specialist in

New York had examined her and pronounced her deafness incurable. This result suggests that the rays may be of value in some forms of deafness.

In conclusion, I am convinced that:

1. The *x*-rays are of undoubted value in the treatment of certain cases of both superficial and deep-seated carcinoma and tuberculosis.

2. The more a case has been tampered with, the less likely it is to yield to the influence of the *x*-rays, *e. g.*, Case 6.

3. Daily treatments carefully and properly given will produce the best results, *e. g.*, Cases 1 and 4.

4. We should try never to produce a dermatitis beyond a simple erythema.

5. There are idiosyncrasies in certain people which render them most susceptible to the *x*-rays, and in these people deeper burns may occur in spite of the most careful treatment, *e. g.*, Case 3.

6. Epithelioma involving the mucous membrane is much less likely to yield to the effect of the *x*-rays than when it simply involves the skin, *e. g.*, Case 6.

7. There is not likely to be any interference with the sense of sight, even though the *x*-rays are used directly over the eye, *e. g.*, Case 4.

8. Tuberculosis, whether of the skin or of the glands, will yield at least in certain cases to the effects of the *x*-rays, *e. g.*, Cases 7, 8 and 9.

9. The *x*-rays may be of value in certain cases of deafness.

10. The *x*-rays will give better cosmetic results than any other form of treatment in simple epithelioma of the face, *e. g.*, Case 5.

11. Epithelioma of the mucous membrane should be removed as early as possible by the knife, and this followed by *x*-ray treatment.

12. Operable cases should be operated on and in each instance followed by a course of *x*-ray treatment, and *x*-ray treatment should be given at the first sign of a recurrence.

13. Inoperable cases should be given a trial with the *x*-rays, since even hopeless cases sometimes yield to this form of treatment, *e. g.*, Case 4.

14. It is desirable to produce a distinct reaction in the treatment of lupus, *e. g.*, Case 9.

Clinical Reports.

TWO CASES OF TRANSITORY COMPLETE BLINDNESS OF BOTH EYES.*

CHARLES J. KIPP, M.D.
NEWARK, N. J.

IDIOPATHIC OPTIC NEURITIS.

CASE 1.—Mrs. K. S., aged 32, has been married eleven years, and is apparently in robust health. She has given birth to two children, both of whom are living and in good health; the younger is 4 years old. Patient has always been in good health. No history of syphilis. For some weeks she has suffered more or less from frontal headache.

She came to the Newark Eye and Ear Infirmary for the first time Dec. 22, 1901, complaining of loss of vision of left eye and impairment of vision of right eye. The examination made at this time showed that the exterior parts of the eye were perfectly normal in appearance, and that the movement of the eyes were unimpeded in any direction. On closer examination it was found that the left eye was blind; this eye could not distinguish between light and darkness. The pupil of this eye was larger than that of the right eye and it did not react

to light. It did, however, respond to light thrown in the right eye. The ophthalmoscope showed clear media and a well-marked papillitis. The optic disc was swollen and apparently enlarged. The retinal veins were full and tortuous, while the arteries were of about normal caliber. The retina immediately surrounding the disc was hazy, but otherwise it was normal. The right eye also showed swelling and enlargement of the optic disc, but not quite so marked as that of the other eye. The retina was transparent except very close to the optic disc. The pupil of the right eye was of normal size and responded readily to light. The sight of this eye was reduced to 5/12. An examination of the urine showed it to be free from albumin and sugar. Close questioning failed to reveal disease in any other part of the body.

She was given iodid of potassium in small doses. She was seen again the next day, when it was found that both eyes were now totally blind. She could not recognize the light of a candle held close to her eyes. The swelling of the optic discs had increased. Their surfaces were best seen with a +3 dioptric glass. The pupil of the right eye was now as large as that of the left eye, and neither pupil reacted to light.

She remained in this condition until December 28, when she could distinguish between light and darkness. December 30 it was found that the pupil of the right eye responded very sluggishly to light, but contracted in convergence. The pupil of the left eye responded even more sluggishly than the right eye to light, but also contracted in convergence. She can see movements of hands in front of her eyes, but can not yet count fingers close to her eyes. She was at this time taking fifteen grains of the iodid of potassium three times daily.

Jan. 8, 1892, it was found that the pupils of both eyes reacted to light in convergence fairly well. The vision of both eyes remained about the same as at last, January 14. Can now count fingers with either eye at four or five feet. She still has frontal headache. The optic discs are less swollen and the retina adjoining the disc is more transparent.

January 22: Her vision is now 5/18 in the right eye and 5/36 in the left. The visual field is intact in both eyes. She is taking twenty grains of the iodid three times daily. After this time her vision gradually improved so that on February 5 it was 5/9 in either eye. The discs were now almost flat and quite white. The headache had subsided.

March 15 her vision was about normal in both eyes and the visual field intact in both. During the following five years I saw the patient at intervals of several months and always found her in good health. Her vision when I last examined her five years after the attack was normal in both eyes and so were the visual fields. Her health was excellent. I have been trying to find her since, but have not succeeded in so doing.

The absence of the pupillary light reflex for a while made it probable that the seat of the lesion was in the visual tract, this side of the corpora quadrigemina.

The visual centers may also have been involved, but this can not be proved, as the conduction of visual impressions was interrupted by disease of the tract. The ophthalmoscope showed the existence of a marked optic neuritis in both eyes. Cases in which an optic neuritis has caused total blindness of both eyes and in which the pupillary light reflex was absent for a time, with subsequent restoration of normal vision, and very seldom encountered. With the exception of a frontal headache my patient was in excellent good health at the time the optic neuritis developed, and as she remained in good health during the next five years it seems reasonable to assume that the eye disease was not the result of a general affection which was latent when she became blind.

AMAUROSIS, PROBABLY FROM AUTOINTOXICATION.

CASE 2.—Mrs. S., aged 52, stout, of typical bilious temperament, is subject to frequent attacks of headache; is constipated most of the time; is fond of rich food and indulges freely in eating.

July 27, in the evening, she was attacked with a sudden intense pain in the head, with nausea and vomiting. The attack was so severe that her attending physician was obliged to give her ¼ grain of morphin hypodermically. She was re-

* Read at the one hundred and thirty-seventh annual meeting of the Medical Society of New Jersey.