myself. We thus used stimulants with and after our dinner meal. We all three marched to Coomassie and back; never felt sick or sorry, although as staff we had many calls on us day and night, and I am of the three the only one who has suffered, having been attacked with remittent fever on my landing in England, but this happily passed off in four days under the daily use of quinine in small doses.

(Signed) B. W. TROUP, M.B.,

Surgeon, 42nd Royal Highland Regiment
("The Black Watch").

Portsmouth, May 8th, 1874.

(To be continued.)

A CASE OF URETHRAL CALCULUS INCARCERATED BETWEEN TWO STRICTURES; EXTERNAL AND INTERNAL URETHROTOMY; GOOD RESULT.

By W. F. TEEVAN, B.A., F.R.C.S.,

SURGEON TO THE WEST LONDON HOSPITAL, SURGEON TO ST. PETER'S HOSPITAL, AND LATE LECTURER ON ANATOMY AT THE WESTMINSTER HOSPITAL.

THE following case is one of much interest, not only on account of the rarity of its occurrence, but also from the different operations required, and the fact that the calculus simulated a traumatic stricture.

William C—, a powerful, healthy-looking stoker, aged thirty-three, gave the following account of his complaint:—When only thirteen years old he was kicked in the perineum, which caused "an awful lot of blood" to come from the penis, and the testicles to swell, but no difficulty in micturition followed. He caught a gonorrhœa when twenty-one years of age, and was long troubled with a gleet; both complaints were treated by medicines only. Ten years ago he began to be troubled with difficulty in passing his water, and went into the Southampton Infirmary, where his stricture was gradually dilated up to No. 11. During the treatment a very small calculus passed from him one day, and about two years ago he withdrew another small stone in the eye of his catheter, and was attacked with rigors a few hours later. On Feb. 6th, 1874, the patient sought my advice at St. Peter's Hospital on account of the increasing difficulty in micturition and the incessant calls to urinate, which forced him to discontinue his work. With some trouble I was enabled to pass a small olivary elastic bougie into the bladder, and to determine the existence of apparently three strictures by external manipulation. One of them was situated in the perineum, another behind the scrotum, and the third in the penile urethra, just in front of the scrotum; and each was about half an inch long. Between Feb. 14th and March 14th, the patient paid me seven visits, but I was only able to pass a bougie two sizes larger than on the first occasion, and I advised him to allow me to perform internal urethrotomy, to which he readily consented; and on March 28th he entered the hospital for the purpose. He was kept quiet in bed, and the perineum was well leached, which enabled me to dilate the first stricture sufficiently to permit me to divide it with Civiale's instrument, on April 8th, in the presence of Mr. Heycock, &c. The patient, who stood up against the wall when the operation was performed, said he suffered but little pain. Only a few drops of blood escaped, and no rigors ensued from the passage of the urine over the freshly divided mucous membrane. I now tried to dilate the second stricture, but without success, and, on May 2nd, I introduced a fine whalebone bougie into the bladder, and used it for a guide for the catheter urethrotome described by me at p. 724 of THE LANCET for May 23rd. Before passing the knife into the urethra, I moved the supposed stricture up and down on the staff, and thus discovered that it was a small calculus, although it had been looked upon as a well-marked specimen of traumatic stricture by those who, like myself, had hitherto examined it externally, without the assistance of a metal instrument in the urethra. I at once passed a urethral forceps and grasped the stone, and managed to bring it forward to about half an inch in front of the scrotum. As I was unable to withdraw the calculus, I introduced Nélaton's urethral "brise-pierre," but I could not get the male blade home, and tried without success to extract the stone by means of the beak of the female blade. I then gave the

instrument to Mr. Heycock to hold, whilst I slid the skin away from the raphe one inch to the right, and removed the calculus through a small incision in the median line.* The skin was now allowed to roll back to its normal position, so that the external incision appeared to be at the side of the penis, and not in the median line where it had really been made. No instrument was left in the urethra, and no rigors followed. Two days later I passed one of the smallest elastic catheters into the bladder, and tied it in for twelve hours, in order to rapidly dilate the deep stricture. On May 6th, four days after the operation, the wound was quite healed. The patient left the hospital on May 23rd, passing No. 14 French catheter, with ease, for himself. His stream of urine was of normal size, and he could hold his water all night, and for four hours in the day. He called on me on July 28th, when he passed No. 16 in my presence. As the deep stricture is very tough, and the introduction of a full-sized catheter a procedure of some difficulty attended with pain, I have recommended the patient to allow me to divide it when he can spare the time to enter the hospital again.

The case presents some important clinical features. When the patient first came to me no metal instrument could be passed, but a small whalebone bougie could be introduced into the bladder with some difficulty, and, although it felt tight, it gave no evidence of the presence of a calculus. Even when the first contraction had been divided no stiff catheter could be insinuated through the second obstruction, and had it not been for the great assistance afforded by the whalebone bougie in enabling the catheter urethrotome to be slipped over it the true character of the apparent second stricture would not have been cleared up, as the calculus was embedded in a recess, and when a metal instrument was passed down to it no click was elicited. With all the facts of the case now before me, I would say that the anterior stricture was, from its nature, the result of the long-standing gleet from which the patient had suffered; whilst the posterior one in the perineum was caused by the kick. How was it, then, that the calculus was arrested by the more recently formed stricture instead of by the older one? The answer is not difficult, I think. Calculi are usually stopped in their descent in the anterior portion of the canal simply because it is much narrower than the deeper portion. When the stone had advanced as far as the front stricture the deeper one still went on contracting, so that it could neither come forward nor recede. The calculus was extracted by an external vulvar incision, and the wound was healed in four days. Now, one of the objections to the performance of external urethrotomy in the penile urethra is the alleged probability of the formation of a fistula, but this may be entirely obviated by adopting the manœuvre I have described. The skin of the penis is so loosely connected with the fibrous envelope of the organ that it can be easily shifted to a great extent in any direction. Thus it was that, although the cicatrix is to be seen one inch to the left of the raphe, it was in reality made in the median line so far as all the tissues, save the skin, were concerned. The case also demonstrates a surgical point, now generally accepted, to the effect that no instrument need be left in the urethra after internal or external incision, for neither rigors nor abscess occurred after the operations, although no catheter was left in on either occasion.

Portman-square, W.

EPILEPTIC HEMIPLEGIA.

By J. CRICHTON BROWNE, M.D., F.R.S.E.,

MEDICAL DIRECTOR, WEST RIDING ASYLUM.

UNDER the title of epileptic hemiplegia Dr. Todd described a number of cases in which transient paralysis of one half of the body followed convulsions of the epileptic kind, generally limited to the side which became paralysed, but sometimes involving the whole body. Combating the crude and easy pathology which would ascribe such hemiplegia, as well as all other obscure brain symptoms, to local congestion, he argued that it ought to be attributed to a state of exhaus-

* The stone was phosphatic, olive-shaped, and weighed, when dried, twelve grains. It measured three-quarters of an inch long, and just filled No. 23 hole of the French gauge.

tion and weakened nutrition in the deeper strata of the white matter of the hemispheres, in the corpora striata, and optic thalami, induced by the morbidly excited polarity of these and other parts, upon which he believed the epileptic paroxysm to depend.

Dr. Hughlings Jackson, in reviewing the clinical facts as to the association of paralysis with convulsions, so accurately observed by Todd, has offered another explanation of them, suggesting that the temporary loss of power is caused by the strong discharge of the cerebrum in which the fit consists. Numerous cases verifying by their phenomena Dr. Hughlings Jackson's theory are continually falling within the experience of those who see much of nervous diseases, but a certain number of cases are encountered for which it does not, at first sight, adequately account. He has himself referred to and dealt with one class of such cases, in which the paralysis precedes the convulsions; and there is another class, well worthy of attention, in which the paralysis follows the convulsions, but is not temporary but gradually progressive, advancing at a rate determined by the number and severity of the convulsive seizures. In this class of cases Dr. Todd thought that weakened nutrition had been continued into a state of white softening. There was a tendency, he believed, in the fit-exhausted brain towards that variety of degenerative change which, when it involved the convolutions, compromised mental power and induced dementia, and when it involved the ganglia at the base of the encephalon compromised muscular power and induced paralysis. Unfortunately for this view, however, post-mortem examinations, in some cases in which progressive hemiplegia has resulted from epilepsy, have not discovered any softening, but rather an opposite condition of induration, affecting both convolutions and ganglia. There has been no atrophy in these cases, for the brain has looked as if increased in bulk, swelling out when the skull was removed; there has been no excess of serous effusion, no vestige of clot, no tumour; and the difficulty has been to find something to account for the hemiplegia. The fibroid induration which Dr. Bucknill's remarkable investigations first taught us to recognise in epileptics, and which was present in such cases, has seemed to be generally diffused throughout both hemispheres, and has not therefore appeared available as an explanation of palsy of one side, until some recent experiments of my own have shown that appearances, as regards this pathological change, are deceptive, that it is not so generally and equally diffused as has been supposed, and that it may affect the two hemispheres in very different degrees. One striking case, which occurred in my practice some time ago, illustrated very clearly the possible dependence of hemiplegia upon a hyperplastic overgrowth of connective tissue in the brain chiefly involving one hemisphere, and afforded a clue to the relation subsisting between the discharging and the destroying lesion when epilepsy eventuates in paralysis. It presents so many interesting features as to justify a full report.

Mary N—, aged thirty-three, single, needlewoman, from Leeds, was admitted to the West Riding Asylum Feb. 10th, 1871. She was said to belong to a healthy family, untainted by nervous disease. Her mother died of old age, in her seventy-eighth year, and her father of emphysema when aged sixty-six. She had never sustained any injury to the head, nor suffered from any serious illness, having been indeed exceptionally healthy up till her nineteenth year, when she was suddenly attacked by epilepsy, for which her friends could suggest no more probable cause than the sedentary habits necessitated by her occupation. From her nineteenth up to her thirty-third year fits of considerable severity occurred from time to time, and gradually, in spite of various methods of treatment adopted, became more frequent, and sapped her mental powers. For twelve months previous to her admission to the asylum she had had one or two fits every second day, and had become stupid and quarrelsome, without, however, exhibiting any distinct exacerbations of excitement. When admitted, she was described as rather below the middle height, of dusky, sallow complexion, and plump and well nourished. A conversation with her proved that her memory was greatly circumscribed, that her power of apprehension was defective, and that all her mental processes were slow and laborious. Her muscular

movements were seen to be clumsy and uncertain, and it was remarked that she was right-handed, but by no means dexterous in either writing or sewing. Both pupils were dilated, and there was a growth of hair on the chin and upper lip. The heart-sounds were feeble, the lungs healthy, and, according to her own statement, subsequently confirmed, the catamenia regular.

The first fit in the asylum occurred on Feb. 12th, two days after admission. There was no warning of its approach, but on its incursion the patient raised both hands behind her head, screamed loudly, fell over, and was violently convulsed, principally on the left side. Her head was drawn to the left, and clonic spasms affected the face, neck, left arm, and left leg, spreading subsequently and in a less degree to the right arm and leg. During the seizure the tongue was bitten, and after its cessation there was a prolonged period of unconsciousness, followed by confusion and bewilderment of thought lasting some hours. All the numerous fits through which Mary N— afterwards passed were nearly identical in character with that just described. In all of them the clonic spasms chiefly affected the left arm and leg, which were soon found to be decidedly weaker than the limbs of the opposite side. The grasping power of the left hand was less than that of the right to a very perceptible extent, and the left leg was dragged heavily in walking. This weakness of the left side gradually increased until there was drooping of the left shoulder, and a distinct halting gait. Immediately after her admission to the asylum, indeed on the 12th February, the patient was put under the bromide of potassium treatment, and was ordered to take thirty grains of that salt thrice daily. For a time this remedy seemed to have but little effect, as the fits recurred with their customary frequency, though perhaps with scarcely their customary violence. Up till the end of February there were ten fits, during March there were eighteen, and during April eighteen. Then, however, a striking improvement was inaugurated. In May there were only six fits, and in June and July there were none. A continued immunity from fits, accompanied by augmented mental power and nimbleness, was enjoyed up till Aug. 21st, when, however, the pent-up torrents began to overflow. On that day there were two fits, and on the 22nd, 25th, 27th, 28th, and 30th there were also two. On Sept. 1st there was one fit, and then came a temporary lull until the 8th, when the genuine eruption commenced with three fits, followed on the 9th by five, on the 10th by ten, on the 11th by twelve, on the 12th by thirteen, and on the 13th by fifteen, causing continuous unconsciousness, and placing the patient, indeed, in the status epilepticus. That was the climax of the attack, and from that point subsidence commenced. On the 14th there were no fits, on the 15th ten, on the 16th eight, on the 18th nine, on the 19th six, and so on by gradual stages of diminution until the 30th, when only one fit occurred, and when the patient seemed to be nearly restored to her usual condition. In October there were no fits, and in November there was only one comparatively trifling explosion, extending from the 10th to the 14th, and including fifteen fits distributed over these four days. During the early part of December there were no fits, but on the 17th of that month the final explosion began. On that day there were six very severe fits, on the 18th there were two, on the 19th one, on the 20th four, and on the 21st twenty. A state of stupor had been induced by the fast-coming fits. The patient could not understand what was said to her, but remained in a drowsy, half-conscious state, with saliva trickling from her mouth. Stimulants failed to rouse her, and after fifteen more fits on the 22nd, she again passed into the status epilepticus, marked by deep coma, hurried stertorous breathing, livid features, upturned injected eyes, fixed dilated pupils, profuse perspiration, and a small feeble pulse, beating 120 times per minute. During the 23rd there were no fits, but the coma continued unmitigated, and exhaustion was obviously setting in, notwithstanding the copious supplies of nourishment which were being administered by the stomach-pump and by injections. On the evening of that day it was thought advisable to resort to dry cupping, as ice to the spine, sinapisms, ergot, &c., had failed; but during the operation a fit came on, and in the fit breathing ceased, and artificial respiration had to be kept up for some time before breathing was re-established. During the 24th there were fifteen fits, with ever-increasing exhaustion. On the 25th her pulse was almost imperceptible, her bronchi were loaded with mucus,

and her face and neck were of a dull-purple colour. She remained deeply unconscious, and died on the morning of the 26th, at 1.30, the convulsive movements up to the last having been always most marked on the left side, which continued to be partially paralysed.

(To be concluded.)

CASE OF INTUSSUSCEPTION IN AN INFANT, ENDING FATALLY.

By CHARLES J. CULLINGWORTH, M.R.C.S.,

SURGEON TO ST. MARY'S HOSPITAL FOR WOMEN AND CHILDREN, MANCHESTER.

ON the evening of the 27th July, 1874, Mr. and Mrs. C— left Blackpool by train for their home in Manchester, bringing with them their baby, a fine healthy boy, fifteen weeks old. During their fortnight's sojourn at the seaside the child had been in his usual health, except that his stools had been somewhat curdy and of a greenish colour, which the mother attributed to the change of milk. Half an hour from the time of starting, the baby awoke from his sleep, and vomited rather violently. Immediately after the attack his mother called the father's attention to a sudden change which had taken place in his appearance: he looked sunken and pale, "like death." During the remainder of the journey, which occupied two hours, he was restless and fretful, uttering, almost without cessation, an unusually weak, piteous cry. I saw him very soon after his arrival at home, and found him still collapsed, the surface was blanched, cold, and perspiring; the features sunk, and the pulse feeble. He closed his eyes for a few seconds, and then roused himself suddenly, as though in pain, with the weak, painful cry which his parents had before noticed. I told them that I feared a sudden displacement of the bowel had occurred. He was immediately wrapped in a small blanket wrung out of hot water, and then enveloped in a dry outer blanket and laid upon a couch. A teaspoonful of milk, to which a drop or two of brandy had been added, was ordered to be given every ten minutes. He died, however, without rallying, at 5.30 A.M. on the 28th, not quite nine hours after the symptoms were first observed.

An examination of the body, twelve hours after death, revealed an invagination of the ileum, an inch and a quarter in length, situated at about its central portion, and lying deeply and posteriorly on the right side of the abdomen. The displacement was from above downwards. There were no external signs of inflammation, either in the part itself or in its neighbourhood. Wishful to preserve the specimen, I left the parts undisturbed, except that the inner fold was withdrawn (with the exercise of some force) to the extent of two or three lines, when it was observed to be distinctly congested, especially at its upper margin, where there had been some constriction. The stomach and upper part of the small intestine were a good deal distended with flatus, but were otherwise healthy. The canal had not been occluded by the displacement; for brandy and milk were found, apparently unchanged, in the parts below, and, indeed, the father assured me that a little had passed per anum.

MEETING OF PUBLIC ANALYSTS AT CANNON-STREET HOTEL.

A NUMEROUSLY attended meeting of public analysts took place at Cannon-street Hotel on Friday, August 7th. The chair was taken at a little after 4 o'clock, by Prof. Redwood, who opened the proceedings in a sensible and temperate speech. He pointed out that the meeting had been convened for the purpose of considering the Report of the Select Committee on the Adulteration of Food Act (1872), and of suggesting amendments to that Act. A printed draft of resolutions to be submitted to the meeting had been sent round some days before, and to this the chairman referred. These resolutions were then put to the meeting *seriatim*.

Many verbal amendments were inserted, and some fresh resolutions were added, and ultimately, after full discussion, the great majority were carried almost unanimously.

The first discussion arose in regard to the proposed reference to Somerset-house Laboratory in disputed cases. Dr. Dupré pointed out that the certificate of a public analyst had very seldom been proved to be wrong, and that magistrates very rarely refused permission to have a second analysis made in the interest of the tradesman. If a court of referees must be established it would at any rate be necessary that the actual analyst should be present at the adjourned trial, and should give his evidence on oath. Mr. Wanklyn took a similar view, asserting that a reference to Somerset House would be a reference from a higher to a lower authority. He argued that only one court of reference would be desirable—namely, a court consisting of the public analysts themselves. Mr. W. L. Scott, Mr. Allen, and Mr. Blyth also supported, in the main, this view of the case.

The third resolution had reference to the proposed examination, in doubtful cases, of candidates for the post of public analyst. Universal objection was expressed to the proposed selection of South Kensington as an examining centre, but it seemed to be generally admitted that some very loose appointments had been made, and that cases must from time to time occur in which some definite test should be enforced. Dr. Tripe was warmly supported in his suggestion, that any such test should not be retrospective in its action, while Mr. Heisch proposed that a certificate of having worked two years in the laboratory of any competent analyst should be accepted as a qualification. Mr. Bell urged that if a board of examiners were established, it should consist entirely of public analysts. It was ultimately settled that either a sufficient probation in an analytical laboratory or the passing of a suitable examination should be esteemed a qualification.

The vexed question of the facing of tea followed, and produced a lively discussion. Mr. Scott, Mr. Jones, Dr. Dupré, Mr. Burge, and Dr. Tripe objected to facing as an adulteration, and as such to any legalisation of it. On the other hand, Mr. Heisch, Mr. Wanklyn, Dr. Stevenson, Mr. Allen, and Mr. Estcourt urged that the facing of green tea, as usually practised, did not materially affect the quality or the value of the tea, and, at any rate, that it would be undesirable or even impossible to arrest the practice suddenly.

Mr. Wanklyn suggested that the ash should not be permitted to rise above 7 per cent., equal to about 2 per cent. of facing, and this suggestion was ultimately carried by a narrow majority.

The next resolution dealt with the question of mixtures. After one or two verbal alterations had been discussed, it was decided that such mixtures should be labeled, and that the percentage of the ingredients, under the name of which the mixture is sold, should be stated prominently on the label, any excess of foreign matter being considered an adulteration.

One or two resolutions referring to the duties of inspectors were then carried with little opposition, and finally an Association of Public Analysts was formally established, with Professor Redwood for president, and a committee consisting, as we understood, of Messrs. Allen, Bernays, Dupré, Escourt, Hassall, Heisch, Stevenson, Wanklyn, and Wigner. Another meeting is to be held before long, and in the meantime the committee has undertaken the difficult task of preparing a definition of the term "adulteration," and some resolutions to be founded upon it.

We abstain from any comment in this place on the conclusions reached by the new Association. Some of them we cordially endorse, while from others we differ; but we congratulate the public analysts of England on the action they have taken, and wish them every success in their enterprise.

DONATIONS, &c., TO MEDICAL CHARITIES.—The executors of the will of Mr. John Bagshawe, of Tideswell, have paid £500 to the Devonshire Hospital, Buxton, and the General Infirmary, Sheffield, respectively. The Wolverhampton and Staffordshire General Hospital has become entitled to £500 under the will of Miss Ann Stubbs, of Walsall. The Kent and Canterbury Hospital has received £462 10s. from "J. M., on behalf of a friend."