INSECTICIDES.

By H. M. LEFROY.

(Imperial College of Science, South Kensington.)

It would be interesting to collect information as to the insecticides in general use in horticulture in different countries in any one season and to determine as far as possible what were the circumstances leading to their use. There is, I believe, a fashion in insecticides: at one time Paris Green and London Purple ruled; then came Lead Arseniate in America; this is apparently giving place to Zinc Arseniate and Barium Arseniate, while there is reason to hope that Lead Arseniate, and perhaps Lead Chromate, will be more familiar in this country.

So, too, for contact poisons. I remember the era in America of paraffin emulsions and rosin washes; then came whale oil soap; lime salt and sulphur followed, then the heavy oils and then "miscible oils." The vogue of the lysol and similar mixtures on the Continent has to be noted, and we may note that miscible oils are being used more freely in Australia and South Africa.

In estimating the causes of these successive fashions one would need to distinguish the effect of the recommendations of State Departments of Entomology and the influence, increasing in recent years, of the weight of pure advertising by firms interested in the sale of the particular products they were able to handle. In this paper I omit entirely the "patent" or "proprietary" insecticides in so large use, since it is not always clear on what ingredients their action depends, and I confine myself to insecticides of definite known constitution, or those which depend upon the presence of some known chemical compound.

It is not easy to ascertain exactly what insecticides are at present used on a large scale in this country, but I have tried to do so from three sources: the first is the published recommendations of horticultural journals; the second is the recommendations in the leaflets of the Board of Agriculture; the third is those that I gather from general

information, observation, reports, papers in actual use. For the first I attach extracts from 16 consecutively weekly issues of the *Gardeners' Chronicle* in 1913, these occurring in the contributions of head gardeners or in editorial advice. I think they reflect accurately current ideas and practice.

- Hardy Fruit Garden .. The tree syringed with a very mild insecticide. In
For aphides some cases it will be found necessary to use tobacco
powder, which should be left on for two or three days,
and then thoroughly syringed off with clear soft water.
p. 268, 26. 4. 13.
- Abies* diseased Spray three times at intervals of four days, with a
Aphis. Editorial solution of soft soap and quassia chips, and repeat
the spraying in September. p. 279, 26. 4. 13.
- Broad beans Syringe the plants frequently with a fairly strong solution
Black fly of soft soap. p. 280, 26. 4. 13.
- Cool orchard house .. The eradication of insect pests can be done either by
fumigation with nicotine, or by spraying with some
good insecticide. p. 288, 3. 5. 13.
- Market Fruit Garden .. For some unknown reason, insect pests are troubling me
Insect pests at present much less than usually. It is many years
since I found so few Apple suckers. Most varieties
of Apples in my orchards are almost free from this
enemy, and many are nearly exempt from the attacks
of aphides and caterpillars. It would be interesting
to learn whether this comparative immunity is general
or not. If it is not so, perhaps persistence in des-
troying the pests for years past is meeting with a fitting
reward. But it may be that suckers and aphides
hatched prematurely during the mild winter or in
March were killed by frost. Or, again, spraying with
a strong solution of lime-sulphur just before the buds
burst may have done something in coating over eggs
or destroying mother queen aphides. It is too early,
however, to assume that the comparative immunity
will continue. There can hardly be a bad attack of
the sucker after this; but the aphis and the caterpillar
have plenty of time during which to appear in strong
force. Already the hope that Plums would escape the
usual infestation of aphis has been disappointed. On
April 24 no attack could be observed on any but two
varieties; but on the 29th there were aphides on all.
As the leaves are curled over the pest, spraying would
be a mere waste of time and money.
- The first summer spraying Owing to the partial immunity of Apples from insect
pests, the usual spraying before the expansion of

blossom was to a great extent dispensed with, only varieties which were badly attacked by aphid, sucker, or caterpillar, or all three pests being sprayed. This was a great relief, and it is not by any means certain that it might not have been extended to all varieties without any bad results. For on every occasion rain fell a few hours after the spraying was done, probably washing most of the spray fluid off the trees, and off the insects that had been wetted by it. There was no question of waiting for settled fine weather, as the work had to be done before the expansion of the blossom. Arsenate of lead was combined with the soft-soap wash for aphid and sucker, in order to poison the food of caterpillars. If pests should attack the varieties not sprayed, they will be treated after the fall of the blossom, and in any case varieties subject to scab will be sprayed with lime-sulphur of summer strength.

Results of spraying

One of the most unsatisfactory features of spraying is the difficulty of determining its results, and that difficulty has proved particularly great this season. Even in the cases of Apples most infested, such as those in which five or six out of twenty trusses of blossom were found to contain aphides, and three or four a few suckers, I could not find more than one dead aphid or sucker in fifty trusses examined a day or two days after the spraying. I found more live ones, but very few even of these. The question, then, is whether the pests, killed or weakened by the wash, were washed out of the trusses by heavy rain. It is commonly assumed that aphides are washed off the trees by rain, and particularly by a thunderstorm; but, so far as I know, this is an assumption which has never been actually proved. Some trusses found to be infested by aphides could be labelled, some of them being sprayed and others not sprayed, care being taken to avoid any more thorough spraying than would be given on a large scale. If the insects could be counted beforehand, so much the better. Examination could be made 24 hours after the spraying, results being noted; and again after the next fall of rain, if it occurred soon enough. Rain might hold off too long, and the experiment would then have to be repeated. In noting the results after rain, if any pests were found to have been washed off the trees, it should be noticed whether more were ousted from the sprayed trees than from the others. As to a thunderstorm, there can hardly be any special

effect from it, unless the rainfall happens to be more violent than in an ordinary storm, as it is not at all likely that a specially electrical condition of the atmosphere affects the aphid. A busy fruitgrower can hardly find time for such experiments as are suggested. They should be carried out by the authorities paid out of public funds to undertake research. As matters stand, I do not know whether the expense incurred in spraying has been useful or almost useless. There is too much reason to fear that the arsenate of lead has been all, or nearly all, washed off the trees by rain, and that the caterpillars will be able to go on feeding with impunity. They, like the other pests, were in the trusses of blossom buds and the surrounding leaves. "A Southern Grower." p. 301, 10. 5. 13.

Roses	Should the pest appear on the leaves, syringe the trees with a mixture of soft soap and sulphur. The sulphur will also check mildew. Caterpillars are particularly numerous this season and unless they are diligently sought for they may do much damage. p. 308, 10. 5. 13.
Aphis	
Hardy Fruit Garden ..	If the pest has already taken a hold on the bushes, they should be syringed at once with a good insecticide. Some mixture containing a certain proportion of nicotine will answer the purpose best, it will kill the grub in a few minutes. The trees should be afterwards syringed with clean, soft water. p. 309, 10. 5. 13.
Gooseberry caterpillar	
Editorial	Place freshly tarred paper under the vines, then approach the vines with a lantern in the hand, giving each rod a slight shaking. . . . It is difficult to kill the larvae in the soil for they seem proof against any but the strongest and most dangerous applications. p. 320, 10. 5. 13.
Vine weevils	
Fruits under glass ..	The vines will need syringing occasionally with an insecticide to destroy red spider. p. 326, 17. 5. 13.
Red spider	
Hardy Fruit Garden ..	A small yellow caterpillar is a great pest: this pest should be hand-picked and destroyed. After a thorough search syringe the trees with a weak solution of nicotine. p. 327, 17. 5. 13.
Apricot	
Market Fruit Garden ..	Our trees were sprayed with Buxton lime and in the case of Warner's King, which I have always found the most susceptible to sucker, I used salt with the lime. The spraying was done just before the flower buds parted from the cluster ready for expansion. p. 332, 17. 5. 13.
Apple sucker	
Editorial	The trees should have been sprayed with arsenate of lead. . . . as soon as the petals have fallen, spray with 1 lb. of arsenate of lead paste to 25 gallons of water. As
Apple blossom	
Bud moth	

effect from it, unless the rainfall happens to be more violent than in an ordinary storm, as it is not at all likely that a specially electrical condition of the atmosphere affects the aphid. A busy fruitgrower can hardly find time for such experiments as are suggested. They should be carried out by the authorities paid out of public funds to undertake research. As matters stand, I do not know whether the expense incurred in spraying has been useful or almost useless. There is too much reason to fear that the arsenate of lead has been all, or nearly all, washed off the trees by rain, and that the caterpillars will be able to go on feeding with impunity. They, like the other pests, were in the trusses of blossom buds and the surrounding leaves. "A Southern Grower." p. 301, 10. 5. 13.

Roses	Should the pest appear on the leaves, syringe the trees with a mixture of soft soap and sulphur. The sulphur will also check mildew. Caterpillars are particularly numerous this season and unless they are diligently sought for they may do much damage. p. 308, 10. 5. 13.
Aphis	
Hardy Fruit Garden ..	If the pest has already taken a hold on the bushes, they should be syringed at once with a good insecticide. Some mixture containing a certain proportion of nicotine will answer the purpose best, it will kill the grub in a few minutes. The trees should be afterwards syringed with clean, soft water. p. 309, 10. 5. 13.
Gooseberry caterpillar	
Editorial	Place freshly tarred paper under the vines, then approach the vines with a lantern in the hand, giving each rod a slight shaking. . . It is difficult to kill the larvae in the soil for they seem proof against any but the strongest and most dangerous applications. p. 320, 10. 5. 13.
Vine weevils	
Fruits under glass ..	The vines will need syringing occasionally with an insecticide to destroy red spider. p. 326, 17. 5. 13.
Red spider	
Hardy Fruit Garden ..	A small yellow caterpillar is a great pest: this pest should be hand-picked and destroyed. After a thorough search syringe the trees with a weak solution of nicotine. p. 327, 17. 5. 13.
Apricot	
Market Fruit Garden ..	Our trees were sprayed with Buxton lime and in the case of Warner's King, which I have always found the most susceptible to sucker, I used salt with the lime. The spraying was done just before the flower buds parted from the cluster ready for expansion. p. 332, 17. 5. 13.
Apple sucker	
Editorial	The trees should have been sprayed with arsenate of lead. . . as soon as the petals have fallen, spray with 1 lb. of arsenate of lead paste to 25 gallons of water. As
Apple blossom	
Bud moth	

- there are some aphides in the trusses add 2 lb. of soft soap to the 25 gallons of wash. p. 336, 17. 5. 13.
- Editorial Spray in winter with a wash of 3 lbs. soft soap, 5 gallons water, and 1 pint of paraffin thoroughly mixed. p. 336, 17. 5. 13.
- ‘Spruce gall’
- Hardy Fruit Garden .. Black and green aphides will soon make an appearance on the ends of the shoots and should be destroyed by syringing with insecticide. Leaves that are infested with caterpillars or (? of) the Pear moth, should be gathered and burned. p. 367, 31. 5. 13.
- Pear trees on walls
- Editorial To destroy aphis, it is best to fumigate with a nicotine preparation. For red spider, the leaves may be sponged with a little soapy water—half an ounce of soft soap to one gallon of water—mixed with half a pound of flowers of sulphur and kept stirred. Or flowers of sulphur may be applied dry with the Malbec bellows. p. 375, 31. 5. 13.
- Aphis and red spider on vines
- Roses Syringe roses with an insecticide at least once a week to keep aphis in check. p. 383, 7. 6. 13.
- The flower garden
- Fruits under glass .. The trees should be syringed frequently with an insecticide. p. 383, 7. 6. 13.
- Red spider
- Public parks and gardens The Goat moth (*Cossus ligniperda*) is far more common over the London area than is generally supposed, and this may also be said of the Wood Leopard moth (*Zeuzera aesculi*), both of which attack the Elm, Poplar, Ash and Oak. The larvae of both moths, which are deposited on the bark of the tree in July or August, tunnel into the wood and do much damage to the timber as well as the health of the tree. Placing cyanide of potassium in the hole and closing the aperture is the best method of destroying the caterpillar, though we have used gas tar in a similar way with good results. Dislodging the caterpillar by means of a bent wire has been successfully carried out. Lime trees all over London are attacked by the caterpillar of the Lime Looper moth (*Hybernia defoliaria*), while the Thorn fly, an Aphis, attacks whole hedges of the Hawthorn as well as specimens standing singly. Sponging with tobacco water is to be recommended for the latter, and for the Lime trees caterpillar bands of any sticky substance painted around the stems will prevent the insects ascending to the branches. Red spider (*Tetranychus telarius*) is another formidable insect pest that not infrequently attacks the leaves of several species of hardwooded trees, and is often the cause of death of the Ivy, particularly when grown as a ground carpet. The leaves turn a rusty-brown colour, crumple

up, and finally fall off, the whole plant dying in consequence. Large areas of the Ivy used for ground work in various parks and gardens of London have been killed outright of late years. Spraying with soft soap will prevent the insect from spreading. Though the Beech is by no means a common tree in London, yet of the few specimens to be found, many are attacked by *Cryptococcus fagi*, which is alarmingly on the increase in this country. Paraffin or petroleum emulsion will destroy the insect, and scrubbing the bark with a rough brush and soft soap is an excellent remedy. Wholesale destruction to the leaves of the Spindle tree and other species of *Euonymus* is yearly occasioned in some of the London parks by the caterpillar of the small Ermine moth, myriads of the caterpillars appearing on the foliage during summer. So rapidly does this insect increase, and so voracious is its appetite, that a shrub will be completely stripped of its foliage in two or three days. Spraying with XI. All Insecticide or petroleum emulsion is the best remedy. Where Oak trees are found in London, the Oak-leaf Roller moth (*Tortrix viridana*) is usually present about the beginning of June. It is a very destructive insect, and attacks not only the leaves, but buds and inflorescences, and usually works from the top of the tree downwards. Where only a single or few Oaks are attacked spraying may be resorted to, but hardly any remedial measure can be adopted in a clump or plantation of the tree. Starlings, rooks and other birds destroy vast numbers of this pest, and should be encouraged. By "Superintendent." pp. 400, 401, 14. 6. 13.

- Apples and plums .. Spraying lime-wash just before the buds approach the bursting stage in dry weather.
- An important object lesson Very striking evidence as to the advantage of spraying Apples and Plums with lime-wash, as thick as it can be got through the spraying machines, was obtained in Mr Hooper's orchards. In one portion of the farm all the Apple and Plum trees had been sprayed with this wash, nothing being added to the lime, except water, while they were left unsprayed in the rest of it. The lime-sprayed Apples and Plums were free from the aphid blight, while those not sprayed were very badly infested. If memory serves, the contrast applies also to Apple sucker infestation. At any rate, a mere glance was sufficient to allow any one to see it in relation to the aphid blight, and growers elsewhere to

whom this striking case was named said that they had seen similar results from lime spraying. Mr Hooper's explanation is that the lime coats over the eggs of the pests, and extracts moisture from them, thus destroying their vitality. Lime-wash must be strained before it is put into a spraying machine, and it is important to do the work only in dry weather, and to cover every twig. If rain occurs before the stuff has dried thoroughly on the trees, the lime is easily washed off; but it sticks on well after being dried. In the case under notice the spraying was done as soon as there was any sign of an attack by birds upon Plum buds. Apart from this attack, however, just before the buds approach the bursting stage is probably the best time to apply the lime-wash. "*Southern Grower*." p. 414, 21. 6. 13.

- Flower Garden. Aphides on chrysanthemums .. It is wise to spray with some insecticide of moderate strength. p. 436, 28. 6. 13.
- Hardy Fruit Garden .. Red spider, syringe the foliage constantly with water, to which has been added a little quassia extract.
Wall trees
If the cherries are infested with black fly, the affected shoots should be cut away and the trees syringed with an insecticide. After each application a syringing of clean water should follow. This method will be found to be a great improvement on the old plan of applying tobacco powder. p. 437, 28. 6. 13.
- Editorial Soft soap, one quart, in one gallon of boiling water. Add one pint crude carbolic acid and dilute with 30 times its bulk of water. Syringe the ground around the attacked plants. p. 444, 28. 6. 13.
- The spruce aphid .. Spraying with paraffin emulsion is recommended. Soft soap and quassia extract may also be tried. p. 5, 5. 7. 13.
- Hardy Fruit Garden .. If there are bad infestations the growths most affected
Black aphid .. should be cut off and burned and the tree syringed afterwards, first with an insecticide, then with clean water. p. 9, 5. 7. 13.
- Editorial Next winter drench the trees thoroughly with soft soap
Conifer coccus .. and paraffin emulsion. p. 20, 5. 7. 13.
- Fruits under glass .. The foliage should be syringed thoroughly or washed by
Peaches .. means of the garden engine two or three times a week. This will cleanse the leaves of insect pests. p. 29, 12. 7. 13.
- Editorial Pinch the maggots in the leaves and remove the leaves.
Celery fly .. .Afterwards you can make the plants distasteful to the flies by syringing them with quassia extract, or you may scatter soot, lime, etc. p. 60, 19. 7. 13.

- Editorial Remove the plants and spray the exposed roots and soil
 Ripersia on palm roots with carbon bi-sulphide, using a small glass spraying
 apparatus. The process must be repeated. Another
 plan is to have holes in the soil and pour one tea-
 spoonful of the carbon bi-sulphide in each hole. p. 76,
 26. 7. 13.
- Editorial. Rose leaves 'Extract of Quassia.' p. 76, 26. 7. 13.

Board of Agriculture Leaflets.

Mangold fly	paraffin	1 gallon	} Formula 1.
	soft soap	$\frac{1}{2}$ lb.	
	water	10 gallons	
Onion fly	paraffin	3 pints	} Formula 2
	soft soap	$\frac{1}{2}$ lb.	
	water	7 gallons	
Surface caterpillars	soot and lime		
	{ Paris green	1 lb.	
	{ water	50 gallons	
Celery fly	{ paraffin	1 quart	} Formula 3.
	{ soft soap	$\frac{1}{2}$ lb.	
	{ water	10 gallons	
	{ carbolic acid	1 pint	
	{ soft soap	$\frac{1}{2}$ lb.	
Carrot fly	{ water	10 gallons	
	paraffin	2 gallons	} Formula 4.
	soft soap	$\frac{1}{2}$ lb.	
	water	60 gallons	
Red spider	sulphur		
	paraffin	7 gallons	} Formula 5.
	soft soap	6 lbs.	
	liver of sulphur	$2\frac{1}{2}$ lbs.	
	water	100 gallons	
	sulphate of iron	8 ozs.	} Formula 6.
	caustic lime	4 ozs.	
	paraffin (solar)	16 to 24 ozs.	
	water	10 gallons	
	liver of sulphur mixed with quassia wash		
Asparagus beetle	{ Paris green	1 lb.	
	{ water	200 gallons	
	{ arsenate of lead	3 ozs.	
	{ treacle	1 lb.	
	{ water	10 to 12 gallons	
Thrips	{ pyrethrum	1 oz.	
	{ soft soap	1 oz.	
	{ water	2 gallons	
Hop aphid	soft soap	4 to 10 lbs.	
	water	100 gallons	
	or with 6 lbs. quassia chips		

Aphides	..	..	..	{	soft soap	6 to 10 lbs.	}	On dormant trees (Woburn formula).
					water	100 gallons		
					or with quassia 6-8 lbs. added			
					paraffin	2 gallons		
					soft soap	1½ lbs.		
Cabbage fly	..	..	..	{	caustic soda	6 lbs.	}	
					water	28 gallons		
					paraffin	1 cupfull		
					sand	1 pailfull		
					lime	1 lb.		
Black currant gall mite	..	..	..	{	sulphur	1 lb.	}	
					water	20 gallons		
					soft soap	1 quart		
Vine weevils	..	..	..	{	crude carbolic	1 pint	}	
					water	30 gallons		
					Paris green	1 lb.		
Winter moth	..	..	..	{	water	200 to 250 gallons	}	
					arsenate of lead paste	2 lbs.		
					water	50 gallons		
					hellebore	1 oz.		
					flour	2 ozs.		
Gooseberry sawfly	..	..	..	{	water	3 gallons	}	
					Paris green paste	½ oz.		
					water	10 gallons		
					paraffin	2 gallons		
					soft soap	½ lb.		
Apple sucker	..	..	..	{	water	11 gallons	}	Formula 7.
					carbolic acid	2 to 3 gallons		
					soft soap	6 lbs.		
					water	100 gallons		
					caustic soda	10 lbs.		
Woolly aphis (Winter washes)	..	..	..	{	carbonate of potash	10 lbs.	}	
					soft soap	2 or 3 lbs.		
					water	100 gallons		
					soft soap	1½ lbs.		
					paraffin	2 gallons		
				{	caustic soda	6 lbs.	}	Woburn formula.
					water	28 gallons		

Speaking generally, the following are methods in fairly general use:

1. Lead arseniate or Paris green for caterpillars.
2. Soft soap and quassia for hop aphis; soft soap alone or with quassia extract for aphides generally.
3. Sulphur in some form for red spider.
4. Washes containing free caustic alkali for cleaning tree trunks in winter.
5. Paraffin in very weak emulsion with soft soap or paraffin stirred up in water for garden pests.

6. Lime, salt and sulphur in vague proportions for apple sucker, etc.
7. Lime washing for winter use.
8. Nicotine as a general panacea for all pests, as a stomach poison for caterpillars, as a contact poison for apple sucker, aphids, etc.

It will be admitted probably that these insecticides have come into use on purely empirical lines; the use of arseniates has probably come from human toxicology, and a very exhaustive inquiry into the action of some mineral compounds in India¹ has not produced anything more poisonous; it is probably a question of the *form* of arsenic to be used. But what has governed the choice of insecticides that do not act on the internal organs, but those acting from without? I think none but tradition and purely empirical practice, with experience in large scale experiments as a deciding factor.

Is experience on a large scale experiment of much real use? It is in some cases, such as mussel scale, where the insect is one-brooded; I doubt if it is much use in a many brooded or migrating insect, such as an aphid. The factors that control the increase of aphids are many and it would require very careful examination of the conditions to say that the spraying had done the real work, or that the actual nature of the insecticide had been the deciding factor. Suppose a grower of plums sprays for the Leaf Curling Aphid in June; a week after he looks for aphids and finds none; he thinks it is the spraying, whereas it is the migration of the aphids from the plums that has produced the absence.

It is worth while thinking for a moment how these insecticides act; what actually happens when paraffin emulsion is applied; where does the paraffin go to, what tissue does it affect, how does it work? A Board of Agriculture Leaflet states "The presence of the soft soap causes the wash to touch the aphids, and as it dries the thin layer of soft soap clogs the breathing pores and kills the insect" (Hop aphid). Did the author of this statement really know this, or was it simply the generally accepted statement? If so, why do most growers add quassia, and why does quassia kill aphids? what does quassia actually do? I do not think actually that anyone can tell us what the quassia extract does, and I believe that we shall get much better insecticides when we know how it is that these things kill. What is it that happens? If we know that, can we not devise insecticides that make this happen, not that happen to kill?

¹ Lefroy and Finlow. Memoirs of the Agricultural Dept., India. *Entomology*, vol. iv. no. 5.

If we did, it is likely that some of them would be much the same as before, since the sum total of human experience, however got, brings one near the truth. Experience tells us that paraffin kills; it does not tell us how; but knowing how is not going to tell us not to use it. *Paraffin will still kill; only ultimately out of knowing why, we shall know whether paraffin itself is the best or whether there is something better not yet tried.*

In considering the question of insecticides there are a number of points and I would like to suggest two:

1. *Spreading over and wetting the plant.* In many cases it is necessary to spread over or wet the plant in order that the substance contained in suspension or in solution should be spread over the leaf.

If a leaf is sprayed with mercury, the mercury would not spread over or wet it; the mercury would draw into drops and run off.

If it is sprayed with water, the water might spread and it might not 'wet'; *on a cabbage leaf it would not spread or wet, on others it might.*

If you sprayed it with pure paraffin it would spread over and wet all the plant and the paraffin would 'creep.' Why this difference of behaviour? It is partly due to two things: the surface tension of the liquid, high in mercury, middling in water, low in paraffin; the question of the inter-solubility of the liquid of some constituent of the surface of the leaf, *e.g.* the wax on the cabbage leaf, being totally insoluble in the water and causing the water to roll off at once.

To spread, therefore, one must presumably have a low surface tension, and to wet also have a wax solvent or a liquid that will be absorbed by or dissolve the outer coat. Now leaving the question of a leaf with its broad surface, take the case of a bud or a curled leaf: here still more the surface tension question is apparent as the liquid might need to penetrate the curled leaf and act by capillarity. I will quote a case here I had need to go into carefully: we had a disease on indigo in which the plant curls into a compact knot at the tip; in this lives a *Psylla*. It was a question of finding an insecticide that would wet and penetrate. We found by experiment that some insecticides penetrated better than others. We found that they varied according to the concentration, and we eventually found one that penetrated so well as to go right in.

Dr Leather, the Imperial Agricultural Chemist, very kindly measured surface tensions for me and this last had the lowest. It was an "oleic" acid soap at 1 lb. in 12 gallons of water. It was the

best soap we got and it was better than rosin compound, oil emulsions, soda solutions and other things. An oil emulsion in .8 per cent. soap solution has a higher tension than the soap solution. We tested 'penetration' by dipping curled heads of the plants, then cutting them open and seeing what had happened.

Now in this case it was purely one of getting a commercial product of low enough surface tension to wet and penetrate the curled plant. We had to think afterwards of what we could put in to kill the insect. A similar case exactly will be described to you in the paper by Mr P. R. Awati.

Apart from the mere surface tension, what of the different behaviour of a liquid on an ordinary leaf or on a cabbage leaf: is there a reaction between the liquid and the surface depending on the latter? I am not a physicist and cannot say, but I believe there is and, so far as I understand it, I explain it below. But you can get round this in some cases by adding to the liquid a solvent of the wax (if any) in the plant. For instance, a cabbage leaf behaves very differently to a liquid containing a solvent of its wax than it does to the liquid without; for the solvent dissolving the wax eliminates the wetting question since the wax is there already, and it becomes therefore a question of 'spreading.' Spreading power, even more than wetting, depends on low surface tension and we may, I think, distinguish "spreading of a liquid over a surface it never wets" and wetting due to the liquid actually entering into solution with a substance on the leaf surface. You may attain 'spreading' of the liquid by either method but not wetting.

I am indebted to Dr Vaughan-Cornish for a reference to Clark-Maxwell's *Theory of Heat*, in which surface tension is discussed. I have endeavoured to extract the gist of the physics of surface tension so far as it refers to this problem.

The spreading power of a liquid on a solid depends in this way on surface tension:

When a solid body (the leaf) is in contact with two fluids (air, wash) then if the tension of the surface separating the solid from the second fluid (wash) exceeds the sum of the tensions of the other two surfaces, the second fluid will gather itself into a drop and the first fluid spread over the surface, *i.e.*

$$\text{If} \quad \text{S.T. of Wash and Solid} > \begin{matrix} \text{S.T. of Wash and Air} \\ + \text{Air and Solid,} \end{matrix}$$

then the Wash goes into a drop.

If the reverse, the Wash spreads over the whole surface and will wet the Solid.

Assuming then that the surface tension of the air and the solid (the leaf) is always the same, and that the surface tensions of Wash-Solid and Wash-Air vary proportionately according to the nature of the wash, then clearly the measure of the surface tension of Wash-Air is an indication to us of the wetting power, and the lower this is the greater the chance of wetting.

The equation is

$$\text{Wash-Solid} < \text{Wash-Air} + \text{Air-Solid},$$

clearly the lower the two Wash surface tensions, the more likely are we to get the one we want.

The Air and Water surface tensions of these liquids are:

			Air	Water
Water	..	..	8.253	—
Petroleum	..	..	3.233	2.834
Mercury	..	..	55.03	42.58

I have not the surface tensions of any of these with a leaf surface so I cannot work out an equation, but the above figures show why, for instance, mercury and petroleum and water behave differently as regards leaf surfaces.

It will be noticed that it is not only the wash-air surface tension that counts, but the wash-solid and the solid-air. The difference between the wetting power of water on a cabbage and on an apple leaf is due to the difference in surface tension to air and wash between the wax on the cabbage leaf and the substance on the apple leaf. In the first case

$$\text{Wash-Wax} > \text{Wash-Air} + \text{Wax-Air}$$

and no wetting occurs by the wash.

In the second case

$$\text{Wash-Apple leaf} < \begin{matrix} \text{Wash-Air} \\ \text{Apple leaf-Air} \end{matrix}$$

and wetting occurs.

Therefore, the nature of the actual substance on the surface of the leaf determines whether the wash will wet or not.

If then we knew the surface tensions of

plant-air,
plant-wash,
wash-air,

we could always tell if the wash would roll off or not. So for the insect: if we knew the surface tension of

aphis-air,
aphis-wash,
wash-air,

we should know if our wash would wet the aphis or not. As we do not know these, and as our plant-air, or aphis-air, remains a constant, we strive to make the wash a practical success by getting one in which wash-air (measurable) and therefore wash-plant is as low as possible. One imagines that, had insecticides received attention from chemists and physicists, the constants surface tension of plant-air, plant-wash, wash-air could be looked up, as also the more complex surface tensions of emulsions. When drops of oil float in an emulsion they affect the surface tension of the emulsion in relation to air and plant.

2. The next point is this: how do these insecticides act; must the liquid spread over the insect, must they 'wet' it, and, if so, then how do they act?

There are, I think, three points:

- (1) Mere spreading over mechanically.
- (2) Wetting with spreading.
- (3) Toxic action after wetting.

(1) In the first case I take such actions as lime wash, which spread over mechanically and never wet; the action may be due to mechanical causes solely, probably suffocation.

(2) Wetting, including spreading, means the insecticide is in physical contact with the body. What does this really mean? Does it mean that the head, the segments, the antennae and legs are wetted, or that only parts are, or that there is a film of the liquid lying over the whole insect but not actually in contact, say, with the under surface?

I take it to mean that the insect is actually in contact with the insecticide all over without any air-film between. It means, for instance, that the liquid is in contact with the spiracles.

(3) Means that *it also has a toxic action*. Now assuming that we have an insecticide that spreads, that wets and that penetrates to the insect, what happens then we do not know.

With a view to getting some accurate ideas I made a large series of experiments with meal-worms with these results:

1. No action is got by applying any liquid for a short time to the skin, but many act at once if applied to a single spiracle.

2. Interference with the mechanical functions of at least five pairs of spiracles is necessary to produce any symptoms due solely to mechanical interference.

3. Closing all spiracles produces a condition we may call 'rigour,' in which a liquid will enter the spiracles.

4. It is impossible to get a liquid to enter one spiracle unless this condition is produced.

5. Dipping a meal-worm in a liquid produces the condition in which the liquid enters. We can therefore compare the effects of liquids by this means.

6. Of liquids tested the following results were got:

<i>Killed all</i>	<i>Killed some</i>	<i>Killed none</i>
Clove oil	Quinoline	Picric acid in water
Xylol	Carbolic acid	Alcohol 70 %
Turpentine	Formol 4 %	Nicotine 1 % aq.
Nitro-benzene	Pyridine	Acetone
Chloroform	Acetic acid	Ether
Amyl acetate	Eucalyptus oil	Chloral hydrate aq.
Cymene	Methyl salicylate	Water
Pseudocumene	Aniline	
	Acetic ester	

Now results with meal-worms are not of much direct use and we must, for the moment, eliminate the peculiar 'rigour' results. But they show us, I think, that a much greater range of liquids than are generally used have a toxic action and that we must attach far more weight to the details of the structure of the spiracles and tracheal system.

It would look even as if only through the spiracles could one reach most insects with an insecticide, and that the insecticide that did not actually fill the spiracles was not acting. I suggest this is why contact poisons fail against large insects as a general rule; the liquid may wet

but does not wet the whole insect and so the spiracles, or part of them, are free.

The positions of the spiracles, for instance in a Scale Insect, a Psylla, an Aleurodid, and an Aphid, are very different indeed. Their structure is different, they have different arrangements for closing them. In Psylla and Aleurodes the adult is very readily killed with insecticides, partly probably because it has very inefficient spiracles and requires much air, whereas the young form is harder to kill because it has a very efficient closing and protecting apparatus and requires less air.

What I suggest is that the way to tackle the question is first to investigate the tracheal system of the insect. Second, to investigate the wetting action of the insecticide on the plant, unless the insect is fully exposed. Thirdly, to investigate the wetting action on the insect, or the dissolving of its waxes or coating; and fourthly the actual toxic ingredient that would be added to the insecticide to act on the insect when it reached it.

Are any of these factors really taken into account in the recommending of insecticides? I think they should be, and there is here a large field for research.

In preparing this paper I have not attempted in any way to summarise the present practice of insecticides outside this country; this would be a very large subject, complicated by the now enormous use of insecticides sold under fancy names by firms who may or may not have any real knowledge of entomology. But I would like to include abstracts of some papers published in America.

"How contact insecticides kill," G. D. Shafer (*Michigan Sta. Tech. Bul.* xi, pp. 65, pls. 2, figs. 7). This bulletin consists of two parts, the first of which (pp. 8-53) deals with the effects of certain gases and insecticides upon the activity and respiration of insects, and the second (pp. 53-64) with some properties of lime-sulphur wash that make it effective in killing scale insects, especially San Jose scale. Abstracts of these have been previously noted (*E.S.R.*, xxv, p. 665). The investigations conducted and here reported in detail have been summarised by the author in the following general conclusions:

"Usually contact insecticides do not depend upon one property or means alone for their effectiveness, yet as a rule some one property is chiefly concerned. Alkaline washes, corrosive sublimate solution, and other fluids, which are capable either of dissolving or of precipitating certain constituents of insect tissues, are able to penetrate the chitin of insects into the tissues slowly. The weaker the surface tension of the fluid, apparently, and the thinner the chitin with which it is in contact the more rapid the penetration. Gases and vapors may penetrate the chitin of insects, especially through the tracheae into the tissues far more rapidly than liquids.

"It is through absorption into the insect tissues of the volatile portions of kerosene, gasoline, creolin, pyrethrum and such contact insecticides that they mainly become effective agents against insects. Vapors from these insecticides enter the tissues and become effective long before the liquids as such have time to penetrate the chitin. Kerosene, miscible oils, etc., are able to enter the spiracles and tracheae of insects even when a 'closing apparatus' is present; but the comparatively rapid influence which such insecticides exert does not come from the plugging of the tracheae alone.

"The general effects of vapors from gasoline, kerosene, carbon disulphid, creolin, and the rest upon insects are very similar to the effects of the vapor of ether. The nervous system seems to be especially affected. Small amounts of such vapors produce, at first, more or less excitement; then a period of uncertain movements; and finally in larger amounts anesthesia or narcosis. The respiratory activity is usually increased until after the insects become deeply affected, and it is then depressed.

"Certain gases and vapors (*e.g.*, sulphur dioxid, ammonia and hydrocyanic acid gas) when present in respired air continue to be absorbed by insects while they are alive. For the most part, these gases are not given off when the insects are exposed to fresh air, but become rather firmly fixed within the tissues.

"Insect tissues quickly become saturated with any certain percentage of the vapor of carbon disulphid, carbon dioxid, kerosene, gasoline, or similar vapor and no more (at that percentage) is taken up. Then when the insects are exposed to pure air, practically all of such vapors or gases are given off from the tissues again—but not quite as readily as they were absorbed.

"Starvation, serious mechanical injury, and ammonia gas were all found to reduce the value of the respiratory quotient below the value given when healthy strong insects are breathing pure air.

"The vapors of gasoline, carbon disulphid, kerosene and To-bak-ine (*i.e.* nicotine), when present in sufficient amounts to bring the insects near death, cause the value of the respiratory ratio to rise above the value given by healthy, strong insects breathing pure air—*i.e.*, these vapors depress the activity of oxygen absorption more than they do the carbon dioxid excreting activity. The insects tried could continue to give off small amounts of carbon dioxid when no oxygen was present to be taken up, as when they were kept in tested nitrogen, hydrogen, or carbon dioxid.

"The evidence gathered seems to indicate that the vapors of gasoline, kerosene, carbon disulphid and the like, after absorption into the insect body, become mainly effective through some tendency their presence exerts to prevent oxygen absorption by the tissues.

"Lime-sulphur is a special rather than a general contact insecticide. Its strong, persistent reducing power, and its ability to soften the wax about the margin of a scale insect like the San Jose scale, are the important properties that make it efficient as a scalecide." *Expt. Stat. Record*, 1912.

"A contribution to our knowledge of insecticides," C. T. McClintock, E. M. Houghton and H. C. Hamilton (*Rpt. Mich. Acad. Sci.* x (1908), pp. 197–208, pl. I, reprint)...

"The work reported in this paper has to do with contact insecticides only....

"The insecticidal, germicidal, and toxic values (for higher animals) have little or no correlation. It is possible to determine the relative strength or value of insecticides

by immersing test insects in definite strengths of the insecticide, and noting the time required to produce death. The common bedbug (*Cimex lectularis*) appears to be the most satisfactory test insect. As yet the mode of action, the way in which the contact insecticides cause the death of the insects, has not been determined. Apparently, the fewer the number of spiracles, the smaller their size, and the better they are guarded by hairs or valves, the more resistant is the insect to the contact insecticides.

"Chemical standardisation of this class of insecticides is with our present knowledge impossible. With two substances, having essentially the same composition, the insecticidal values may vary enormously. Even the same substance, prepared with what are apparently unimportant chemical variations, gives widely different insecticidal values."

Relative efficiency of contact insecticides.

Kohlentear	125
As ₂ O ₃ soap sol. .. .	50
Terpentin säure .. .	40
Schwafel säure .. .	15
Nikotin	15
Leinol seife	5
KCN	5
Kresyl säure, seifen lösung ..	2.5
Karbol säure, seifen lösung ..	2.0
Morphin sulphat	2.0
Karbol säure, H ₂ O	1.0
HgCl ₂	0.5
HgI	0.5
Formaldehyd	0.4
Alkohol	0.05

"A further contribution to our knowledge of insecticides," C. T. McClintock, H. C. Hamilton and F. B. Lowe (*Jour. Amer. Pub. Health Assoc.*, 1 (1911), No. 4, pp. 227-238, pl. 1): "It is possible to standardise substances which are subject to sophistication or deterioration by comparing the efficient dilution of their vapours with that of a product of known purity. This is particularly applicable to solutions of nicotin and to powdered chrysanthemum flowers. As yet there is nothing from which to conclude what action the vapours have on the insects. If it were merely irritative, formaldehyde would be valuable and the vapours of the burning insect powder without value. If the action were similar to anesthesia, chloroform should have been of greater value. If the action were purely that of poisoning one would have expected the highly poisonous hydrocyanic-acid gas to be of exceptional value for all species of insects."

These show that the way in which contact poisons act is not by any means clear, and that a great deal of research in insect physiology is required before we do know anything about it.

In conclusion, I wish to emphasise the practical bearing of this subject. I may have shed no light on it, but I do think that we owe it

to horticulturists to investigate these problems. The wetting of leaves and plant-buds is one point; the wetting or penetration of insects is another; the toxic action on insects is a third, and the last is to combine the three together in efficient insecticides that really do kill the insect every time and that cause a direct mortality of 100 per cent. In seeking an insecticide for woolly aphis, I would first seek a solvent of its wax. This is easy, but it is far from easy for instance with beech Coccus. In seeking an insecticide for apple sucker we first sought one that would wet apple buds. For mussel scale I would seek either a solvent of the secretory part of the scale, or a liquid that would penetrate under the scale. These are protected insects, but what about naked insects such as bugs, aphides and the like; each has to be tackled individually. So too for stomach poisons. We know that we want to wet the leaves uniformly. Does lead arseniate and water wet well? I think not. In using lead chromate we add 25 % soft soap and 1.5 % of gelatine, simply to secure wetting; it is worth while as we use less liquid to get good even wetting and no dripping. What will wet one plant's leaf may not wet another. We know that for cabbages, for instance, but it is a factor seldom taken into account except with cabbages.

I hope I have shown that there is a field for research in this question. I wish we could get a satisfactory basis of reason on which to proceed, and not have to rely on empirical methods which really are not always successful. And I am inclined to think that in England too much reliance is placed on the tar and water principle to the detriment of the estimation in which applied Entomology is held by horticulturists and fruit growers.