viously stated in my original article, all the evidence is in favor of "these bright bodies" being the ameboid leucocytes circulating through the lymph canals of the cornea. A third and final paper on this subject will dwell on the theory⁸ of chemotaxis and physiology of the white corpuscle, as studied from the cornea.

BIBLIOGRAPHY.

1. Norton: Seeing Capillary Circulation in One's Own Eye; *JOURNAL*, July 15, 1899.
2. Willets: Corneal Corpuscular Activity; *Pa. Med. Jour.*, December, 1894, p. 334.
3. Willets: Address in Ophthalmology, *Pa. Med. Jour.*, September, 1899, p. 145.
4. *Ibid.*
5. Gould: Concerning Corpuscular Phoses and Aphoses; *Phila. Med. Jour.*, July 22, 1899, p. 179.
6. Park: Archives of Ophthal., 1892 or 1893.
7. Stillson, Hamilton: The Subjectoscope; *JOURNAL*, July 8, 1899.
8. Address in Ophthalmology, 1897.

EVISCERO-NEUROTOMY; A NEW OPERATION.*

BY J. G. HUIZINGA, M.D.

Professor of Ophthalmology at the Post-Graduate Medical College;
Chief Ophthalmic Surgeon to the Halsted Street Hospital;
Member of the AMERICAN MEDICAL ASSOCIATION,
the Chicago Academy of Medicine, the Grand
River Valley Medical Society, Etc.

CHICAGO.

The subjects of evisceration and enucleation of the eyeball are just now commanding more attention than in the past, especially in view of the fact that they are to be made the subjects of special inquiry at the next meeting of the International Medical Congress at Paris. Barring special conditions and indications, both enucleation and evisceration have their advantages and disadvantages.

The advantages of the former are a rapid healing of the wound with the minimum of local reaction, and the absolute certainty that it will not be followed by sympathetic inflammation in the other eye. Its disadvantages are that it is more radical than necessary, in that it removes too much tissue and leaves a very poor stump for an artificial eye.

Fig. 1.—Amputation of the cornea.

The disadvantages of evisceration are just the opposite of the advantages of enucleation, viz., a considerable local reaction and an occasional case of sympathetic inflammation. Though the danger from this source is rare, yet it is not unknown, and the operation is to that extent objectionable. Its advantages are a more perfect stump for prosthesis, especially when an artificial vitreous is introduced.

To obtain the advantages of enucleation, viz., absolute certainty that sympathetic inflammation will not occur; as well as the advantages of Mules' operation, viz., a good large stump for an artificial eye, I have originated the following method: Having obtained as aseptic a field of operation as is possible, the cornea is amputated and the opening thus made enlarged laterally by slitting the sclerotic up to the insertions of the internal and external recti muscles.

The contents of the globe are then removed in the same manner as for simple evisceration.

While an assistant keeps the mouth of the wound widely open, a pair of forceps is introduced into the inside of the ball, through the opening made by amputation of the cornea, and the sclerotic coat is caught about midway between the equator and the posterior pole, and button-holed with blunt-pointed scissors.

This opening is then enlarged laterally, parallel with the equator, until it has encircled one-half of the ball. A blunt, curved enucleation scissors is then introduced through this opening and passed back of the globe, and the optic and ciliary nerves are severed. Having thus loosened the eyeball posteriorly, this part can then be drawn forward up to the corneal opening by partly

Fig. 2.—After corneal amputation.

Fig. 3.—Eviscerating the contents of the globe.

everting the sclerotic from behind by the aid of forceps, and that part of it, including the nerves, is removed by continuing the cut parallel with the equator until it extends entirely around the eye. In this manner the posterior segment of the eye, a section somewhat larger than the cornea is removed, and all nerve connection thoroughly and permanently obliterated, while the rest of the sclerotic with its muscle attachments is allowed to remain.

The introduction of an artificial vitreous is very desirable, notwithstanding the fact that it is not always retained and the difficulty occasionally experienced to obtain prompt and perfect union by first intention. Where such unfortunate conditions do not obtain, and they are in the majority, the results are so much better for prosthesis that I believe this practice would be more generally adopted if some material could be obtained

*Read before the Chicago Academy of Medicine, Dec. 8, 1899.

that is more suitable to the purpose than either glass or metal. This subject will be discussed in another paper later. However, the use of the fenestrated aluminium ball in connection with this operation, as advised by Bryant, in implantation, ought also to be followed by good results. As a segment of the sclerotic has been removed, there is no longer any reason why the inside of such a sphere should not fill in with new connective tissue, which, being continuous with that of the orbit, would prevent its extrusion.

Fig. 4. - Removing posterior segment of eyeball.

The wound is closed with five silk sutures, and the after-treatment is the same as in evisceration. The local reaction following this operation has in no case been as severe as that following simple evisceration. In fact, I should say that it has been no more severe than after enucleation.

Fig. 5.—Posterior segment including part of optic nerve removed.

Fig. 6.—Showing sutures and stump after operation.

This operation is strictly in accord with what I consider one of the first rules in surgery, viz., to obtain the maximum in results by the removal of the minimum of tissue.

In conclusion: 1. This method, with the use of an artificial vitreous, leaves as perfect a stump as after Mules' operation. 2. It prevents absolutely the dangers

of sympathetic inflammation. 3. The local reaction appears to be no greater than after enucleation. 4. We obtain the maximum in results by the removal of the minimum of tissue. 5. We obtain, besides this, all the advantages claimed for Mules' operation, as well as those of enucleation without their disadvantages.

100 State Street.

RAILWAY HYGIENE AND EMERGENCY EQUIPMENT.*

BY W. W. GRANT, M.D.

SURGEON CHICAGO, ROCK ISLAND AND PACIFIC RAILWAY CO.
DENVER, COLO.

Coming within the province of duty of railroad surgeons, it may be said with much probability that, unless they take a positive interest in the matter, slow progress will be made in the sanitation and hygiene of railway travel. So far we have devoted ourselves chiefly to the treatment of injuries and the consideration of litigation cases. Especially are the furnishing and ventilation of coaches and first treatment of the injured matters of supreme importance and always worthy of special interest and consideration. It is often a long time after the recognition of an evil before the successful remedy is discovered and applied. The best means in the accomplishment of a given purpose is generally the result of countless theories and suggestions, and careful, patient investigation. The comfort, health and safety of employees and passengers are cardinal elements that must command the serious consideration of railway corporations and the railway medical corps. Should we examine the air of a Pullman coach, with a spectroscope, the myriad living things to the square inch revealed, many inimical to health, would be enough to cause the shades of Schwann, Helmholtz, Kircher, Pasteur and Tyndall to weep for the living.

The ventilation of houses and public buildings has never been easily accomplished, but that of a moving car is a matter of more than ordinary difficulty. The transom-window near the top promptly admits more smoke, cinders and dirt than pure fresh air, and is not a very satisfactory device. The sudden frequent and extreme change of temperature—generally when passengers and porters are asleep—is a fruitful cause of colds and sickness. The current of outside air is much influenced by the motion of the train. The sudden displacement of a large volume of air by the train in rapid motion condenses and gives increased impetus to the air without, which rushes with unusual force through car windows and doors, and always prevents a uniform, comfortable circulation of air within the car.

I may suggest that it might be an improvement if the ventilator windows were opened and closed by a valve-like arrangement, made of the usual glass and wooden frame, and fastened by hinges at one end, preferably toward the engine, on the outside, while on the inside a similar one may be placed, with the fixed, or hinge, end in a reverse direction; a fine wire screen might be placed between. This arrangement would occupy as little space or less than the tilted window now in use. The force and volume of the air current would be broken, and the inside air would be less subject to violent disturbance, and less smoke and dirt would enter the car.

The only perfect remedy for part of the evil is the more complete combustion of fuel and smoke, which is eminently desirable, not only as an economic measure

* President's Address, delivered before the American Academy of Railway Surgeons, Omaha, Neb., Oct. 12 and 13, 1899.