ory, and also because the subliminal self was supremely dominant, and, as is usually the case, no recollection is left of whatever happens during this condition. It would be interesting to know whether, within a few days of the explosion, hypnotism could have recalled to consciousness the recollection of what transpired by bringing the subliminal self again into the foreground.

In the case of S. P., the blow of the ball on the top of the head was the inciting agent. In an instant confused, aimless self-defence—the mere animal instinct of self-preservation—dominated the man; but so entirely motiveless was its demonstration that only purposeless homicidal violence was the result. For years the nervous mechanism, and particularly the subliminal self, had been led to anticipate this moment. Nervous energy was instantly withdrawn from the brain paths whose associative action is identified with normal consciousness, and transmitted with accumulated intensity through other tracts—the combined action of which eventuated in a motor display having for its object self-preservation and defence.

But the utter uselessness of the violence is worthy of notice and an evidence of the automatic and probably unconscious character of the act. In the case of S. P., the violent homicidal onslaught on the innocent and defenceless grandchildren whom he dearly loved, would seem to confirm the probable unconscious automaticity of the act. In G. W., the cyclonic character of the deed in the presence of many people certainly militates against premeditation, and the speechless, dazed condition of the man affords strong confirmatory evidence of its unconscious character. It is also significant that neither at the time nor after did either ever make any allusion to the homicide. The particular period of time seemed as though it were a blank. In none of the furious acts of the fulminating psychoses does the patient ever allude to what transpires at the time. This period is, to all appearances, as much a complete blank to the individual as are the acts of the epileptic when in the midst of his attack. Neither retains the faintest recollection of what has transpired.

In both cases there was rapid heart action, pallor, physical prostration. In the case of G. W., the patient said that everything turned dark—a struggle with a crowding upon him of men, was all that he could remember. This is very similar to the statement of those Malays who have recovered from an amok. I cannot help feeling that the two conditions are quite analogous.

I am well aware that the explanations herein advanced are rather visionary and insusceptible of actual proof. But that there may be a subliminal consciousness, that certain brain paths may be in active operation to the exclusion of certain other tracts that usually are, through their associative action, identified with the normal empirical consciousness—all this seems plausible and not inconsistent with what we already know of brain physiology and pathology.

THE SURGEON-GENERAL OF THE NAVY.—The President has appointed Dr. Newton C. Bates surgeon-general and chief of the bureau of medicine and surgery of the navy, to succeed Surgeon-General Tryon. Dr. Bates has been Mr. McKinley's family physician since the inauguration.

PHTHISIS AND CHILD-BEARING.¹

BY CHARLES W. TOWNSEND, M.D.

THE influence exerted by pulmonary tuberculosis on the course of pregnancy, labor and the puerperium is of considerable interest, owing to the prevalence of this disease and to its somewhat definite course under these conditions.

The following considerations are based on a study of twenty-four cases, all but two being patients of the Boston Lying-in Hospital:

Pregnancy.—During pregnancy the patient often seems better and the disease appears in abeyance, although in some advanced cases severe dyspnea and exhaustion may occur.

The appearance of improvement may have led to the erroneous view that pregnancy is a prophylactic or curative agent in tuberculosis. The subsequent course of the disease disproves this statement and also shows that the improvement is probably deceptive. It is reasonable to suppose that the diminished strength of the pregnant woman offers a favorable opportunity for the beginning of this disease in those who are predisposed.

All of the ten incipient cases went to full term; two of the four moderately severe cases were taken in labor at the eighth month, while, of the nine advanced cases of phthisis, three of them gave birth prematurely at the seventh and eighth month, and in one, labor was induced at that time owing to the feeble condition of the mother.

Labor.—The effect of the disease on the labor is interesting. In the incipient cases, the average duration of the first stage was sixteen hours, of the second stage, forty-three minutes, which is perhaps about the normal duration of labor, considering that one-half were primiparæ. One labor was instrumental. In the four moderately severe cases, one a primipara, the average duration of the first stage was eleven hours, of the second stage twenty-three minutes. In the nine advanced cases, of which three were primiparæ, the average length of the first stage was seven hours, of the second stage twenty-three minutes. In several of these cases coming under my observation the labor was markedly easy.

Instead, therefore, of the labor being prolonged and difficult, owing to the weakness of the patient in advanced cases of this disease, it is short and easy. The muscular power of the uterus seems to be good, while the resistance from the soft parts is slight.

Puerperium.—Nature seems to put forth a supreme effort to suppress the disease during pregnancy and to make the labor easy and short, but after the child is born the disease advances at a rapid rate. This is well shown in all the cases without exception by an examination of the temperature charts. In the incipient cases the temperature in all fluctuated from 98° to 100° or 101°, and in two, at the end of two weeks, the evening temperature rose to 103° and 104°. Weakness and cough were marked.

In the moderately severe cases of phthisis the temperature in all reached 100° and 101° and went to 103° in two cases at night. In the advanced cases a daily evening rise to 102° to 104° and even to 105° was noted.

Notwithstanding the absence of uterine symptoms it was often difficult to exclude sepsis as a cause of the

¹ Read before the Obstetrical Society of Boston, May 18, 1897.

temperature. In all doubtful cases, therefore, the uterus was explored and the result of this examination, together with the subsequent course of the disease, made the diagnosis certain. It is better to consider a case puerperal sepsis until it is proved to be tuberculosis.

All of the incipient and moderately severe cases passed from observation, but of the nine advanced cases, six were observed until their death, which took place in eight days in one case, in twelve days in one case, in one month in two cases, and in two months in the other two cases. In all of these the advances of the disease in the lung were very rapid.

One case remains to be considered. Here the patient was well up to the time of labor with the exception of a slight cough during the last three weeks of pregnancy. The labor was rather a long one, and on the fourth day the patient had a chill, followed by fever reaching 104° and 105° at night. The pain was referred to the chest and there were no uterine symptoms. The uterus was found to be normal, while signs of disease in the lungs were detected which rapidly increased. Tubercle bacilli appeared in the sputa. There was slight improvement during a stay of two weeks at the Massachusetts General Hospital, to which she was transferred from the Lying-in Hospital; her subsequent history is unknown.

This case is an example of tubercular disease of the lung originating in the weakened state of the woman during the latter part of pregnancy or early in the puerperium.

It is probable that some of the incipient cases began in pregnancy, although it was impossible in many cases to state the exact time of origin.

In this connection the observations of Flint on eighty-seven cases of phthisis in married women under forty years of age is of interest. Ten of these developed during pregnancy and twelve others within a few weeks or months after child-birth.

Pregnancy and the puerperium are dangerous periods, therefore, to those with a tendency to tuberculosis.

The Child.—The effect of the existence of tubercular disease in the mother on the condition of the child at birth is next to be considered. In the cases of incipient phthisis all the infants arrived at full term and all were discharged well at the end of two weeks, and one of these was known to be alive and well at the end of a year. The average birth-weight of these ten babies was six pounds and eleven ounces.

Of the moderately severe cases, two were a month premature, one of these giving birth to twins, both of whom died very soon, weighing only three pounds, and two pounds twelve ounces respectively. The other premature child and the two other children were discharged well in two weeks. The premature child weighed five and a half pounds, the others five and a half, and six pounds six ounces respectively.

In the class of advanced cases of phthisis, excluding the four premature cases, the average weight was six and a half pounds.

With the exception of one still-born child weighing two pounds, and one that died in three weeks, all the children in this class, including the premature, were discharged alive at the end of two weeks.

The average weight¹ of full-term children at the Boston Lying-in Hospital is seven pounds nine ounces

for males, and seven pounds five ounces for females, so that the weight of these babies of tuberculous mothers is only about a pound under the average.

A primipara who had been so long sick that she had not menstruated for thirteen months, or for four months before conception took place, who had cavities in the lungs and advanced tubercular laryngitis and who died eight days after a precipitate delivery, gave birth to an apparently healthy child weighing six pounds.

Another case, a multipara, the only out-patient of the hospital included in the list, had cavities in both lungs and laryngitis, and was so weak that she had been in bed for six months with resulting bed-sores. This patient went to full term and gave birth to an apparently healthy child weighing eight pounds and two ounces. The mother died twelve days later. It is certainly remarkable that these patients conceived, and that they gave birth at full term to such large living children. The subsequent history of these children is not known. It is possible that some of the children discharged apparently well may have been suffering from latent tuberculosis resulting from intra-uterine infection. Inherited tuberculosis in the sense of actual intra-uterine infection, the child being born tuberculous, has been proved to occur in so few instances that this accident must be extremely rare. In the absence of congenital tuberculosis, however, there probably exists in these children of tuberculous mothers an enfeebled constitution, rendering them a prey to this and other diseases. These children are further handicapped by the lack of mother's milk and by the constant exposure to tuberculous infection. For the sake of the child as well as to preserve the mother's strength, nursing is contraindicated. Nature generally enforces this rule by refusing to supply milk.

CONCLUSIONS.

(1) Conception may take place even in advanced pulmonary tuberculosis.

(2) The disease is generally held in abeyance during pregnancy, although it may advance or even originate at this time.

(3) Labor is short and easy in proportion to the severity of the disease.

(4) During the puerperium a rapid advance, leading in some cases to speedy death, occurs; or the disease may originate at this time. In either case the temperature chart suggests puerperal sepsis.

(5) Premature labor is more common the more advanced the disease, although pregnancy often goes on to full term even in advanced cases.

(6) The average weight of the full-term children and their general condition at birth is not markedly below that of children of healthy mothers, except in the rare instances of congenital tuberculosis.

A WELL-EQUIPPED MEDICAL SCHOOL.—Among the million or more medical colleges in this great country there is one with thirty-eight professors, twenty-three assistant professors and instructors, and fifty-one students. This gives one professor, assistant professor, or instructor to each student, with ten left over. It is suggested that a new medical journal be started, to give the idle ten something to do.—*Medical Record.*

¹ See Boston Medical and Surgical Journal, May 14, 1896.