standard of morality for the sexes. From the ranks of those so taught, it is quite possible, springs many of those moral and social degenerates, of all classes of society and wealth, who regard with mistaken levity any attempts toward the maintenance of virtuous standards, and whose worthless or vicious offspring contribute a large share to the innumerable types of deterioration. Men should cease to tolerate these pernicious and erroneous ideas, and our youth should understand that a reasonable care for and preservation of the health is a part of morality. Furthermore, in the interests of justice, we should properly require of our sons the full measure of virtue we expect of our daughters. It is an evident truth that what is wrong for a woman is wrong for a man. This truth should receive full recognition in the marriage estate and in all other relations of society, since such recognition is a part of the endeavor to improve the race.

The enduring effects of hereditary laws should convince us that it is necessary that we should understand ourselves, our peculiarities and shortcomings; that we should also heed the lessons of the past; that we owe to every child the birthright of those inherited tendencies which make for strength and health of body and mind; and that proper knowledge should be given children in order that they may be prepared to meet the perplexities of life.

Church and pulpit, school and college, parent and physician, each should contribute a proper share toward the improvement of moral and social influences, and to the publicity and study of various social questions. Fearless and wise education is needful everywhere if the menaces to the general good are to be lessened in any perceptible degree.

Considering the immeasurable influence of heredity on interests of future generations, there is believed to be an urgent necessity for the enactment of legislation which will prevent the perpetuation by hereditary degenerates of their hopeless species. It is certain that the population subject to confinement in our penal and charitable institutions would be greatly diminished through the operation of careful and considerate measures.

When people become fully conversant with the truth, public sentiment will make it possible to correct these evils with justice to all, no matter how radical and unpleasant at first sight such remedies may appear.

PERITHELIAL SARCOMA OF THE UTERUS.*

W. A. NEWMAN DORLAND, A.M., M.D.

Assistant Instructor in Obstetrics, University of Pennsylvania;
Fellow of the College of Physicians of Philadelphia.
PHILADELPHIA.

According to von Wenczel¹ the ratio of the frequency of carcinoma to sarcoma of the uterus has been placed by the authorities as 24 to 1. The site of sarcoma is generally in the corpus or fundus. For each time that sarcoma begins in the cervix there will be eight cases in which it will be found to have originated in the corpus uteri.

The uterine sarcomata spring most frequently from the mucosa—much more commonly from this structure than from the outer wall of the uterus. When a sarcoma

does find its origin in the external wall, it usually originates in a degeneration of a fibromyoma, and is then of exceedingly rapid growth and commonly attains a considerable size.

Sarcomata developing in the uterine wall are most commonly of the spindle-celled variety. This form is four times as common as the round-celled sarcoma. Of the round-celled sarcomata found in this organ the small-celled type is much rarer than the large-celled variety. These general observations will emphasize the special interest attached to the following case:

Von Wenczel¹ reports an instance of the small round-celled variety of sarcoma developing in the cervix of a woman 46 years of age. The tumor, which was ulcerating, had reached the size of a child's fist. The cervix was amputated, but after the report of the microscopic examination a vaginal hysterectomy was performed. No evidence of fibromyoma was visible. The tumor sprang solely from the cervix. The connective tissue of the mucosa was inflamed and hypertrophic and the inflammatory infiltration passed into the substance of the tumor without showing any sharply defined line of demarcation. Accordingly, it was assumed that the tumor developed from the wall of the cervix itself, and not from mucosa, and that it was, therefore, a primary growth.

Lazarus-Barlow,² while speaking of perithelial tumors, remarks that the "group of the endotheliomata must be widened to include certain varieties of growth (occurring especially in the cervix uteri and the breast, but also occurring more rarely in other parts of the body) which are commonly, if not invariably, described as squamous or spheroidal cell carcinomata." From the microscopic appearances of von Wenczel's case, which I have recorded above, I feel justified in believing it to have been a perithelioma of the cervix, which should be included in this group of endotheliomata of Lazarus-Barlow.

Last October I encountered a case in my private practice which was one of more than ordinary interest.

Patient.—An unmarried woman, 65 years of age, for a little over two months had been complaining of pain in the pelvic region. She had visited a woman physician of considerable ability, who had examined her vaginally, and had pronounced the condition rheumatism of the pelvis. The patient was placed on a suitable course of treatment, but without an amelioration of the symptoms. In fact, the pain grew more severe and the patient was referred to me.

Examination.—I found the woman anemic and thin almost to gauntness. The skin was shrivelled and on the abdominal surface could be pinched up in thin folds. It was sallow or pale yellowish in color. After considerable difficulty due to the modesty of the patient, I succeeded in making a vaginal examination, in which the following points were elicited: The vagina was small and undergoing senile changes. The vulvar mucosa showed the characteristic smooth yellow appearance of senility. The uterus lay anteriorly, was small, although larger than would be expected from the age of the patient, and slightly sensitive to the touch. There was a very slight discharge, sanguineous, but without odor or any decided appearance of hemorrhage. At no time had the patient suffered from bleeding, nor had she at any time noticed any odor to the small amount of leucorrhea of which she complained. The history of the patient, her age, the condition of anemia, and the local signs impressed on me the importance of a microscopic examination of the intrauterine curettings and after considerable difficulty this was agreed to. Under ether anesthesia the uterus was scraped at the patient's home and a considerable amount of debris obtained. The examination at this time confirmed the findings of the previous exploration and also demonstrated the absolute mobility of the uterus and the healthy condition of the ovaries and tubes. The scrapings were sent to Dr. A. O. J. Kelly, who reported that

*Read before the West Philadelphia Branch of the Philadelphia County Medical Society, April 11, 1908.
1. *Zentralbl. f. Gynäk.*, 1907, No. 34.

2. *Glasgow Med. Jour.*, 1907, lxxviii.

they showed an unusual condition of great malignancy which warranted the immediate removal of the uterus. He at this time pronounced the growth a perithelial sarcoma, the small tumor-cells showing an unusual degree of grouping around the vessel walls.

Operation.—Two days later I removed the uterus and tubes, performing a panhysterectomy. This, at the request of the patient, was done at her home. I was assisted in the work by Drs. H. A. Slocum, G. A. Knowles and F. A. Faught, and two skilled nurses. The operation was simple and uncomplicated, taking not more than thirty minutes for its completion. The patient made an uninterrupted recovery for three days. Then symptoms of bowel obstruction intervened and despite the use of eserine and the other remedies generally employed in this condition the patient sank steadily and died of paralytic ileus on the evening of the fifth day. Lavage was not resorted to on account of the age and feeble condition of the patient. A water-color drawing (reproduced in the illustration) was made of the fresh specimen, after which it was forwarded to Dr. Kelly for a pathologic report which I here append in full.

Pathologic Report.—"The scrapings consist of a new growth made up of a framework of fibrous connective tissue supporting large cylindrical cells with clear vesicular nuclei. The connective-tissue stroma is quite scanty, and in some places altogether wanting. The tumor cells, for the most part, are arranged radially about and attached directly to capillary blood vessels—whence, morphologically, the term 'perithelial sarcoma' seems justified. In other places there is a diffuse infiltration of the tissue—endometrium and musculature—by the tumor cells. The uterus is rather small and reveals at the fundus a rough excrescence about two centimeters in diameter projecting into the cavity and infiltrating the wall of the uterus to a depth of seven to ten millimeters. Microscopically, the growth consists of a connective-tissue stroma enclosing epithelial cells arranged after the manner of the glands of the endometrium. In general this glandular arrangement is more or less well preserved but in many places it is altogether wanting, and the epithelial cells infiltrate the tissues irregularly, as is common in this type of tumor. The tumor must be called an adenocarcinoma of the uterus. The growth in the uterus is typical microscopically. The scrapings were quite atypical, and serve well to illustrate the wide variations that tumor growths may undergo—indeed, the metaplasia to which their cells are sometimes subject."

In the light of the foregoing report my case assumes a considerable degree of importance, inasmuch as it is the first case reported of perithelial sarcoma of the fundus of the uterus, or of a tumor most closely approximating that unusual neoplasm. While fundamentally an adenocarcinoma, with symptoms differing somewhat from the usual, owing to the metaplasia which its superficial cells had undergone, it gave under the microscope the typical picture of a perithelioma—so much so that in the hands of a very able pathologist the first diagnosis was that of perithelial sarcoma of the uterus.

The literature of perithelial sarcoma—also known as perithelioma, endothelioma, perivascularis and angiosarcoma (A. plexiformis, A. peritheliale) is very scant. I have been able to find but eighteen references to papers dealing with the condition as occurring in various portions of the body. Zietzschmann's paper³ is an exceedingly interesting treatise, covering the recorded information of perithelioma as occurring in lower animals. In the human body authentic cases have been noted as occurring in the cervix uteri (never as yet in the corpus—whence the special interest attached to my own case), on the face, on the lip, on the eyelid, in the pelvic connective tissue, in the testes, behind the ear in the parotid gland, in the brain, on the optic nerve, in

the lymph-glands, in certain serous membranes, in the soft palate, in the subcutaneous connective tissue, in the muscles, in the kidneys, mammae and ovaries, in the bony system, and in the carotid body (the latter very rarely, the tendency of growth here being upward). Wherever occurring, these tumors show a marked tendency to undergo a hyaline degeneration.

The English pathologist, Bayon,⁴ has probably given the best pathologic description of peritheliomata and assigned to them their proper oncologic position. He states that "all growths presenting a radial disposition of cells around blood vessels may be termed peritheliomata, even if it cannot be conclusively shown that they originate from the perithelium."

There has been some misconception as to the exact definition of the various terms employed. Generally speaking the term "angiosarcoma" may be regarded as generic, embracing all sarcomatous tumors occurring in connection with the blood vessels, whether capillary or of larger caliber. This generic term may then be regarded as including the two sub-groups, the endotheliomata or those tumors springing from the endothelial cells and the peritheliomata or those arising from the perithelium, which, according to Bayon,⁴ "is a purely topographical expression."

Endothelial cells are those cells which line the lymphatics and blood vessels, and these take their origin in the cells which form the perithelium so called, the layer of cells which surrounds the capillaries and smaller vessels. The perithelium has been demonstrated conclusively around the vessels of the brain and cord. Eberth and Iwanoff, I believe, were the first to find that the vessels of the pia mater showed the perithelium, and they proved that this structure constituted the external boundary of the vessels which ran through the so-called perivascular spaces and that it did not form the external wall of these spaces. The perithelium has also been described as existing around the vessels of the pineal gland, the mammary gland, the suprarenal capsules, the thymus gland, the salivary glands (Ebert, Leydig, Henle, Kölliker and Boll), and in the vessels of the testicles (Ebner, Leydig, Henle, Kölliker and Boll). A similar, although probably not identical, layer of cells was described by Waldeyer and Sertoli in the coccygeal gland and by Paltauf in the carotid gland.

Bayon⁴ states that "the consideration of perithelium as a special class or quality of cells has in no case met with the approval of any histologist so far as he knows. It is also uncertain as to what and how far layers of cells disposed round vessels in various organs deserve to be called by a special name."

An endothelioma should accordingly be defined as a growth taking its origin "from the endothelial layer of cells of blood vessels and lymph sinuses, the growth and these cells being both contiguous and continuous." The name should be exclusively reserved for those growths which really show similarity in disposition, position and general morphology to genuine endothelium (which it must be remembered, is solely and alone the lining membrane of blood vessels and lymphatics). From a morphologic standpoint it is admissible to make use of the expression "peritheliomata" to designate a peculiar class of growths springing from the delicate external layer of cells around the minute vessels and showing the characteristic radial disposition of the tumor cells.

3. Beitrag zur Kenntniss der Peritheliome angiosarcome bei Thieren. Ztschr. f. Thiermed., 1904, viii, 417-427.

4. Brit. Med. Jour., 1907, ii, 1400-1402

According to Vignard and Mouriquand,⁵ who report a case of perithelioma of the face in which ligation of the primitive carotid was practiced, with death in four days, the greater portion of a perithelioma is composed of capillary vessels more or less distended. The characteristic cellular proliferation begins at the external wall of the vessel and not in the lumen of the vessel, as in simple hemangio-endothelioma. These peritheliomata are characterized by a radial disposition of cells around the vessel, as has already been stated, and, according to von Hansemann, this disposition is due to the growth-cells being more readily nourished in that position.

Bayon states that metastases are not at all frequent with such tumors, which is diametrically opposed to the general expectation when there is taken into consideration the extreme vascularity of the tumors, their intimate approximation to the blood vessels, and their multiplicity of small cells. They are, however, characterized by infiltrative and rapid expansive growth, but on complete and thorough operative procedure do not show a tendency to recur.

An interesting question associated with peritheliomata and, indeed, with tumors generally, is that which concerns the metaplasia to which their cells are subject, and which my own case showed to a remarkable degree. In this connection Bayon offers a very important suggestion. He wishes to know whether sarcomata showing this perithelial arrangement of cells are in the early stages of development, and whether they ultimately become carcinomata? This question he answers as follows:

It can most definitely be proclaimed that there exists at this moment no conclusive proof of the fact that a carcinoma cell is ever anything else but a carcinoma cell, from the very beginning of its existence in the human or animal body. The same may be said of sarcoma, though with sarcoma a certain amount of circumstantial evidence exists which renders its origin from preexistent, apparently normal cells probable; this, I repeat, is not the case with carcinoma.

Here is a question of vital interest in histology and pathology which may ultimately exercise an important influence on the study of oncology.

The other papers, bearing on the subject, which I have been able to collate from the world's literature are as follows: Schmincke,⁶ who reports two cases originating in the left arm of men 62 and 65 years of age; Toyosumi,⁷ of Tokio, a case of papillary cystic lymphangio-endothelioma of the abdominal wall in a woman 52 years of age; Faucon and Augier,⁸ an endothelioma of the jaw, both melanotic and exceedingly vascular, in a woman 56 years of age; Nitsch, who publishes a brochure⁹ reporting a case of endothelioma of the pelvic connective tissue; Wingrave,¹⁰ cases occurring in his laryngologic practice; Znojensky,¹¹ a cystic endothelioma of the face; Lazarus-Barlow,¹² a case with general observations; Looser,¹³ one case of endothelioma; A. Bevacqua,¹⁴ a case of retroauricular perithelioma; A. Carless,¹⁵ a general article on the subject; F. Cinotti,¹⁶

a case of telangiectatic cervicofacial tumor in a man 60 years of age, invading the jaw; and some peritheliomata of the osseous system by Ritter,¹⁷ Müller,¹⁸ and Crile and Howard,¹⁹ who report four new cases and nineteen culled from the literature of the world.

1623 Walnut Street.

Clinical Notes

OVERDOSAGE OF VERONAL.

JOHN E. CLARK, M.D.
DETROIT.

The following instance of overdosage of veronal is, I believe, sufficiently unusual and interesting to be worthy of record:

Patient.—A business man, aged 35, with no organic lesion, was greatly distressed during the recent panic by an investment of several hundred thousand dollars in mining securities, which were steadily decreasing, and had resorted to an occasional "bracer" as a mental stimulant.

History.—He consulted me recently, suffering from melancholia and persistent insomnia, results of worry and too liberal indulgence in the aforesaid "bracers." I had treated him twelve months before for similar trouble, when he responded to comparatively large doses of trional, which relieved the insomnia at that time. The sequelae, however, were such that this time I determined to use another hypnotic and selected veronal, hoping to overcome the vertigo and incoordination of movement which had followed administration of the trional. Knowing the patient thoroughly, I had no hesitation in prescribing veronal in ten-grain doses, to be repeated not oftener than once every three hours. He was instructed not to take more than three powders without telephone consultation.

Result of Veronal.—The following morning I was called to the telephone about 8 a. m. by his wife, who said: "My husband took ten of your powders last night; he is still sleeping." I at once went to the patient and found him in a deep sleep. Pupils were dilated, respiration was shallow, pulse 72; there was no cyanosis, but it was impossible to arouse him. At 2 p. m. the same condition existed. At 9 p. m. he was sleeping quietly and could not be aroused. His wife stated that he had swallowed a little coffee and broth. The kidneys had not acted.

Subsequent History.—On Monday morning the patient was still sleeping and apparently in the same condition as on the previous day. Urine had passed in slight quantities. Pupils were still dilated and the patient seemed to respond to shouting in ear, but did not respond to slapping or pinching. In the evening, pupils were still dilated, and sluggish reaction; pulse was somewhat weaker, temperature was 97.4 F. The surface of the body was moist. He had swallowed coffee and some egg-nog during the day. During the evening he slept continuously, but toward night he became slightly restless, moving from one side of the bed to the other, and moaning at times.

Tuesday morning: The man had passed a quiet night; he responded to shouting in ear; pulse was full, 75; respiration was 20 and deeper. He awakened sufficiently to recognize his wife. At noon he was raised to a semi-recumbent position and partook of broth and malted milk. Urine examination showed specific gravity 1.022, acid reaction, no albumin, no sugar. No other symptoms appeared.

Tuesday evening: The patient recognized me. He complained of feeling tired and wanted to sleep; he slept continuously until Wednesday morning, when he had recovered sufficiently to complain of vertigo. He went to the bath-room

5. Rev. de chir., Paris, 1905, xxxii, 467-478.
6. Zentralbl. f. Chir., 1907, xxxiv, 1292.
7. München. med. Wochenschr., 1907, liv, 1985.
8. Jour. d. sc. méd. de Lille, 1907, T. 233.
9. Jena, 1906, B. Vopelius.
10. Jour. Laryng., 1907, xxii, 41-44.
11. Znojensky, J.: Casop. česk. lékařn., v. Praze, 1907, xli, 10, 37, 74.
12. Arch. Middlesex Hosp., 1906, vii, 79-87.
13. Beitr. z. klin. Chir., 1906, lli, 261-274.
14. Glor. med. d. r. esercito, 1906, liv, 337-348.
15. Practitioner, London, 1906, lxxvi, 228-242.
16. Perithelioma all' ascella di un gatto n. ercolani, Pisa, 1905, x, 25, 47, 81.

17. Deutsch. Ztschr. f. Chir., 1899, l, 348.
18. Arch. f. Chir., 1901, lxiii, 348.
19. Ann. Surg., September, 1907, 358.