fessor Henslow and others that it was an anatomical impossibility, was entirely disproved by the large size of the œsophagus before the Society.

November 13, 1855.

Dr. Gray, F.R.S., in the Chair.

1. ON A NEW GENUS AND SPECIES OF TROCHILIDÆ FROM
ECUADOR. BY JOHN GOULD, F.R.S. ETC.

The remarkably fine species of Humming Bird which I am about to describe, I have lately received from Ecuador. This new bird is remarkable for its large size, deeply forked tail and the harmonious hues of its plumage, which, although less glittering and metallic than in many other species, is nevertheless strikingly beautiful. I consider this bird to be new to science, both generically and specifically, and as the name of *Victoria regia* has been given to one of the finest flowers of the same part of South America, I am desirous of dedicating this new Humming Bird to the Empress of the French, and I accordingly propose to name it *Eugenia imperatrix*. Its native habitat is the vast Andean forests in the neighbourhood of Quito in Ecuador, where it procures its insect food from the bell-shaped flowers of the *Datura*.

Genus EUGENIA.

Gen. Char. Bill rather lengthened, straight and strong; wings long and pointed; tail lengthened and very much forked, the feathers narrow and rigid; tarsi clothed with feathers nearly to the toes; feet small.

EUGENIA IMPERATRIX.

Male. Face and fore part of the neck brilliant grass-green; crown of the head, back, neck, chest and upper part of the flanks very deep-green; on the centre of the throat a gorget of reddish-violet; abdomen and under tail-coverts shining, greenish-yellow; wings purplish brown; tail deeply forked, the feathers black, narrow and rigid; some tufts of white downy feathers across the lower part of the abdomen; thighs brown in front, white behind.

Total length $6\frac{1}{4}$ inches; bill $1\frac{1}{4}$; wing $3\frac{1}{3}$; tail 3.

Female. Upper surface green; throat, chest and abdomen grayish-white spangled with green, the spangles being very minute on the throat and gradually increasing in size downward to the flanks; tail blackish-brown; tarsi white.

Total length $5\frac{1}{2}$ inches; bill $1\frac{3}{8}$; wing $2\frac{3}{4}$; tail $2\frac{3}{8}$.

Hab. Ecuador.

J. Wolf. lith.

M & N. Hanhart, Imp^t

BUCCO HYPERRHYNCHUS. Bp.

J Wolf, lith.

M. & N. Herbert, Imp^t

BUCCO PULMENTUM. *Ep. et Verr*

2. CHARACTERS OF SOME APPARENTLY NEW SPECIES OF BUCCONIDÆ, ACCOMPANIED BY A GEOGRAPHICAL TABLE OF THE FAMILY. By PHILIP LUTLEY SCLATER, M.A., F.L.S.

(Aves, Pl. CV. CVI.)

1. BUCCO HYPERRHYNCHUS. (Pl. CV.)

Tamatia hyperrhynchus, Bp. MS. et Consp. Vol. Zygodact. p. 13.
B. supra fulgenti-niger; fronte lata et supercilii anticis albis:
subtus albus, nigro late torquatus; lateribus nigro radiatis:
rostro maximo.

Long. tota 10·5; alæ 4·5; caudæ 3·5; rostri a rictu 2·1.

Hab. In regionibus fl. Amazonum superioris (*Hawxwell*). Mus. Paris. et P. L. S.

When I drew up the characters of *Bucco macrorhynchus*, as given in the 'Annals of Nat. Hist.' for May last (p. 357), I had not in my possession specimens of the true *macrorhynchus* from Cayenne, and consequently confounded with it the present bird. But the much larger size of the bill and whole body, the greater extension of the white colour on the front, the narrower black band and the total absence of any fawn-coloured tinge on the belly and vent are quite sufficient to distinguish this *Bucco* from its Cayenne representative.

I have lately ascertained, through the kindness of Prince Bonaparte, that this is the species included under the name *Tamatia hyperrhynchus* in his *Conspectus Volucrum Zygodactylorum*, published in the 'Ateneo Italiano' of May last, and I have therefore adopted his specific designation. But no descriptions have yet appeared of the many new species of which the names only are inserted in that and other similar recent publications of the Prince.

The type specimens of the present bird are in the French National collection.

2. BUCCO DYSONI.

Tamatia gigas, Bp. Consp. Vol. Zygodact. p. 13?

Bucco dysoni, G. R. Gray in Mus. Brit.

B. supra fulgenti-niger; fronte usque ad oculos et collari postico albis: subtus albus; vitta pectorali lata nigra; lateribus nigro variis; rostro pedibusque nigris.

Long. tota 9·7; alæ 4·5; caudæ 3·4; rostri a rictu 1·8.

Hab. In America Centrali, Honduras (*Dyson*). Mus. Brit.

Obs. Species a *Buccone macrorhyncho* fronte latius albo, rostro majore, et ventre pure albo, a *Buccone hyperrhyncho* rostro minore et fronte minus albo diversa, et inter has duas media locanda.

A single specimen of this bird in the British Museum was procured by Mr. Dyson in Honduras. In my Synopsis of this family I have confounded it with its near affines, from which I now think, as might have been expected from the locality, it will bear separation. It is very probable that Prince Bonaparte's name, *'gigas'* (which was applied to a bird brought by Delattre from Nicaragua), was intended

for this same species, but as the type has disappeared, and no specific characters have been published for the name, it is difficult to be certain on that point.

3. BUCCO PULMENTUM. (Pl. CVI.)

Tamatia (Nyctactes) pulmentum, Bp. et Verr. MS.

B. supra fusco-brunneus; fronte et superciliis rufescentibus; torque angusto nuchali inconspicuo albido; dorsi medii alarum uropygiique plumis partim fulvo terminatis: subtus albus; gutture inferiore pallide rufescente; plaga utrinque gutturali magna cum maculis crebris pectus totum et ventrem (præcipue ad latera) occupantibus atris: rostro nigro.

Long. tota 5·0; alæ 3·1; caudæ 2·5.

Hab. in Peruvia Orientali et regionibus fl. Amazonum superioris: Pebas (*Cast. et Dev.*): Chamicurros (*Hawxwell*). Mus. Paris., Joh. Gould et P.L.S.

Obs. Sp. *Bucconi tamatiæ* affinisissima, sed gula pallidiore et maculis ventris majoribus et intensioribus differt.

This appears to be a western representative of the *B. tamatia* of Cayenne, from which, however, I think it may be fairly separated. MM. Verreaux of Paris have lately received a considerable number of specimens of it from the Upper Amazon. They all present the same distinctive characters as are above noticed.

4. MONASA PERUANA.

Monasa peruana, Bp. et Verr. MS.

M. plumbescenti-nigra, capite et gutture intensioribus; fronte et menti summa parte albis: rostro ruberrimo.

Long. tota 11·0; alæ 5·0; caudæ 4·5.

Hab. In Peruvia Orientali in regionibus fl. Amazonum superioris: Chamicurros (*Hawxwell*). Mus. Joh. Gould et P.L.S.

Obs. Simillima *M. personata* sed rostro clarius rubro, mento ad ipsam apicem solum albo haud nisi dubie disjungenda.

My specimen of this bird was obtained from the MM. Verreaux, and carries the MS. name above quoted, which I have thought it as well to adopt. The characters which separate it from its well-known Brazilian representative are certainly very slight, but appear to be constant in at least a dozen examples I have examined from the same locality.

5. BUCCO PICATUS.

B. supra niger; plaga in summis scapularibus utrinque magna et maculis in pileo rotundis cum loris albis: subtus albus; vitta lata pectorali nigra: cauda nigra, rectricibus tribus utrinque extimis in medio et harum omnium apicibus albo maculatis: rostro pedibusque nigris.

Long. tota 6·7; alæ 3·2; caudæ 2·3.

Hab. In reg. fl. Amazonum superioris; Chamicurros (*Hawxwell*). Mus. Joh. Gould.

Obs. Species *Bucconi tecto* forsan nimium affinis, et ob crassitiem majorem, torquem pectoris latiore et caudam minus albo maculatam non sine dubio constituenda.

Mr. Gould's collection contains two examples of this bird, which he has entrusted to me for comparison with its Cayenne representative. It is not, however, without hesitation that I have determined to separate them from it. Besides their larger size and broader breast-band, the white medial square spot extends in the present species only through the three lateral rectrices, with a slight trace of it in the fourth.

In the Cayenne bird the outer five pairs are all strongly marked thus. The whole plumage of the bird is also generally more intensely black.

6. MALACOPTILA NIGRIFUSCA.

Malacoptila fusca, ex Bogota, Sclater, P. Z. S. 1855, p. 136.

M. nigricanti-brunnea, plumarum scapis pallide fulvis; loris et plumis mystacalibus cum plaga triangulari super-pectoralis albis: ventre medio crissoque fere unicoloribus, albicantioribus; rostri basi læte aurantio, apice nigra; pedibus nigricanti-brunneis.

Long. tota 6·5; alæ 3·5; caudæ 2·5.

Hab. In Nova Granada, Santa Fé de Bogota. Mus. Brit. et Joh. Gould.

Obs. Sp. *Malacoptilæ fuscae* affinissima sed statura minore et coloribus nigricantioribus: rostri basi lætius aurantio.

This New Grenadian bird, which in my Synopsis of this family and List of Bogota birds I united with the true *fusca* of Cayenne, certainly presents considerable claims for specific distinction. The body is generally smaller, the bill in particular is shorter and not so strong, and at the base is of a deep orange colour instead of pale yellow, the black not extending so far towards the base of the upper mandibles; also, the markings on the head, throat and breast are much blacker, and I have therefore named the bird *nigrifusca*. There are specimens of it in the British Museum and in Mr. Gould's collection.

The East Peruvian or High-Amazon examples on the other hand (which are held distinct by some naturalists under Du Bus' title *inornata*) resemble the Cayenne bird much more nearly. After remarking that the white lore-spot is nearly obsolete, and the skins are rather finer and larger, it is in truth difficult (at least with my present examples) to see further differences, and I therefore regard *M. inornata* as a very doubtful species.

Rio Napo specimens are still more like the true *fusca*.

BUCCONIDARUM SCHEMA GEOGRAPHICUM.

[illegible]

3. ON SOME NEW SPECIES OF FRESHWATER TORTOISES FROM NORTH AMERICA, CEYLON AND AUSTRALIA, IN THE COLLECTION OF THE BRITISH MUSEUM. BY DR. J.E. GRAY, F.R.S., F.R.G.S. ETC.

Fam. I. EMYDIDÆ.

The freshwater Tortoises which have been referred to the genus EMYS, as it is at present constituted, may be divided into two very distinct genera; and this is the more advisable as it is extremely difficult to distinguish the American species of which it is composed, and the separation of any of them by organic characters must facilitate the process. The genera may be thus named and defined:—

1. EMYS.

The lower jaw rounded beneath, and covered with, the hinder part of the horny beak; the toes strong, covered with broad band-like scales.

This genus includes *E. ornata*, *E. scripta*, *E. Holbrookii*, and many other species, both Asiatic and American.

2. PSEUDEMYS.

The lower jaw flattened beneath and covered with a soft skin. The toes weak, slender, covered with small scales above, and very broadly webbed.

1. *Pseudemys concinna*.

2. *P. serrata*.

The genera *Batagur* and *Malaclemys* have nearly similar feet, and they appear, like *Pseudemys*, to be the most aquatic animals of the family.

The species which have hitherto been referred to the genus CISTUDO differ considerably in their habits, some being nearly terrestrial and others almost exclusively aquatic. The examination of the animal shows that there are good external characters by which they may be divided into natural groups agreeing with their habits and their geographic distribution.

I. The more terrestrial have the front of their legs covered with thick, imbricate, triangular scales, the toes only slightly webbed, and the sternum broad, hiding the legs when withdrawn, as—

1. CISTUDO.

The head rhombic, the forehead flat, and eyes lateral: confined to N. America; as

C. Carolinensis, with four, and *C. Mexicana*, with only three toes on the hind feet.

2. LUTREMYS.

The head oblong, depressed, with the eyes on the upper part of the cheek. Found in Europe, as *L. Europæa*.

II. The more aquatic kinds have the front of the legs covered with small scales and some broad, transverse, lunate plates; the toes webbed. They are confined to Asia; as

3. CUORA.

The head rhombic, the eyes lateral, the sternal lobes broad, hiding the legs when contracted, as *C. Amboinensis* and *C. trifasciata*.

4. CYCLEMYS.

The head depressed, eyes subsuperior, the sternal lobes rather narrow, not hiding the legs when contracted, as *C. dentata* and *C. platynota*. The latter species was referred to the genus *Testudo* by F. Müller, and when I first described it I considered it as an *Emys*, but the examination of a series of specimens of different ages shows that it is a species of Box Tortoise nearly allied to *C. dentata*.

It has been hitherto believed that there was only a single species of the genus KINOSTERNON, as now restricted, found in the United States; and all the adult specimens I have received from that country are, I must own, exceedingly alike, so much so that I cannot undertake to say that we have adult specimens of more than a single species. On examining the young specimens of this genus from the United States, in the Museum Collection, it is evident that there are at least three most distinct species found in that country, which probably in their eroded and discoloured adult state are so alike as to be mistaken for one another.

They may be thus described:—

1. KINOSTERNON PENNSYLVANICUM.

Head brown-dotted; temples with two parallel distant streaks of white spots, from the upper and lower edge of the orbit, and a third streak across the lower jaw; neck white-dotted; back deep brown; lower side of marginal shields, the axillary and inguinal plates and each of the sternal plates with a large yellow spot; sternum broad, rounded before and behind.

Hab. North America, Florida, *E. Doubleday, Esq.*

2. KINOSTERNON HIPPOCREPIS.

Head brown, with a broad white streak on each side, from the end of nose over the eyebrows to the sides of the nape; back pale and sternum brown; dorsal shield with a single apical and some scattered black spots; under side of each marginal and sternal plate rather paler in the middle; sternum rather broad, rounded in front and slightly truncated behind.

K. Pennsylvanicum, Holbrook, N. Amer. Herp. t. 21.

Hab. North America, New Orleans, *E. Doubleday, Esq.*

3. KINOSTERNON PUNCTATUM.

Head brown, minutely white-dotted, without any streaks; the back brown, discal shield with a very distinct apical, and some scattered black spots; margin with a very narrow white line; under side whitish,

with minute scattered black dots and line; sternum narrow, contracted at each end, and with straight sides behind, rather truncated in front and more distinctly and broadly so behind.

Hab. North America.

There are several specimens of the first species of different ages, from various parts of the States, in the British Museum; I have therefore retained for it the more general name; and two young specimens and a half-grown one of the second species, and only a single young specimen of the third species; the latter is so distinct, by the narrow form of its sternum, from the other two, that it might be referred to the genus *Aromochelys* if the pectoral plates were not triangular; it may be considered as the species passing towards that genus, and I should think that the adult animal must differ considerably from the common form of *K. Pensylvanicum*.

AROMOCHELYS.

The Musk Tortoise, or, as it is more commonly called, the Stinkpot of North America, is easily distinguished from the other *Kinosterna* by the narrowness of the sternum and the humeral plates being square, like the pectoral one, instead of triangular, as they are in *K. scorpioides* and *K. Pensylvanicum*. For this reason I have proposed to divide them into a distinct group under the name of *Aromochelys*.

I am the more inclined to do so, as there are two most distinct species in the British Museum Collection, which have either been confounded together by the American naturalists, or have been most unaccountably overlooked. They may be thus defined:—

1. AROMOCHELYS ODORATA.

Head moderate, with two streaks from the nose, one above and the other under the eyes, to the side of the neck; the back oblong-convex, the vertebral line rather flattened; the gular plate small, triangular, the humeral plate rather oblique, shield brown, purple-brown spotted.

Holbrook, N. Amer. Herpet. t. 22.

Hab. United States and Louisiana.

2. AROMOCHELYS CARINATA.

We have four specimens of this species in the Museum Collection. Cat. Tortoises B.M. t. 20 a.

Head very large, black-dotted, without any lateral streaks; back oblong, very high, the vertebral line high and acutely keeled the whole length, shields grey-brown, spotted and lined with purple-brown; the gular plate very small, linear, transverse marginal, the humeral plate square, transverse, parallel to the pectoral plates.

Hab. North America, Louisiana.

There are two species of North American Tortoise which are referred to the genus *CHELYDRA*, which are so differently organized that they are evidently the types of two very distinct genera, which may be thus characterized:—

1. CHELYDRA.

Head moderate, rather depressed, covered with a soft skin, chin bearded, neck granular; back with two slight keels; marginal plates in a single series.

Chelydra serpentina.

2. MACROCHELYS.

Head large, angular, contracted in front, covered with symmetrical horny plates, neck with several series of spinose warts; back with three sharp continued tubercular keels; the lateral marginal plates in a double series.

M. Temminckii.

Fam. II. CHELYDIDÆ.

When Australia was first visited by Sir Joseph Banks, he brought home with him from New Holland a freshwater Tortoise, which Dr. Shaw described under the name of *Testudo longicollis*. This has been made the type of the genus CHELODINA. Recent travellers in Australia have shown that the genus is distributed over the country; each part appears to have a species peculiar to itself. In Capt., now Sir George Grey's Travels, I described and figured a species from Western Australia under the name of *Chelodina oblonga*. In a collection which we have lately received from Haslar Hospital, there are two very large specimens of the genus sent from Swan River by the late Mr. Collie, which, though similar in several respects to *Chelodina oblonga*, may be considered as a distinct species, which I shall proceed to shortly characterize.

The species of the genus hitherto described have the thorax covered with very thin smooth shields, so transparent that a peculiar black reticulated appearance, which exists between the shields and the bones of the thorax, can be distinctly seen through them. This character is common to *C. longicollis* of New Holland, *C. oblonga*, and Mr. Collie's species from Swan River, which I propose to call, in honour of my late friend and excellent collector—

1. CHELODINA COLLIEI.

The shield oblong, elongate, contracted and revolute on the sides; under side uniform pale yellow.

Hab. Swan River, *Alexander Collie, Esq.*

This species agrees with *C. oblonga* in the uniform colour of the back and sternum, which is only varied by the dark lines of the netted appearance before referred to; but it is easily known from that species by its larger size, the much narrower shape, and the lateral margin becoming strongly revolute, and the edge over the hinder limb raised up and rather expanded.

The British Museum have lately received, with some other specimens, from the Australian continent—but unfortunately the special habitat was not indicated—the shell of a Tortoise which has all the characters of the genus as at present defined, except that, instead of

the shields on the thorax being thin, submembranous and semitransparent, they are thick, horny and concentrically grooved like the shields of many other genera. It is not accompanied by the head or limbs of the animal, so we have not the means of determining if they offer any characters which, with the peculiar structure of the shell, might render it desirable to form it into a separate genus. It may be defined and thus named :—

2. *CHELODINA SULCATA*.

Shell depressed, roundish ovate, brown; shield horny, thick, distinctly concentrically grooved.

Hab. Australia.

Fam. III. *TRIONYCIDÆ*.

The species of this family, which have the hind legs covered with moveable flaps affixed to the sides of the hinder lobe of the sternum, named *Cryptopus* by MM. Dumeril and Bibron, may be divided into two very distinct geographic genera.

1. *EMYDA*, Gray.

The margins of the upper shield strengthened with bones; the sternum with three pairs of callosities and a small odd one behind the anterior pair. Asia.

2. *CYCLANOSTEUS*, Peters.

The margin of the upper shield flexible, without any bones; the sternum with four pairs of callosities and an odd one behind the two anterior pairs; the pair on the hinder lobes small, far apart. Africa.

It has been usually stated that the only known species of the genus *EMYDA* was generally distributed over India; we have in the British Museum specimens only from the Valley of the Ganges. The young specimens all agree in the head and shell being variegated.

We have lately received a specimen of this genus from Ceylon, collected by Mr. Thwaites, which differs in both the above particulars; and in the Museum of the Society there is the shell of an adult animal, sent home from Ceylon by Dr. Kelaart, which shows that it is a most distinct species. They may be thus characterized :—

1. *EMYDA PUNCTATA*.

Back and upper part of the head pale spotted; the odd anterior callosity small, roundish triangular; the hinder callosity of adult ovate, inner edge semicircular; of young triangular, far apart.

Hab. India, Ganges.

2. *EMYDA CEYLONENSIS*.

Back and upper part of the head (in spirits) dull pale olive; lips, chin and lower part of the body whitish. The odd anterior sternal

tubercle large, oblong, transverse; the hinder pair of callosities large, close, in adult nearly united, with straight parallel inner edges.

Emyda punctata, Kelaart, Prod. Faun. Ceylon. 179.

Hab. Ceylon.

Dr. Kelaart, in his work on the Ceylon animals, was not aware of the distinctness of this animal from the continental species; he observes that the head is black-lined when alive.

The new species described in this paper will be figured in the Illustrated Catalogue of Tortoises in the Collection of the British Museum, which will shortly be published.

4. DESCRIPTIONS OF TWENTY-THREE NEW SPECIES OF *ACHATINELLA*, COLLECTED BY MR. D. FRICK IN THE SANDWICH ISLANDS; FROM MR. CUMING'S COLLECTION.
BY DR. L. PFEIFFER.

Sect. *BULIMELLA*.

1. *A. NOBILIS*, Pfr. *A. testa subimperfata, dextrorsa, turrata, solida, striatula, nitida, fulvida vel griseo-virente, saturatius strigata; spira exacte conica, apice alba, acuta; sutura leviter marginata; anfr. 6 $\frac{1}{2}$ —7 planiusculis, ultimo $\frac{2}{3}$ longitudinis sub-æquante, infra medium obsolete subangulato; apertura obliqua, obauriformi, intus alba; perist. expansiusculo, margine dextro intus late labiato, columellari subadnato.*

Long. 23, diam. 11 mill.

Hab. Oahu.

2. *A. HANLEYANA*, Pfr. *A. testa subrimata, dextrorsa, globoso-conica, solida, lævigata (sub lente exilissime spiraliter striata)* nitida, fulva, castaneo-radiata; spira conica, apice minuto, corneo; sutura filo crenulato marginata; anfr. 6, superis planis, sequentibus modice convexis, ultimo spiram fere æquante, rotundato; apertura obliqua, truncato-auriformi, intus alba; plica colum. supera, valida, dentiformi; perist. intus labiato, margine dextro breviter reflexo, columellari dilatato, calloso, subadnato.*

Long. 18, diam. 11 mill.

Sect. *NEWCOMBIA*.

3. *A. ZEBRINA*, Pfr. *A. testa imperfata, dextrorsa, tenuiuscula, lævigata (sub lente minutissime decussata), nitida, albida, strigis fulgurantibus nigro-castaneis elegantissime picta; spira subturrata, apice obtusa, fusca; sutura filomarginata; anfr. 5 modice convexis, ultimo $\frac{3}{4}$ longitudinis fere æquante; apertura vix obliqua, truncato-oblonga, intus lilacina; plica colum. supera, torta, callosa, vix prominente; perist. simplice, recto, fusco-limbato.*

Long. 12 $\frac{1}{2}$, diam. 6 mill.

4. *A. CHLOROTICA*, Pfr. *A. testa subperforata, ovato-conica, solida, ruguloso-striata, albida, epidermide virente strigatim variegata; spira conica, acutiuscula; anfr. 6, superis convexiusculis, penultimo turgido, ultimo $\frac{2}{3}$ longitudinis subæquante, rotundato; apertura obliqua, truncato-ovali, intus alba; plica colum. supera, levissima, fere nulla; perist. simplice, recto, margine colum. parum dilatato, sublibero.*

Long. 18, diam. $10\frac{1}{2}$ mill.

Sect. LAMINELLA.

5. *A. FERUSSACI*, Pfr. *A. testa sinistrorsa, perforata, turrita, solida, subruditer striata, nuda, coccinea; spira elongata, subconcavo-conica, apice fusca, obtusiuscula; sutura profunda, subcrenata; anfr. $7\frac{1}{2}$, superioribus planiusculis, sequentibus convexis, ultimo globoso, $\frac{1}{3}$ longitudinis non æquante; apertura diagonali, sinuato-semicirculari; columella distincte biplicata; plica supera obliqua, obtusa, altera compressa, illi subparallela; perist. nigro-limbato, margine externo recto, acuto, columellari dilatato, patente.*

Long. 24, diam. 11 mill.

Hab. Oahu.

6. *A. ALBIDA*, Pfr. *A. testa perforata, ovato-turrita, tenui, irregulariter striata, albida, epidermide pallide fulvescente strigulata; spira conica, apice acutiuscula; sutura levissime crenulata; anfr. 6, superis planis, sequentibus convexioribus, ultimo $\frac{2}{3}$ longitudinis vix æquante, rotundato, peripheria obsolete subangulato; apertura parum obliqua, sinuato-ovali; plica colum. mediana, laminæformi, obliqua; perist. simplice, recto, margine colum. breviter reflexo, sublibero.*

Long. 17, diam. $9\frac{1}{2}$ mill.

7. *A. INFLATA*, Pfr. *A. testa imperforata, conico-globosa, subruditer striata, albida, epidermide nigra infra suturam late fasciatim detrita obducta; spica inflata, in conum brevem, acutum terminata; anfr. $5\frac{1}{2}$, ultimis 3 perturgidis, ultimo spiram æquante; apertura obliqua, sinuato-ovali, intus alba; plica colum. subtransversa, alte ascendente, perist. recto, intus labiato, margine colum. dilatato, calloso, adnato.*

Long. $16\frac{1}{2}$, diam. 10 mill.

- β . *Minor, anfractu ultimo superne castaneo, basi pallido, vel coloribus obversis.*

8. *A. GONIOSTOMA*, Pfr. *A. testa rimata, dextrorsa, turrita, solida, oblique striatula, sub epidermide decidua fulva carneo-grisea; spira elongato-conica, apice acuta; sutura subcrenata; anfr. 8 planiusculis, penultimo convexiore, ultimo $\frac{1}{3}$ longitudinis vix superante, rotundato; apertura obliqua, elliptica, utrinque angulata; plica colum. subbasali, compressa, obliqua; perist. recto, basi producto, marginibus callo junctis, dextro intus sublabiato, columellari subdilatato, adnato.*

Long. 16, diam. $7\frac{2}{3}$ mill.

9. *A. EXTINCTA*, Pfr. *A. testa perforata, ovato-turrita, solida, striatula, cretacea; spira elongata, sursum attenuata, acuta; anfr. 7 vix convexiusculis, ultimo $\frac{1}{3}$ longitudinis non attingente, juxta perforationem subcompresso; apertura vix obliqua, rhombo-ovali, basi angulata; plica colum. compressa, fere a basi ascendente; perist. simplice, recto, marginibus callo crasso, subnodifero junctis, columellari dilatato, libero.*

Long. 16, diam. $7\frac{1}{2}$ mill.

Subfossilis lecta in insula Oahu.

10. *A. TRANSVERSALIS*, Pfr. *A. testa subrimata, ovato-conica, solida, confertim striata, lineis spiralibus impressis distantibus subdecussata, nigro-rufa, epidermide fulvo-grisea, guttatim et fasciatim interrupta, obducta; spira inflato-conica, acuta; anfr. 6 convexiusculis, ultimo $\frac{2}{3}$ longitudinis subæquante; apertura parum obliqua, sinuato-ovali; plica colum. lamellæformi, subbasali, fere transversa; perist. recto, intus albido vel roseo tenuiter labiato.*

Long. 12, diam. 6 mill.

11. *A. GROSSA*, Pfr. *A. testa imperf., dextr., conico-ovata, solida, subruditer striata, castanea; spira ovato-conica, apice acuta, nigricante; sutura pallida, crenulata; anfr. 7, summis planis, lævigatis, sequentibus sensim convexioribus, ultimo $\frac{2}{3}$ longitudinis subæquante; apertura obliqua, sinuato-semiovali, intus albida; plica colum. laminæformi, subtriangulari, fere transversa; perist. recto, acuto, marginibus callo albido junctis, dextro intus sublabiato, columellari adnato.*

Long. 23, diam. 11 mill.

12. *A. LUCTUOSA*, Pfr. *A. testa imperf., dextr., oblongo-conica, solida, striatula, nitidula, bicolore; spira inflato-conica, apice acuta; sutura crenulata; anfr. 6, summis 4 nigris, subplanis, penultimo pallido, convexiore, ultimo $\frac{2}{3}$ longitudinis subæquante, superne pallide fulvescente, infra peripheriam nigricante; apertura parum obliqua, sinuato-semiovali, intus alba; plica colum. laminæformi, subbasali, triangulari; perist. recto, acuto, nigrolimbato.*

Long. 16, diam. 8 mill.

13. *A. GRAYANA*, Pfr. *A. testa imperf., dextr., ovato-conica, solidula, nitidula, griseo-carnea, fasciis castaneis deorsum evanescentibus varie picta; spira inflato-conica, apice acuta, nigra; anfr. $6\frac{1}{2}$, superis planis, radiato-plicatis, penultimo convexo, ultimo spiram fere æquante, leviter striato; apertura subverticali, acuminato-ovali; plica colum. mediana, subtransversa, triangulari; perist. simplice, recto, margine colum. vix dilatato, adnato.*

Long. $21\frac{1}{2}$, diam. 12 mill.

14. *A. GLUTINOSA*, Pfr. *A. testa imperf., dextr., ovato-conica, solidula, arcuata, striatula, sub epidermide glutinosa, lutescente alba; spira subregulariter conica, apice obtusula; sutura levi,*

crenulata; anfr. 7 planiusculis, ultimo spira vix brevior, basi rotundato; apertura subverticali, sinuato-semiovali, intus alba; plica colum. compressa, torta, fere transversa; perist. recto, marginibus callo junctis, dextro sublabiato, columellari calloso, reflexo, adnato.

Long. $11\frac{1}{3}$, diam. 6 mill.

15. *A. UMBILICATA*, Pfr. *A. testa angustissime sed aperte umbilicata, dextr., ovato-conica, tenui, striata, opaca, fusca; spira concaviusculo-conica, apice acuta; anfr. 6 vix convexiusculis, ultimo spira paulo brevior, basi angulato; apertura elliptica, utrinque angulata; plica colum. compressa, profunda, subtransversa; perist. simplice, acuto, margine colum. subdilatato, omnino libero.*

Long. $10\frac{1}{2}$, diam. $5\frac{2}{3}$ mill.

Sect. ACHATINELLASTRUM.

16. *A. CLEMENTINA*, Pfr. *A. testa imperf., dextr., ovato-conica, solida, sublævigata, nitida, virenti-fusca, infra suturam lutescenti-fasciata; spira regulariter conica, obtusa; sutura castanea, impresso-marginata, crenulata; anfr. 6 superne turgidulis, ultimo $\frac{2}{3}$ longitudinis subæquante; apertura obliqua, truncato-auriformi, intus alba; plica colum. supera, mediocri, alba vel carnea; perist. recto, marginibus callo junctis, dextro substricto, intus incrassato-labiato, columellari dilatato, adnato.*

Long. 19, diam. 10 mill.

17. *A. CUNEUS*, Pfr. *A. testa imperf., sinistr., oblongo-turrita, solida, striatula, nitida, albida vel lutescente, fasciis angustis fuscis varie ornata; spira convexo-conica, apice alba, acutiuscula; sutura filo castaneo marginata; anfr. 6 planiusculis, ultimo $\frac{1}{3}$ longitudinis paulo superante; apertura fere diagonalis, semiovali, intus alba; plica colum. mediocri, oblique subtorta; perist. acuto, intus labiato, margine columellari adnato.*

Long. 18, diam. 8 mill.

Sect. AMASTRA.

18. *A. IRREGULARIS*, Pfr. *A. testa imperf., dextr., tenuiuscula, longitudinaliter confertim plicata, nigricante, interdum albidofasciata; spira irregulari, conica, obtusa; sutura subcrenata; anfr. $5\frac{1}{2}$ celeriter accrescentibus, convexis, penultimo gibbo, ultimo $\frac{2}{3}$ longitudinis subæquante, infra medium læviore, interdum corneo; apertura vix obliqua, ovali-oblonga; plica colum. supera, parum prominente; perist. simplice, margine dextro antorsum subarcuato, columellari dilatato, adnato.*

Long. 9, diam. $4\frac{1}{3}$ mill.

Sect. LEPTACHATINA.

19. *A. DIMIDIATA*, Pfr. *A. testa imperf., ovato-conica, solidula, striatula, nitida, superne nigro-castanea; spira convexo-conica,*

apice obtusula; sutura levi, subcrenulata; anfr. $6\frac{1}{2}$ planiusculis, ultimo spira paulo brevior, a medio ad basin pallide corneo; apertura vix obliqua, sinuato-ovali, intus concolore; plica colum. callosa, subobliqua; perist. albo, recto, intus labiato, margine colum. angusto, adnato.

Long. 11, diam. $5\frac{1}{2}$ mill.

β. *Anfractu ultimo unicolore castaneo, spira pallidior.*

20. A. SEMICOSTATA, Pfr. *A. testa subumbilicata, dextr., turrita, solidula, confertim costata, nitida, castaneo-cornea; spira elongata, convexo-conica, obtusula; sutura impressa; anfr. $8\frac{1}{2}$ parum convexis, ultimo $\frac{2}{5}$ longitudinis vix æquante, infra medium lævigato, pallido, attenuato, basi saccato-compressa; apertura parum obliqua sinuato-oblonga; plica colum. lata, obliqua, alba; perist. simplice, recto, margine colum. reflexo, omnino patente.*

Long. $9\frac{1}{2}$, diam. 4 mill.

21. A. TERES, Pfr. *A. testa imperf., dextr., cylindraco-oblonga, tenuiuscula, leviter striatula, nitidissima, cornea; spira sensim attenuata, apice obtusula; sutura crenulata, saturate castanea, in anfractu ultimo impresso-marginata; anfr. 7 planiusculis, ultimo $\frac{1}{3}$ longitudinis paulo superante, antice subascendente; apertura subobliqua, sinuato-semiovali; plica colum. subcallosa, torta, obliqua; perist. simplice, recto.*

Long. 10, diam. 4 mill.

22. A. ORYZA, Pfr. *A. testa subrimata, fusiformi, striatula, cornea (!); spira convexo-conica, apice obtusula; sutura levi, simplice; anfr. 7 vix convexiusculis, ultimo $\frac{2}{3}$ longitudinis subæquante, basi attenuato; apertura vix obliqua, acuminato-ovali; plica colum. subcompressa, acuta; perist. simplice, recto, margine colum. subreflexo, adnato.*

Long. 8, diam. $3\frac{3}{4}$ mill.

Hab. Subfossilis in insula Oahu.

23. A. MARGARITA, Pfr. *A. testa subrimata, dextr., ovato-oblonga, tenui, striatula, pellucida, pallide cornea; spira ovato-conica, obtusula; sutura submarginata; anfr. 6, superis planiusculis, penultimo convexiore, ultimo $\frac{3}{4}$ longitudinis subæquante, basi subattenuato; apertura subverticali, elliptico-ovali, basi subangulata; plica colum. mediana, parvula, dentiformi; perist. simplice, recto, albido-limbato.*

Long. $6\frac{1}{3}$, diam. 3 mill.

Sect. AURICULELLA.

To this section may be referred:—

24. ACHATINELLA LURIDA, Pfr. (*Tornatellina castanea*, Pfr. Mon.)

25. ACHATINELLA OBELISCUS, Pfr. (*Balea Newcombi*, Pfr. Mon.)

5. DESCRIPTIONS OF SIXTEEN NEW SPECIES OF ACHATINELLA,
FROM MR. CUMING'S COLLECTION, COLLECTED BY
DR. NEWCOMB IN THE SANDWICH ISLANDS.
BY DR. L. PFEIFFER.

1. *A. DWIGHTI*, Newc. *A. testa perforata, sinistrorsa, solida, oblongo-conica, ruditer striata, striis confertis spiralibus decussata, fusculo et albo variegata s. marmorata; spira conica, obtusula; anfr. $6\frac{1}{2}$ planiusculis, ultimo spira paulo brevior; apertura obliqua, obauriformi; plica columellari obliqua, levi; perist. albo, intus crasse labiato, margine externo expanso, columellari dilatato, calloso, sublibero.*

Long. 24, diam. $11\frac{1}{2}$ mill.

Hab. Molokai.

2. *A. MANIENSIS*, Newc. *A. testa subperforata, ovato-oblonga, tenuiuscula, longitudinaliter irregulariter striata, sub lente striis spiralibus confertissimis decussata, haud nitente, albida, strigis fusco-corneis serratis variegata; spira regulariter conica, apice obtusa; sutura levi, vix submarginata; anfr. $5\frac{1}{2}$ planiusculis, ultimo $2\frac{1}{2}$ longitudinis subæquante, basi subattenuato; apertura parum obliqua, oblonga; plica colum. obsoleta; perist. simplice, recto, fusco-limbato, margine colum. fornicatim reflexo, subadnato.*

Long. 13, diam. 6 mill.

Hab. Mani.

3. *A. TETRAO*, Newc. *A. testa subperforata, sinistrorsa, globoso-conica, pallida, infra suturam late rubro-fasciata, undique epidermide virenti-fusca dense reticulata; spira conica, acutiuscula; anfract. $6\frac{1}{2}$, summis subplanis, reliquis perconvexis, ultimo globoso, spira paulo brevior; apertura obliqua, semiovali, intus alba; plica colum. laminæformi, subobliqua; perist. simplice, recto, acuto, margine columellari roseo, sublibero.*

Long. 16, diam. $8\frac{1}{2}$ mill.

Hab. Ranaï.

4. *A. REMYI*, Newc. *A. testa subimperfecta, sinistrorsa, oblonga-conica, plicatula, rubella, lineis fulguratis fuscis amœne variegata; spira elongata, apice nigricante, acuta; anfr. 7 modice convexis, ultimo $\frac{1}{3}$ longitudinis vix superante; apertura fere diagonali, sinuato-ovali, intus purpurascens; plica colum. parum prominente, subduplicata; perist. simplice, recto, acuto, margine colum. rubro, subadnato.*

Long. $17\frac{1}{2}$, diam. 8 mill.

Hab. Hawaii.

5. *A. HUMILIS*, Newc. *A. testa subperforata, oblongo-conica, solidula, subruditer striata, griseo-carnea; epidermide nigra fere omnino obducta; spira convexo-conica, apice nigra, acuta; anfr. $6\frac{1}{2}$ infra suturam turgidis, medianis pallide reticulatis, ultimo $\frac{1}{3}$ longitudinis paulo superante; apertura parum obliqua, sinuato*

ovali, intus livide purpurascente; plica colum. mediocri, compressa, obliqua; perist. simplice, recto, acuto, margine colum. dilatato, reflexo, subadnato.

Long. 18, diam. $8\frac{2}{3}$ mill.

Hab. Molokai.

6. A. PETRICOLA, Newc. *A. testa subperforata, dextrorsa, ovato-conica, tenuiuscula, striatula, fusca; spira convexo-conica, acutiuscula; sutura levi; anfr. $5\frac{1}{2}$ convexiusculis, ultimo $\frac{2}{3}$ longitudinis subæquante; apertura parum obliqua, sinuato-ovali, intus concolore; plica colum. tenui, compressissima, parum obliqua; perist. simplice, recto, intus albo limbato, margine colum. reflexo, subpatente.*

Long. $9\frac{1}{2}$, diam. 5 mill.

Hab. Modonai.

7. A. DUNKERI, Cuming, MSS. *A. testa imperforata, oblongo-turrita, solidiuscula, lævigata, pallide carnea, strigis confertis fusculis ornata, lineis 1 vel 2 nigris sæpe cincta; spira turrita, apice acutiuscula, alba; sutura leviter marginata, pallida; anfr. $6\frac{1}{2}$ vix convexiusculis, ultimo $\frac{2}{3}$ longitudinis fere æquante; apertura obliqua, truncato-auriformi; plica colum. alba, valida, dentiformi, subsulcata; perist. recto, acuto, pone limbum fuscum albo labiato.*

Long. 24, diam. 11 mill.

8. A. FULVA, Newc. *A. testa subperforata, turrito-conica, solida, striatula et sub lente subtilissime decussata, nitida, luteo-fulva; spira subconcavo-conica, apice alba, acuta; sutura filomarginata; anfr. $6\frac{1}{2}$, superis planis, sequentibus convexis, ultimo $\frac{1}{3}$ longitudinis paulo superante, infra suturam turgido, basi saccato; apertura obliqua, subtetragono-auriformi; plica colum. supera, nodiformi, aurantiaca; perist. recto, intus labiato, margine dextro substricto, columellari dilatato, fornicato-reflexo.*

Long. $18\frac{1}{2}$, diam. 9 mill.

9. A. OVIFORMIS, Newc. *A. testa sinistrorsa, imperforata, ovato-conica, solida, striatula, sub epidermide nitida, olivacea alba, fusco 1-3 fasciata; spira convexo-conica, apice acutiuscula, alba; sutura profunde marginata; anfr. 5 vix convexiusculis, ultimo $\frac{2}{3}$ longitudinis subæquante; apertura fere diagonali, auriformi; plica colum. mediocri, dentiformi, rubella; perist. recto, marginibus callo tenui junctis, externo intus labiato, columellari dilatato, adnato.*

Long. 17, diam. $8\frac{2}{3}$ mill.

Hab. Oahu.

10. A. UNDULATA, Newc. *A. testa sinistrorsa (rarius dextrorsa), imperforata, solida, striatula, nitida, fulva, strigis undulatis fuscis vel nigro-castaneis picta; spira conica, apice acuta, alba; sutura marginata; anfr. $5\frac{1}{2}$ modice convexis, ultimo $\frac{2}{3}$ longitudinis subæquante, latere subcompresso; apertura fere diagonali, obauri-*

formi, intus alba; plica colum. valida, dentiformi; perist. simplice, recto, fusco-limbato.

Long. 16, diam. 9 mill.

Hab. Oahu.

11. *A. NIGRA*, Newc. *A. testa imperforata, globosa-conica, tenuiuscula, confertim striata, albida, epidermide nigricante fere omnino obducta; spira concavo-conica, apice acuta; anfr. 6, superis planis, penultimo convexo. ultimo spiram subæquante, inflato, plicis spiralibus irregulariter minuto; apertura obliqua, sinuato-ovali; plica colum. subtransversa, linguæformi; perist. recto, acuto, intus tenuiter labiato, marginibus callo tenui junctis, columellari obsoleto.*

Long. 17, diam. $10\frac{1}{2}$ mill.

Hab. Mani.

12. *A. PULLA*, Newc. *A. testa imperforata, acuminato-ovata, solidula, distincte striata, lutescente, epidermide fusca varie fasciata, vel præter fasciam pallidam omnino obducta; spira inflata, versus apicem acutum attenuata; sutura pallida; anfr. 6, primis planis, penultimo modice convexo, ultimo $\frac{3}{4}$ longitudinis subæquante; apertura parum obliqua, sinuato-ovali; plica colum. dentiformi, subacuta, alba; perist. simplice, recto, margine columellari adnato.*

Long. 11, diam. 6 mill.

Hab. Ranaï.

13. *A. SUCCINCTA*, Newc. *A. testa imperforata, ovata, tenuiuscula, dense striolata, pellucida, cornea, linea suturali et fascia lata fuscescentibus ornata; spira ventroso-conica, obtusula; anfr. $6\frac{1}{2}$ vix convexiusculis, ultimo $\frac{3}{4}$ longitudinis subæquante, basi lævigato; apertura vix obliqua, sinuato-ovali; plica colum. obliqua, compressa, violaceo-submarginata; perist. simplice, recto, intus tenuiter albido-labiato.*

Long. 13, diam. $6\frac{1}{2}$ mill.

Hab. Wahi.

14. *A. ELEVATA*, Newc. *A. testa imperforata, oblongo-turrita, solidula, longitudinaliter (ad suturam distinctius) striata, nitida, corneo-lutea, castaneo-bifasciata; spira turrita, apice obtusa; anfr. 8 planiusculis, ultimo $\frac{1}{3}$ longitudinis subæquante, basi subattenuato; apertura verticali, elliptico-ovali; plica colum. compressa, obliqua, pallide lilacea; perist. recto, obtuso, margine dextro antrosum subdilatato.*

Long. $11\frac{1}{3}$, diam. $4\frac{1}{3}$ mill.

15. *A. OBTUSA*, Newc. *A. testa imperforata, oblonga, sublævigata, nitida, pellucida, castaneo-cornea; spira cylindrico-turrita, apice obtusa; sutura submarginata; anfr. $6\frac{1}{2}$ subplanulatis, ultimo $\frac{2}{3}$ longitudinis subæquante; apertura verticali, acuminato-ovali, basi subangulata; plica colum. levi, obliqua; perist. simplice, recto, pallide limbato, marginibus callo tenui junctis.*

Long. 10, diam. 4 mill.

16. *A. APICATA*, Newc. *A. testa imperforata, globoso-conica, solida, levigata, nigricante vel fusca, pallide irregulariter radiata et subfasciata; spica conica, apice acutiuscula, cinnamomea vel hepatica; anfr. 6, superis planis, penultimo convexo, ultimo globoso, $\frac{2}{3}$ longitudinis vix æquante; apertura obliqua, subtetragono-ovali; plica colum. supera, dentiformi, lilacea; perist. recto, intus lilaceo-labiato, margine colum. perdilatato, incrassato, adnato.*
Long. $18\frac{1}{2}$, diam. 12 mill.

6. DESCRIPTIONS OF FIVE NEW SPECIES OF TERRESTRIAL MOLLUSCA, FROM THE COLLECTION OF H. CUMING, ESQ.

BY DR. L. PFEIFFER.

1. *LEPTOPOMA LOWI*, Pfr. *L. testa perforata, globoso-turbinata, tenui, sub lente confertissime spiraliter striata, pellucida, fulvo-cornea, ad suturam fascia alba subopaca ornata; spira turbinata, acuta; anfr. 5 convexis, medianis liris nonnullis obsoletis notatis, ultimo rotundato; apertura obliqua, subcirculari, intus margaritaceu; perist. albo, subrectangule expanso, marginibus callo tenuissimo junctis, columellari superne subexciso.*

Diam. maj. 14, min. 12, alt. 11 mill.

β. *Minor, anfr. ultimo conico-variegato, medio albido-subfasciato.*

Hab. Isle of Labuan (Mr. Hugh Low).

2. *PLECOTREMA BICOLOR*, Pfr. *Pl. testa subumbilicata, ovato-conica, solida, spiraliter confertim sulcata, rugulis longitudinalibus irregulariter tuberculosa, vix nitidula, brunnea, pallide fasciata; spira convexiusculo-conica, apice pallida, acuta; anfr. 9 vix convexiusculis, ultimo spira paulo longiore, superne turgidulo, basi attenuato, circa perforationem non perviam compresso, antice crista elevata munito et contracto; apertura vix obliqua, oblonga; plicis parietalibus 2 parallelis, inferiore extus breviter bicruri; plica colum. parvula, transversa; perist. vix patulo, marginibus callo junctis, dextro bidenticulato.*

Long. 6, diam. $3\frac{1}{2}$ mill.

Hab. Port Jackson (Mr. Strange).

3. *CLAUSILIA BLANDIANA*, Pfr. *Cl. testa non rimata, fusiformi-turrita, solidula, oblique obsolete striata, diaphana, cornea; spira turrita, obtusa; sutura levi, rufo marginata; anfr. 9 regulariter accrescentibus, parum convexis, ultimo angustato, deorsum protracto, basi rotundato, antice rufescente, distinctius striato; apertura oblique piriformi-circulari; lamellis convergentibus, superiore valida, marginali, altera conspicua, tenuiore; lunella distincta, angusta; plica palatali 1 supera, ad lamellam superiorem producta; subcolumellari inconspicua; perist. albo, tenui, expanso et reflexiusculo, margine sinistro angustiore.*

Long. 19, diam. $4\frac{1}{2}$ mill.

Hab. Santa Fé de Bogota, Columbia.

4. *ACHATINA SCULPTA*, Pfr. *A. testa conico-oblonga, solida, plicis longitudinalibus subarcuatis validis sculpta, straminea; spira regulariter attenuata, obtusa; anfr. 8 planiusculis, ultimo $\frac{1}{3}$ longitudinis paulo superante, antice subascendente; apertura subverticali, sinuato-ovali; columella breviter arcuata, oblique truncata; perist. simplice, recto, margine dextro antrorsum subarcuato, columellari tenuiter calloso.*

Long. $8\frac{1}{2}$, diam. $3\frac{1}{8}$ mill.

Hab. Oahu, Sandwich Islands (Mr. Frick).

5. *ACHATINA COMORENSIS*, Pfr. *A. testa ovato-turrita, solida, leviter et irregulariter striatula, sub epidermide decidua, fulva alba; spira elongata, apice obtusula; anfr. 7, superis subplanis, penultimo conveziore, ultimo $\frac{1}{3}$ longitudinis paulo superante, basi subattenuato; apertura subverticali, elliptica, basi subcanaliculata, columella ad basin aperturæ protracta, oblique subtruncata; perist. obtuso, marginibus callo crassiusculo junctis.*

Long. 34, diam. 13 mill.

Hab. Comoro Islands.

Nov. 27th, 1855.

Dr. Gray, F.R.S., in the Chair.

The following papers were read :—

1. ON A NEW SPECIES OF SOMATERIA, AND THE FEMALE OF *LAMPRONETTA FISCHERI*, BRANDT. BY G. R. GRAY, F.L.S. ETC.

(Aves, Pl. CVII.—CVIII.)

I have the pleasure of exhibiting Mr. Adams's drawing of a new species of *Somateria*, of which several specimens were collected by that gentleman during the voyage of H.M.S. *Enterprise*. The large drawing exhibits the male and female, while the lesser drawing contains, in addition to the two sexes, the male in change. It will at once be seen that this species is very similar to *Somateria mollissima*, but is easily distinguished from it by being larger, and by a black mark on the throat, which is like the one found on that part of the *Somateria spectabilis*; and by the white longitudinal mark on the top of the head being narrower and thereby less conspicuous than that found on the head of *Somateria mollissima*; while the yellow on the sides of the head, which is tinged with green, extends below the black towards the eyes; the black is less rounded posteriorly. The bill of the male also differs by being orange-yellow with a white nail; while that of the female is dusky green as in the com-