Original Articles.

DOUBLE CONGENITAL DISPLACEMENT OF THE HIP; DESCRIPTION OF A CASE WITH TREATMENT RESULTING IN CURE.

(With Plates.)

BY BUCKMINSTER BROWN, M.D.,

Consulting Surgeon of the House of the Good Samaritan, etc.

WHEN in Paris during the winter of 1845 and 1846 I was much interested in observing, among the cases in the wards of M. Guérin at the Hôpital des Enfants Malades, three or four of congenital dislocation or displacement of the hip. I was not able to follow them a sufficient length of time to learn the degree of success attending the treatment. M. Guérin was sanguine that the result would be favorable.

In 1843 "l'ancien conseil général des hôpitaux et hospices civils de Paris" appointed a commission to follow, for a time not less than one year, the treatment pursued by M. le docteur Jules Guérin in the Children's Hospital of Paris in cases classed under the head of Orthopedic Surgery. The commission was instructed to report upon the cases so followed and examined. It was composed of MM. Rayer, Serres, Louis, Breschet, Jobert, and Blandin. The council also appointed M. Orfila as president.

In 1848 the report of this committee was published.¹ The fourth class of cases reported upon was congenital luxations of the femur. Of this description of cases there were five submitted to the commission for observation and examination. Of these five, four were examples of single, and one of double, dislocation. The first two patients presented were girls, one six and the other seven years of age. In each, one joint only was affected. These cases were under the notice of the reporters from December 3, 1843, to December 28, 1845. The result was not a permanent replacement of the head of the femur in either of the cases. But, much to the surprise of the committee, there was found in each case an elongation of the long bones of the leg to the extent of two centimeters. In this way compensation was established for the shortening produced by the luxation.

Such was the result of the examination of the committee and testified to in their report.

Upon the point above referred to I will quote a personal observation made about the same period that this committee was pursuing its investigations:—

"There is yet another class of cases. In certain children the resistance to the means employed for producing a descent of the head of the femur is so great as to render these efforts wholly ineffectual. What is very curious in these cases is that there is an elongation of the bone itself. Thus we have still the signs of dislocation on examining the hip-joint, but on comparing the two limbs we find them of the same length."²

The third case reported upon, after fourteen

months' treatment promised success. Most unfortunately, however, the patient had typhoid fever attended by severe cerebral symptoms, and by general convulsions and contraction of the muscles of the trunk and limbs. The displacement returned worse than it was originally. After extension, the trochanter was again brought into position, and hopes were entertained of a cure. Without known cause the luxation returned, and at the date of the report the case was still under treatment. Of the two other cases one was not treated, on account of the occurrence of some other malady.

The fifth subject, being the double displacement, was under treatment at the time of the report.

Thus we have five cases of this malformation presented by M. Guérin to an especially appointed commission. In two of these, as I have stated, the treatment was terminated at the end of two years. In neither of them was the malposition remedied. The third and fifth were still under treatment, and in the fourth, as before mentioned, no attempt was made at restoration.

Contributions have been made from English, French, and Italian writers to elucidate the pathology and aetiology of this class of cases.

Dr. Carnochan, of New York, has written a valuable monograph upon this subject, in which he describes and exemplifies, with plates, the anatomical relations of the displaced bones and the changes which occur in the pelvis after a lapse of years. He likewise gives plates, copied from a report of M. Pravaz, representing apparatus for extension and for exercising the joint, which, judging from the diagrams, must be extremely complicated, unwieldy, and expensive. These, however, Dr. Carnochan has not put into any practical use so far as can be gathered from his work. No case illustrative of treatment and cure is related by him.

Brodhurst has had a number of these cases under his care. He advises in all cases, except in children under two years of age, the division of "all the muscles" which are inserted into and about the trochanters, especially the glutei and the rotators.

This author relates a case of single displacement in which he did this operation, and the patient, with the assistance of a thickened sole, walked well. Similar operations were performed by M. Guérin in the cases to which I have previously referred.

Dr. Sayre recommends his instrument for extension of the hip during locomotion and also Darrach's wheel-crutch.

"M. Pravaz, Jr., has put on record the case of a girl, aged seven years and a half, in which he claims complete success in the reduction of a dislocation of this nature by a prolonged treatment."³

In this instance there was unquestionably a well-defined acetabulum, as, when the head was reduced, it went in with a "distinct snap." In this respect the case of M. Pravaz essentially differs from the one hereafter to be described in this paper. So important a fact as the existence of a pronounced socket places the parts to be acted upon in an entirely dissimilar condition. Dr. Carnochan likewise alludes to the report upon this case of M. Pravaz.

¹ Rapport Adressé à Monsieur le Délégué du Gouvernement Provisoire sur les Traitements Orthopédiques de M. le docteur Jules Guérin à l'hôpital des Enfants Pendant les Années 1843, 1844, et 1845, par une commission composée de MM. Blandin, P. Dubois, Jobert, Louis, Rayer, et Serres; Président, M. Orfila. Paris, 1848.

² Orthopedic Surgery in Europe. Letter dated Paris, May 19, 1846. Buckminster Brown, M.D. Boston Medical and Surgical Journal, July 1, 1846.

³ The Surgical Treatment of the Diseases of Infancy and Childhood, p. 219. T. Holmes.

Mr. H. A. Reeves, of London, in a work just published (1885) says: "My opinion is that in a large majority of cases anything other than palliative measures, such as the use of appropriate apparatus for grasping the trochanters and keeping them as far as possible in place, is of little use. Pravaz claimed to have made cures by the use of his special apparatus, but Bouvier and others have fairly shown that these were not permanent cures."⁴

Dr. James Knight, in his recent work on Orthopaedia, relates a case where "after two years' treatment the patient was enabled to walk independently of support to body or limbs." Further particulars are not given; as whether after treatment the position of the femoral head was normal or otherwise, or if there was a cotyloid cavity before or after treatment. So far as my knowledge extends these patients are usually able to walk.

Dupuytren wrote a memoir on this subject and expressed his opinion that there was no remedy, and not even a palliative for these displacements.⁵

Mr. Thomas Bryant, in his standard work on Surgery, in writing on the treatment of cases of this malformation, says: "Nothing can be recommended with the view of cure, for art cannot supply a natural deficiency or make up for a defect of structure in the bones of the joint. . . . Mechanical appliances are to be condemned as useless, if not worse. They have been employed on a wrong principle, or rather on a want of due appreciation of the conditions of parts, and therefore with only a vague hope that they may do good. These observations apply as much to the use of an extension apparatus as to operative interference, although with more force to the latter."⁶

Coxectomy has been performed for relief in this affection.

I quote from the *Medical News* for October, 1884: "Congenital dislocations of the head of the femur, especially when they have existed for some time, are notoriously unamenable to permanent relief, the means resorted to for this purpose being as a rule directed to limiting the tendency of the bone to further displacement by wearing a strap of webbing around the hips.

"At the recent *Versammlung der Deutscher Naturforscher und Aerzte*, in Magdeburg, Dr. Heusner, of Burmen, demonstrated that such cases are open to more radical measures, and placed on record the first excision of the hip for congenital dislocation. The operation was done for excessive suffering in the left joint of a girl twenty years of age. The ligamentum teres was absent, the head of the femur was about as large as that of a child of six years, ovoidal in shape, and much flattened, and its surface was uneven and the neck of the bone formed an obtuse angle with the shaft. The acetabulum was fully developed and of normal size, although its rim was somewhat flattened and polished above and below.

"The head, neck, and nearly an inch of the shaft of the femur having been excised, the acetabulum

was made deeper with the chisel and the parts brought into apposition. At the expiration of eleven weeks, with the aid of a cane the patient was able to walk without fatigue for half an hour, the pain was relieved, and there was every prospect of future good use of the limb.

"The case briefly narrated shows that the theoretical objection to operative interference in congenital dislocations of the hip, based upon the absence of a well-formed cotyloid cavity, is not well founded. Even if the acetabulum were superficial or deficient a cavity of sufficient size to accommodate the divided shaft of the femur could be chiseled out of the ilium." Operations similar to the one above described have been performed by Margary, of Turin, E. Rose, and C. Reyher.

The behavior of the hip-joint in other circumstances, as in traumatic unreduced dislocations, etc., would unquestionably lead to the belief that a cure could be effected by following the teachings, and taking advantage of the hints thus obtained, and by the combination of physiological and mechanical means, without having recourse to the heroic measures above described. My confidence, however, in the success of treatment has been so slight, and the circumstances of the patients who have presented themselves have been such, as not to warrant me in testing a course of procedure tedious and wearing at the best, whether operative or mechanical, and of which the history of cases, where any minute record has been given, has not been for the most part encouraging.

Connate malposition of the head of the thigh-bone is comparatively a rare affection and occurs most frequently double. Dupuytren says: "Simultaneous displacement of both femurs is observed in most individuals affected." Carnochan also states, as the result of his researches, that dislocations on both sides are the most frequent.

The prominent symptoms in those cases where both hips are displaced are, first, the peculiar waddling walk, and second, the extreme spinal incurvation.

The swaying walk is due mainly to the extent of surface traversed by the heads of the femurs. The acetabula being absent or deficient, the capsular and round ligaments, if present,⁷ are so stretched as to have no influence in limiting the motion of the femoral heads, which are held alone by the muscles and fascia.

Thus, in raising the foot in walking, the weight of the limb extends the muscles connected with the neck of the femur and trochanter, and the head of the thigh-bone sinks one or more inches. In placing the foot upon the floor the weight of the body causes it (the head) to slide upward again as far as the insertion of the muscles will permit. In this way a smooth sac is formed either by the capsular ligament, or if this has been ruptured, as sometimes occurs, immediately beneath the muscles on the external surface of the ilium. This is probably bursal-lined, and from the difficulty of producing adhesion between its sides is an additional impediment to permanent restoration.

⁴Bodily Deformities and their Treatment, p. 308. By Henry Albert Reeves, F.R.C.S.E. Philadelphia: P. Blakiston, Son & Co. 1885.

⁵Mémoire sur le déplacement originel ou congénital de la tête du fémur.

⁶A Manual for the Practice of Surgery, p. 769. By Thomas Bryant, F.R.C.S. Second American Edition.

⁷"According to Cruveilhier, this latter ligament is often undeveloped and sometimes wanting." The Mechanism of Dislocation and Fracture of the Hip, p. 21. By Henry J. Bigelow, M.D. It may be that in some cases of congenital displacement of the hip such a state of things exists.

Plate 2.

Plate 1.

Plate 3.

Plate 4

Plate 6. June, 1884.

Plate 5.

Lewis & Co., Photog.

Plate 7.

Andrew P. Smith, Phil.

Plate 8.

The lordosis is simply the natural attempt to gain an equilibrium. The upper part of the body being thrown from the median line by the displacement backward and upward of the femoral heads, a counterpoise is necessitated where only it can take place, and an unavoidable anterior lumbar curve is the result.

The following records of an interesting and rare case are written in the hope of their being of service to members of the profession, who, in undertaking a similar case, may be glad to have the narration of such an experience to turn to, to aid them in their labors. With this in view the apparatus, which was found fully to answer its purpose, has been accurately photographed and described, and every important step toward restoration carefully noted. The apparatus used was designed as the objects to be accomplished and the exigencies of the treatment presented themselves.

On the fourth of April, 1882, a little girl four years of age was brought to me to be treated for what was thought to be an unusual example of spinal curvature. On examination, I found a double hip displacement.

As on former similar occasions and for reasons already mentioned, I decided against active treatment. The state of the case was clearly placed before the parents, accompanied by the statement that an attempt at restoration would involve a course of treatment not unattended with risk, at least in some of its phases, necessarily prolonged, and so extremely uncertain in its results that I did not feel justified in advising its adoption.

The parents of this child, however, were not to be dissuaded from making an attempt to relieve their daughter from so distressing a condition.

The thought of its continuance with its blighting influence through life was unbearable. After weeks for mature deliberation they decided that the trial should be made, if I were willing to undertake the treatment. The condition of the hips at this time was as follows: No trace of a cotyloid cavity could be discovered. When the patient was recumbent, if the shoulders were firmly held by an assistant, traction on the legs would readily bring the heads of the femurs into their normal position, and they could as easily be pushed back again, or they would slip upward by the natural contraction of the muscles. There was no spasmodic or spastic contraction. The muscles acted powerfully, but not abnormally. There was no such action of a muscle or group of muscles as would demand or warrant the use of the knife. I have said there was strong muscular action, and after prolonged extension, if the hand was laid upon the nates or pelvis, it was curious to feel the fibrous, fascicular, wormlike movement by which the limb was slowly drawn upward over the ilium. This was the simple, natural, muscular contractility brought into play after resistance had been removed. The head glided upward over a smooth surface. The walk was the extreme of that sideward movement which is expressed by the term waddle. It was a much exaggerated waddle. Plate 1 represents the displacement viewed laterally. The great trochanter is seen as a rounded prominence high up on the ilium, bordering on the crest. The muscles are stretched

and bagged up by the pressure of the weight of the body on their tissues. The excessive lordosis is conspicuous, showing the obtuse angle at the junction of the lumbar vertebrae and sacrum.

The consequent prominent, overhanging abdomen and compensatory bend of the knees are well seen in this side view. Likewise the shortening of the stature is evident. Plate 2 represents the appearance of the child seen anteriorly. It depicts the height of the trochanters on a level with the upper rims of the ilia, and the projection in front. The splayfoot, or valgoid tendency, the usual accompaniment of this condition, is also well marked.⁸

The child was brought to Boston, and on the thirtieth day of December, 1882, she was placed upon the bed. I will here premise that for many of the dates and for much of the minute history of the case I am indebted to a diary kept by the mother of the patient.

Obviously my treatment must be divided into distinct stages. First, it was requisite to relax and extend the large mass of glutei and other pelvic muscles.

It has been thought by some surgeons, and perhaps with justice, that continuous, warm, moist applications have an effect in relaxing and weakening the muscular fibre. How deep the influence of such applications extends from the surface may be a question. In order to omit no measure, however apparently trifling, which might be supposed to assist to the end in view, I commenced with warm poultices. These enveloped the pelvis and hips, and were renewed twice a day for one week.

January 6, 1883, a firm leather belt three inches wide, padded and lined with chamois-skin, was buckled round the hips. To the upper edge of this belt four long straps were attached, extending from just above the anterior superior spinous process of the ilium on each side in front and from just above the sacro-iliac synchondrosis on each side at the back. These long straps were buckled to the rail at the head of the iron bedstead. Two perineal straps were fastened by four buckles to the lower edge of this belt, in such a manner as to bring the soft, padded portion into the groove of the groin. This gave a perfectly firm, unyielding counter-extension.

Direct extension was made by means of stiff leather bands, three and a half inches wide, padded and lined, encircling the legs above the knees and above the ankles, accurately fitted. To the lower edge of the bands leather loops three or four inches long were fastened, one on each side. A cord to which weights were suspended passed over parallel pulleys at the foot and on a level with the bed, and were tied to the loops. See *a, a*, plate 3, and *a, a*, plate 4. The use of the bands alternated: sometimes those above the knees, and sometimes those above the ankles, were employed. In this way the pressure around the limbs could be frequently varied, thereby affording great relief.

It is important to be thus minute in relation to

⁸ Plates Nos. 1 and 2 are from photographs of a former patient, a complete facsimile of the one I am now describing. On comparing these photographs with the subject of this paper, at my first examination, April 4, 1882, they were deemed, both by the parents and myself, to be exact representations of the case before us and no others were considered necessary.

the various appliances, as all expectation of success depends upon their being rendered as little painful and irksome as possible.

The knee-bands were preferably employed. By them the extension drew directly upon the pelvic muscles. The object here being to tire out and to permanently stretch these organs, and thereby remove one important obstacle to restoration. Weights weighing three pounds were used at first. These were afterward increased. After a few days the head of the bedstead was raised by means of wooden blocks, eight inches high. The child was thus, to a certain extent, suspended by the perineal straps.

In the course of ten or twelve days sufficient relaxation of the muscles was produced to allow of easy manipulation of the bones, and each morning they were placed in their natural position, where they were maintained by the weights for some hours. On the following morning, however, they were invariably found displaced. About this time, on two or three occasions when drawing upon and rotating the left femur, a sensation was communicated to the hand as if the head of the bone was passing over an inequality. Whether this sensation was due to its slipping over a segment of the rim of an undeveloped acetabulum, or whether it arose from its passing over a portion of an elongated round ligament was a question which was difficult to decide. I was disposed to think that the former supposition was correct. Dr. H. H. A. Beach, when examining the patient at this time, also detected this slight jar on rotating the left femur, previous to the upward movement by which it passed on to the dorsum illi, and was inclined to the opinion that such a segment as above suggested did exist.⁹

This inequality was at no time observed on the other side, and even here could not always be discovered.

If a detached remnant of the cotyloid ligament did exist, it was so imperfect as to offer no resistance to the free movements of the head of the femur, and escaped detection at the earlier examinations.

About the middle of February it was found that the tendency to slip upward was greatly diminished, and that days elapsed without this occurrence taking place.

Accurate measurements, together with the hint derived from the slight section of a ridge on the left side just referred to, were the guides for placing the heads of the femurs and electing the spot for the formation of the hoped-for sockets.

As an assistance to this end I resorted to a position of the limb which I had seen employed by Dr. H. J. Bigelow in a case of acquired hip dislocation which in facility of displacement resembled the one we are now considering. This position was likewise resorted to in a case which is narrated in his work on "The Hip," and is described on page 56.

The case was one of traumatic dislocation of the hip of eight months' standing. The limb was reduced by flexion, abduction, and eversion, but was readily displaced. Dr. Bigelow says:—

"A very slight inversion sufficed to reproduce the dislocation: in fact, the limb could not be

trusted to itself. After the bone had thus repeatedly slipped out, the patient was placed in bed on her back, and the dislocation again reduced by flexion, abduction, and eversion, which brought the flexed thigh and knee down to the mattress on their outer side. The knee was then tied to the bedstead in this position by a towel, and the foot secured to the knee of the sound side until the socket should be excavated by absorption."¹⁰

To return to my patient. The femoral heads now being in normal position the thighs were flexed to nearly right angles with the body and then spread in the manner described by Dr. Bigelow, and the knees tied to the edge of the bedstead on each side and the feet secured to the opposite side. The heads of both thigh-bones were thus brought to bear forcibly on the pelvis at the points selected, trusting that absorption might be produced as a commencement of the desired depression.

This position was painful and could be borne but two or three hours in the course of the day. The strain on the anterior crural and other nerves in the neighborhood affected the whole nervous system. After the ties were removed from the knees the legs were weighted as before and so remained the rest of the twenty-four hours. The traction bands were alternated, as previously mentioned, from those that encircled above the knees and those encircling the ankles.

Two long narrow strips of wood were now attached to the foot of the bedstead a few inches from the corners. (See *b, b*, plate 3, and *b, b*, plate 4.) Near the top of each a pulley was inserted twenty-seven inches from the mattress. Cords were tied to the loops of the encircling knee-bands and run through these pulleys, to each of which was suspended a three-pound weight. This was traction on a line opposed to the dislocation, namely, forward and downward, and was alternated with the parallel extension, one or the other being used day and night. At no time were the limbs allowed, even for a moment, without extension. When the bands were changed from knees to ankles, and *vice versa*, an assistant always held the limbs firmly extended until the weights were reapplied. This manual traction was likewise maintained when the patient was temporarily moved to another bed. The three-inch wide pelvic belt, previously described, was constantly worn, and beneath it, on each side, narrow, firm, graduated compresses, made of soft linen, were placed across the dorsum illi, immediately above the trochanter major. One of the compresses was nine inches long, four and a half wide, and a quarter of an inch thick; the other, five and a half inches long, five inches wide, and a quarter of an inch thick.

By means of this belt and compresses pressure was maintained with the object of obliterating the sacs in which the bones had played.

The limbs were now rarely displaced. All the unavoidable movements attendant on change of apparatus, etc., were borne without injury, indicating that the heads of the thigh-bones were to a very considerable extent secured in their new position.

Thus much having been gained, the next object was by passive flexion and extension, in imitation

⁹ I will here mention that Drs. J. C. Warren and H. H. A. Beach saw this patient at intervals during the course of the treatment, and later Dr. D. W. Cheever also examined her from time to time.

¹⁰ The Hip, p. 56.

of the natural motions of the joints, to still further promote absorption, and perhaps cause some degree of irritation in the immediate neighborhood, and consequent effusion of adhesive matter.

To accomplish this three sets of pulleys were employed, namely, the parallel at the foot of the bed (*a, a*, plate 3, *a, a*, plate 4), the high ones at the foot of the bed (*b, b*, plate 3, *b, b*, plate 4), and two additional pulleys at the head of the bed (*c, c*, plate 3, *c, c*, plate 4). The high pulleys (*b, b, b, b*) were used, as before, for oblique traction from the knee-bands. The cords from the high pulleys (*c, c, c, c*) were likewise attached to the knee-bands in front. To this a light weight was suspended, which served as a handle, and when drawn upon raised the thigh to an angle with the body. The cords over the parallel pulleys (*a, a, a, a*) were tied posteriorly to a soft roll of padded linen, which passed round the upper extremity of the thigh. When the joint was flexed to an obtuse, or right, angle by drawing down the weight on pulley *c, c, c, c*, it is evident that the weight on *a, a, a, a* acted upon the femur as upon the extremity of a lever, preventing the head from rising when the upper pulleys (*c, c, c, c*) were in action. It will be seen that by this arrangement we have flexion and extension of the articulation, with every safeguard against displacement. An attendant stood, or was seated, at the head of the bed and drew the small weight downward, by which action the knee was raised toward the abdomen. The weights at the foot then pulled it back again. At the same time the pelvic leather belt buckled over the summits of the trochanters caused strong pressure against the pelvis. Four hours a day this passive exercise was continued, two hours for each limb. This may properly be termed the stage of excavation. The blocks from beneath the legs at the head of the bed were placed under the foot. The trunk then served as counter-extension. The long straps were removed from the head-rail. As an additional security a belt eight inches wide, made of soft, pliable leather lined with thin leather, and cut going to fit the hips, was used outside of the narrow belt. This wide belt had five small buckles and straps or billets of webbing, each one and a half inch wide, and round, padded perineal straps. The wide belt encroached somewhat on the upper part of the thighs, and a slit was required for the passage of the perineal straps to buckles in front. During the passive exercise the outer belt was removed. The course I have here indicated was pursued, with slight variations, until July 20, 1883. The time allotted for exercise had been gradually lessened. The high pulleys at the foot were used one day and night. The next day and night the patient wore the wide belt, thus alternating from one to the other.

The hips remained in position. The last time either had been displaced was on March 28, 1883.

June 2, 1883. The patient was removed to the seaside to a hotel in the neighborhood of my summer residence. Every precaution was observed in the removal. She wore the narrow belt, with the compresses above the trochanters, and the wide belt over it. She was placed in a hospital basket. The long straps from the narrow belt were secured to the head end of the basket. The perineal straps

were buckled in place. The straps from the knee-bands were fastened to the lower end of the basket.

In this way the patient was made a solid fixture to the basket, and could be moved in any way without disturbing her position. She arrived safely at her destination. This same arrangement was used through the summer. By it she was enabled to be upon the piazza or beneath the shade of the trees almost daily.

As an indication of the firmness with which the bones were held in their new position, I will mention that on one occasion much alarm was caused by the child rolling out of bed in the night. When reached, her head and hands were upon the floor, and her body was suspended obliquely by the foot-weights, which brought up against the bed-rail. No injury arose from this accident.

July 20, 1883. The first attempt was made to test the strength of the tissues by which the heads of the femurs were retained in their sockets. I pushed the limbs upward, and then upward and backward, with considerable force. There was firm resistance in each direction. From this date this experiment was frequently renewed, more and more forcibly, until finally the body was moved by pushing against the rim of the acetabulum, which was now unquestionably formed.

Dr. Beach saw the patient again in October, 1883, after her return to Boston. He judged that the upward pressure thus employed would amount, calculated by pounds, to more than the weight of the child. The patient now used active movements of the limbs in imitation of the previous passive exercise with weights. In order to avoid the friction of the foot upon the bed the heel rested in the hand of an attendant in its to-and-fro movement. This was done with a weight of two pounds attached to the ankle, for five minutes, twice a day.

Dr. D. W. Cheever, at my request, examined the patient at this time and freely manipulated the limbs.

The fourteenth day of January, 1884, the little girl sat up in bed, for the first time for thirteen months.

The period had now arrived for an initiation into the so long forbidden and forgotten faculty of locomotion.

A cart was constructed, of which plate 5 is a representation. It is very simple in its make, and perfectly answered its purpose. The plate requires but little explanation.

The heavy leather strap with the saddle in the centre is the important feature. The patient sat astride upon the saddle, which when first used was buckled upon the cross-bars, at such a height as only to allow the extremity of the toes to touch the floor. The crutches, as is plainly seen, were made to slide up and down and fixed by thumb-screws. They were of elastic wood and were sprung into and fitted the axillæ, where they were secured by a webbing belt. They relieved the perinæum of a portion of the weight. From time to time the saddle and the crutches were let down until she walked squarely upon her feet.

At first she only imitated walking, but soon was able to propel herself through two large rooms and a hall with ease. The hip-joints worked perfectly.

As a safeguard against accidents the narrow leather belt was still worn. In April the patient bore her own weight for a few moments at a time, and on the 11th took her first drive sitting up. May 14th she began to walk in her carriage, without the saddle, and on the 16th without either saddle or crutches, taking hold of the side-bars. On the 18th she walked, supported on each side by a person holding a towel which was passed through a loop in a belt around her waist, steadying herself by grasping a cane held above and in front. I will here quote a few minutes from a journal kept by the patient's mother, of which I have availed myself for many of the data in the previous pages.

It is right to add that to her faithful, unwearied, and determined perseverance the success which has attended the case may to a great extent be attributed.

Note for June 9, 1884. "While sitting in her chair, got off the chair, turned round, and stood up; turned round, then turning back again, sat down.

"June 11th. Walked by herself without support a distance of three feet. Did this six times.

"June 12th. Walked twelve times without support, the distance being a little greater.

"June 19th. Having more confidence in herself she walked about the room alone.

"June 24th. Light corsets with steel ribs instead of whalebone were put on to support the spine, which seemed weak.

"June 28th. Commenced to use canes, which gave her confidence in walking."

On the thirtieth of June, 1884, the patient left Boston for her home in the southern part of the State.

It is worthy of remark that the patient, with slight exceptions, remained in excellent health throughout the long confinement, was always in good spirits, retaining her flesh and a healthy color in her cheeks. On account of her appreciation of the importance to herself of the course pursued, she much facilitated the treatment.

Plate 6 is from a photograph taken just before leaving the city. It is a back view. The child is reclining on her left side to show the hip and the symmetrical outline of the body at this period.

My later notes of the case are as follows:—

January 20, 1885. The child walks well without canes. There is some lateral movement of the shoulders due to spinal debility which is probably the result of long continuance in the horizontal position. The left knee obliques slightly inward, causing the limb to appear somewhat shorter than the right and producing some inequality in her gait.

February 18, 1885. The walk has much improved, partly in consequence of shoes being worn, which support the weak ankles. There is some degree of muscular atrophy between the trochanter and the crest of the ilium, on each side, where formerly rested the femoral-heads. This depression is also due in part to the pressure of the belt and compresses. The limbs can be pushed upward, and upward and backward, against the lips of the newly formed acetabula with a force which rolls the pelvis, and moves the body without the slightest displacement.

March 23, 1885. The lateral movement of the

shoulders disappeared some weeks since. As strength has increased the left knee has become nearly straight. The tread is firmer and more even.

Nélaton's line, passing over the summits of the trochanters, indicates that the heads of the femurs are in their normal position. They have not been displaced since March 28, 1883.

May 25, 1885. The child's walk is normal, and she enjoys walking; she steps with natural firmness and vigor. Like other children she takes especial pleasure in running.

The photographs from which the two pictures composing plates 7 and 8 were reproduced were taken March 23, 1885, about two years and three months from the commencement of treatment. The first figure, plate 7, is standing in a position to correspond with the figure in plate 1, which latter gives a side view previous to treatment. Plate 8 is a front view to correspond with the second figure of the series seen in plate 2.¹¹

THE SELECTION OF A SUITABLE CLIMATE FOR THE VARIOUS FORMS OF PULMONARY CONSUMPTION.¹

BY J. HILGARD TYNDALE, M.D., OF NEW YORK.

THE law of the beneficial influence of climate in pulmonary consumption may be remembered by this adaptation of a biblical phrase: *There abideth equability, dryness, and elevation, these three; but the greatest of these is dryness.*

The following tables will serve as a guide to the various degrees and qualities of climatic constituents. The humidity table is R. v. Vuierot's; the rest are my own.

TABLES OF THE CONSTITUENTS OF CLIMATE:

Humidity, equability, elevation, and degrees of warmth.

(1) Humidity.

Excessively dry: 1 to 55 % of relative humidity.

Moderately dry: 56 to 70 % of relative humidity.

Moderately moist: 71 to 85 % of relative humidity.

Excessively moist: 86 to 100 % of relative humidity.

(2) Equability.

Very equable: 15° to 35° of monthly range.

Moderately equable: 36° to 50° of monthly range.

Unequable: 51° to 65° of monthly range.

Very unequable: 66° to 85° of monthly range.

(3) Elevation.

Low altitude: 500 to 1,000 feet above sea-level.

Moderate altitude: 1,100 to 2,000 feet above sea-level.

Medium altitude: 2,100 to 4,000 feet above sea-level.

High altitude: 4,100 to 7,000 feet above sea-level.

(4) Temperature.

Hot: 80° to 100° of monthly mean.

Warm: 55° to 79° of monthly mean.

¹¹ The illustrations which accompany this paper are exact reproductions of the original photographs by a new application of the gelatine process, which preserves great delicacy of detail and outline.

¹ Concluded from p. 517.