in addition to their occurrence as nasal polypi originating from the mucous and submucous tissues of the nose, they may originate from these structures elsewhere or in the intermuscular septa, nerves, subserous fat and periosteum.

Symptoms.—The chief symptom is nasal stenosis which increases with the development in size and number of the polypi, until complete obstruction of one or both nostrils results. Mucous or muco-purulent discharge, cephalalgia, aural complications and other symptoms of a catarrhal nature, together with those incident to mouth-breathing, are observed. To quote the words of a sufferer: "It affects the sight, the hearing, the taste, and the smell of course" Spasmodic asthma, paroxysmal cough and sneezing attacks are among the reflex phenomena which are occasionally excited. For diagnostic purposes it is usually only necessary to look with a good light and to feel with a probe in order to establish correspondence with the physical characters just described, but more rarely an accurate knowledge of all pathological states is essential to a precise diagnosis.

The treatment has been outlined in the relation of the cases. It consists first in the establishment of free nasal passage for respiration, drainage, vision and instrumental manipulation, and to this end in the reduction of hypertrophied turbinated bodies, the removal of septal excrescences, hypertrophied tonsils, etc.

To reduce the hypertrophied turbinated bodies the electro-cautery, applied by means of the point, knife, or snare electrode, is the most convenient agent, although chromic acid, carefully used, will suffice. One of the best methods is to employ a rather heavy point-electrode having its end slightly "curved on the flat" toward the side to be cauterized, and to draw this, in one application, three or four times, at a little more than cherry-red heat, through approximately one inch of the length of the turbinated body, passing each time through the same line until the soft tissue is penetrated to the bony base. From three to six treatments, in different lines on each of the lower turbinated bodies, may be required. Fewer suffice for the middle bodies.

Septal excrescences are removed by the nasal saw, chisel, or burr and trephine drills propelled by the surgical engine or electrical motor. The latter apparatus is more useful for drilling away projecting edges and corners after the use of the saw than for execution of the entire operation. A large variety of nasal saws is necessary in order best to conform to individual cases.

If the septum is seriously deflected it may be necessary to correct it.

Adenoid vegetations of the naso-pharynx must be cleared away.

While this work is progressing such polypi as can be reached should be removed, and others as rapidly as access is gained. This is done by the

galvano-cautery snare, the cold snare, or fine polypus forceps, whichever seems best adapted to the individual polyp.

All operating is done under the local anæsthesia, and *retraction of tissues* produced by cocaine applied by spray in solution not to exceed 5 per cent., and on cotton not to exceed 2 per cent., and operating must cease as soon as hæmorrhage obscures the view.

But the real success of the treatment, after having gained access to the polypi, consists in tracing them to their points of attachment and in thoroughly cauterizing these so-called roots; if not at the same sitting, then at the next, remembering meanwhile the exact spot. Knowing the *hiatus semilunaris* to be a favorite point of origin, those polypi which proceed from beneath the middle turbinated body should be followed up by insinuating a fine electrode slightly curved on the flat to this point, and those which spring from the superior meatus posteriorly must be reached by a properly curved point-electrode introduced through the mouth and naso-pharynx, always under the best illumination.

The latter procedure is by far the most delicate part of the operation, and it is often necessary to tie the velum forward or to use a self-retaining palate retractor.

No. 70 Monroe St., Chicago.

THE IMPORTANCE OF LOCAL TREATMENT IN SYPHILIS.¹

Read before the Section on Dermatology and Syphilography, at the Thirty-ninth Annual Meeting of the American Medical Association, May, 1888.

BY JOSEPH ZEISLER, M.D.,
OF CHICAGO.

PROFESSOR OF GENITO-URINARY AND SKIN DISEASES, CHICAGO POST-GRADUATE MEDICAL SCHOOL; MEMBER OF THE AMERICAN DERMATOLOGICAL ASSOCIATION.

There are few diseases like syphilis, for which our therapeutical actions are so well defined, for which we possess specific remedies, upon whose never-failing effect we count with nearly absolute certainty; yet nowhere is the Latin phrase equally true: *duo si faciunt idem, non semper est idem*. The where, when, and how to use those two chief agents, mercury and iodine, seems to be to many a *terra incognita*, and the prescribing of the mixed treatment forms the beginning and the end of their fight against that dread disease, in whatever phase it may be, whether it has attacked the skin only or other more important organs. The result of this *laissez aller* is often a very sad one, and many a perforated septum or palate, or even the loss of a considerable part of the nose is due to the fact that local treatment has not been given a proper place in the management of that case. This may sound somewhat hard, but I speak from ex-

¹ This paper was mislaid by the Section officers into whose hands it came. Hence its late appearance.

perience, and having observed a few such cases of late which impressed me very strongly, I have chosen this theme for my present paper. There is nothing especially new to be said in regard to this subject, at least not for those engaged in the special line of dermatology and syphilis, but it seems to me to be not at all superfluous to call the attention of the practitioners at large to the important rôle which local treatment plays in the management of syphilis, as the text-books usually touch this question in a rather superficial manner. In laying special stress upon local applications, I hope I shall not be understood as underestimating the necessity of constitutional treatment in every single case; this, of course, is a *conditio sine qua non*.

There is in nearly every stage of syphilis more or less opportunity for the successful employment of local remedies, while in some cases, mostly such showing tertiary forms, they are of the greatest importance.

In regard to the primary indurated sore, the majority of physicians to-day refrain from giving constitutional treatment and employ various local measures to secure a speedy cicatrization. The question of excision might well be discussed here, should it not abstract us too much from our theme; there can little be argued against it, when the localization allows it, as healing often takes place by primary intention. As regards its preventive effect opinions are divided. Believing the induration to be the sign of general infection, I fail to see how excision can abort the disease, and the reports of seemingly successful operations are surely capable of different explanations. In the management of hard chancre, it is always desirable to make the induration disappear as quickly as possible. For this end I have found the mercurial plaster a very valuable agent; it constitutes the simplest, cleanest, most practicable and efficacious treatment of the primary sore. As reference will again be made to this plaster, I wish to say that I always use that prepared and spread by Seabury and Johnson, which in every respect proved satisfactory.

After secondary symptoms have appeared, there is a very useful field for local applications. In selecting the plan of a general treatment I have always given preference to the time-honored frictions, which, in spite of all objections, still form the most reliable way to treat the earlier manifestations of syphilis. Besides their constitutional effect they have an undoubted local influence upon the cutaneous eruptions, causing them to vanish usually very rapidly. Patients in private practice will mostly desire before all to have the visible manifestations, as on forehead and hands, disappear as soon as possible; for this purpose a 10 to 20 per cent. ointment of the white precipitate will often work like a charm, and this result will always be highly appreciated by pa-

tients in the better walks of life. Still, the papular syphilides on the palms and soles, the so-called psoriasis specifica, will occasionally prove very obstinate and will call for more energetic measures. Thus chrysarobin has been recommended by Reumont² and by Mosengeil,³ who have successfully used it as an ointment of 10 to 20 per cent. strength. Sigmund⁴ speaks very highly of a 1 to 2 per cent. solution of sublimate in collodion. This should be painted once or twice daily on the infiltrated plaques; fissured places, of course, should not be touched by it, these are best covered with strips of the gray plaster. Gilles de la Tourette⁵ has found warm local baths with an addition of 1 part each of sublimate and of chlorate of ammonia to 2,000 parts of water, as very efficacious for that trouble. Papules appearing on mucous membranes, mostly in the throat, will readily heal when touched daily with nitrate of silver in strong solution or in substance, and under the use of an astringent gargle.

Of far more importance are topical applications for moist papules or condylomata lata, as these not only are the cause of much annoyance and discomfort to the patient, but also form one of the most common sources of syphilitic infection. As they are so easily reproduced by coaptation of opposing surfaces of the skin, as, for instance, in the genito-crural or anal folds, it is before all necessary to keep such places carefully separated by the application of absorbent cotton or gauze, and to scrupulously remove the fetid secretion by frequent cleansing with diluted sublimate solutions. For the removal of those characteristic vegetations developing so often on such places different caustic pastes have been recommended, but their application is attended with a great deal of pain. I usually prefer the cauterization with the silver stick, after previously anæsthetizing the parts by a strong cocaine solution. Moist papules on the corners of the mouth or between the toes are often very painful, but they heal readily when touched with nitrate of silver and covered subsequently with mercurial plaster.

There is rarely occasion to act against the swelling of lymphatic glands. I lately treated a young man in whom a group of infiltrated glands, situated between the lower jaw and the mastoid process, caused a constant pain and great difficulty in swallowing. An ointment containing iodine, iodide of potassium and ichthyol reduced the swelling in two days and brought perfect relief. For suppurating bubos surgical interference will, of course, be necessary.

During any form of treatment with mercurial preparations the mouth will always require attention, as the saliva soon becomes impregnated with the drug and is the direct cause of mercurial stom-

² Chir. Centralbl., No. 3, 1880.

³ Berl. Klin. Wochenschrift, No. 22, 1879.

⁴ Wr. Med. Wochenschrift, No. 41, 1879.

⁵ Progrès Médical, No. 30, 1886.

atitis. The most diligent care of the teeth and gums, abstinence from tobacco and irritant substances, and the frequent use of an astringent mouth-wash will greatly aid to delay this incident. Brockhart⁶ recommends a solution of 5 to 10 parts of tannic acid in 20 parts of glycerine to 80 parts of water for this purpose. The gums may sometimes become so tender as to make mastication very painful and to necessitate an interruption of the administration of mercury. To overcome this the same author has suggested the use of a 10 per cent. cocaine solution, to be painted on the delicate parts a few minutes before meals.⁷

We have thus far seen that, during the earlier manifestations of syphilis, there are many indications for local treatment. In the later stages of the disease it becomes often imperative, and Hutchinson, who regards tertiary symptoms as mere local sequelæ, has gone so far as saying that only local remedies will lead to success in treating them, whereas constitutional treatment is of minor value.⁸ There is much truth in this. While general treatment, especially the use of iodides, is certainly capable of removing many of the symptoms incidental to the so-called tertiary period, it will be entirely unable to arrest the destruction which, for instance, serpiginous ulcers produce on the skin and mucous membranes, and here it is where local treatment is followed by the most brilliant results. Allow me to illustrate this.

Anna S., 16 years old, was brought to my office the 17th of December, 1887, through the kindness of Dr. C. Fenger, who in an accompanying letter asked me to conduct the medical treatment, while he later on should do the surgical part of the work. Without going into the details of the history of this case, which, from its aspect (general cachexia, Hutchinsonian formation of teeth, etc.), and from the anamnesis, had to be regarded as due to hereditary causes, I shall briefly state that the whole upper lip and both alæ nasi were the seat of those characteristic ulcerations resulting from the breaking down of gummatous neoplasms. The cartilaginous part of the septum was perfectly destroyed, only the small anterior cutaneous bridge remaining; accordingly the dorsum was depressed as a result of the defect. The anterior portion of the nose, where not ulcerated, presented that peculiar livid tint, and it seemed only too natural that the necessity of a rhinoplastic operation in the near future was suggested to Dr. Fenger. The affection was said to have begun with a peculiar eruption on the nose about one and one-half year ago, which was followed by a swelling of the upper lip. For the last six months she had been under the care of a prominent specialist for the diseases of the throat and nose. *Sapienti sat!*

I immediately began energetic local treatment

by using the silver stick very liberally every second day. The patient was directed to clean the sores twice daily with a 1 per cent. sublimate solution, and to cover the same in the meantime constantly with the adhesive gray plaster. Iodide of potassium in daily doses of 2.00 was prescribed for internal use. The effect of this treatment was all I could hope for; within ten days the ulcerations were completely healed, a very smooth cicatrization having taken place, and the shape of the nose was preserved. By inserting two small rubber tubes the contraction of the scar tissues around the opening of the nose was prevented. The patient, of course, was only too glad to be saved the trouble of a plastic operation.

Cases like this are, fortunately, in our days not very common; but they must not happen at all, and to let the destruction go on undisturbed until an important organ is endangered, borders on criminal negligence or ignorance.

I have on record the notes of a case where extended, deep syphilitic ulcerations had existed and spread on the scalp of a man for over two years when I began to treat him, in December, 1886, after the principles just mentioned, and in less than two weeks perfect cicatrization had taken place, the patient remaining well ever since.

To the kindness of Dr. M. Mannheimer, of Chicago, I owe a very interesting case of a woman about 30 years old, whom I treated in the winter of 1884. She had been infected about 1½ yrs. ago by her husband, who confessed to having had an indurated chancre, but otherwise did not remember to have suffered from any symptoms of constitutional lues. The woman herself knew nothing of having had a primary sore, nor the usual secondary manifestations, but the first thing she noticed was the appearance, judging from her description, of a generalized rupia about a year ago, which soon became ulcerated. When I first saw her, she was simply covered with those horse-shoe formed sores; the whole scalp was one ulcerated surface, producing an unbearable smell. The back of the right hand was nearly perfectly destroyed, and about sixty more ulcers, from the size of a penny to that of the palm of a hand, were present on the rest of the body. The forehead, nose and chin were the seat of an ecthyma eruption. The patient suffered a great deal, and was hardly able to find comfort in any position. I ordered iodide of potassium 2.00 pro die in solution, a warm bath medicated by 10.00 of sublimate daily for an hour, emplastrum hydrargyri as a local dressing, and white precipitate salve for the face, while the scalp, for the first few days, was to be saturated with salicylated olive oil to facilitate the removal of crusts and detritus, and afterwards washed daily with sublimate soap and anointed with the mercurial salve. The sores were touched every third day with nitrate of silver. Healing soon began to take place on some of the smaller ulcers,

⁶ Monatshefte f. prakt. Dermatologie, No. 8, 1885.

⁷ Ibid., No. 2, 1886.

⁸ Syphilis, Discussion, London Pathological Soc., 1876.

and after a little over two weeks the patient was completely cured, and only continued to take the iodide for some time. She has remained well ever since.

I could cite quite a number of other observations just as striking, but these few examples may be sufficient to show the efficiency of local treatment in these graver forms of cutaneous syphilis.

I wish to add only one word with reference to the cauterization by nitrate of silver, which I regard as a very essential part of the treatment. Its use was formerly accompanied by a great deal of pain; now I always employ previously a 5 to 10 per cent. solution of cocaine for a few minutes, by which it is possible to make that operation without the least objection on the part of the patient.

Ulcerating syphilides may occasionally become phagedenic and resist all kinds of treatment. For such cases Spillmann⁹ has recommended the free use of Volkmann's sharp spoon, subsequent application of Paquelin's cautery and dressing with sublimate solution. I have so far had no occasion to resort to these radical measures.

It would be beyond the scope of this merely suggestive paper to speak at length about the treatment of syphilitic affections of the tongue, the nose, throat and larynx, the eye and ear, the bones, etc.; this must be left to those who make these organs their special study. What has been said in regard to the skin applies still more to them. When necrotic bones are the cause of a chronic ozæna, when a gumma is about to destroy an eyelid, when exfoliative perichondritis in the larynx may at any moment endanger the life of a patient, there is no time to wait for the effect of internal treatment; this will certainly in many respects be a valuable adjuvant, but only *local* applications will be able to arrest the destructive processes in those important organs.

MEDICAL PROGRESS.

FORTY-TWO CASES OF INTUBATION.—DR. F. K. PRIEST reports: In a series of 266 cases admitted to the Willard Parker Hospital, intubation was performed for the relief of laryngeal stenosis, in 42, or a little over 22 per cent. of that number, with a percentage of recoveries of 26.19. The majority of these cases were of undoubted diphtheria, with well-marked symptoms of laryngeal stenosis, at the time of admission, while many of them had been without proper care or medical treatment and well-advanced in the last stages of septicæmia, the operation being performed merely for the relief of the stenosis, with no hope of ultimate recovery.

Of the 42 cases 8 were cases of diphtheria com-

plicating or accompanying other contagious diseases, as follows: 1 case of measles and diphtheritic conjunctivitis; 1 of varicella; and 6 of scarlet fever. In 35 of the 42 cases membrane was present in the throat or anterior nares, or coughed out either before or after the introduction of the tube; in the 7 remaining cases, the presence of membrane was not detected at any time during the course of the disease, although there was more or less swelling of the cervical glands with discharge from the nostrils. Only one of the seven recovered.

On admission in 29 cases, the symptoms of laryngeal stenosis were well advanced, 25 of this number being unquestionably diphtheria, and one case in which the presence of membrane could not be detected, a large piece was coughed out during the introduction of the tube. In 5 cases there was a hoarse, croupy cough without dyspnoea, and in 8 the laryngeal symptoms developed after admission. In all cases where the character of the dyspnoea and the general condition of the patient permitted of the delay, intubation was not resorted to until other means for the relief of the stenosis had been tried and failed to give more than temporary relief.

In 3 cases only, where the urine could be obtained, albumin was not found on examination, several dying from acute nephritis with suppression of urine, as occurred once five days after the removal of the tube, all local evidences of diphtheria having disappeared at the time. This one has been included in the list of fatal cases, as the patient was not discharged from the hospital, although it might be considered as a successful case of intubation.

In those cases that recovered, the average time the tube remained in the larynx was 6½ days. No definite rule, however, was followed as to the time of removal. In cases that were progressing favorably the tube was left in position until the membrane had disappeared from the throat and nose and the fever had subsided. The tube was removed in one case on the fifth day, the membrane being still present on the tonsils; on account of violent and persistent coughing and vomiting after taking fluids, the urine was also scanty, high colored, nearly three-fourths albumin; after which the cough was less troublesome, vomiting ceased, and the child took plenty of fluids; although the breathing was somewhat stridulous there was no return of the dyspnoea.

Coughing after drinking was a constant symptom and, in several cases, it was severe enough to induce vomiting. The above case was the only one, however, in which this symptom was severe or constant enough to seriously interfere with the ordinary feeding or treatment. The most troublesome complication, and one that in several cases was productive of very alarming symptoms, was from the collection of mucus in

⁹ Ann. de Derm. and Syphilogr., 1885, p. 712.