Annals and Magazine of Natural History: Series 6

Publication details, including instructions for authors and subscription information:
<http://www.tandfonline.com/loi/tnah12>

IX.—On newly-discovered East-African Chameleons, with remarks on some other Reptiles described by Dr. Steindachner

G. A. Boulenger

Published online: 02 Oct 2009.

To cite this article: G. A. Boulenger (1892) IX.—On newly-discovered East-African Chameleons, with remarks on some other Reptiles described by Dr. Steindachner , Annals and Magazine of Natural History: Series 6, 9:49, 72-74, DOI: [10.1080/00222939208677273](https://doi.org/10.1080/00222939208677273)

To link to this article: <http://dx.doi.org/10.1080/00222939208677273>

PLEASE SCROLL DOWN FOR ARTICLE

Taylor & Francis makes every effort to ensure the accuracy of all the information (the "Content") contained in the publications on our platform. However, Taylor & Francis, our agents, and our licensors make no representations or warranties whatsoever as to the accuracy, completeness, or suitability for any purpose of the Content. Any opinions and views expressed in this publication are the opinions and views of the authors, and are not the views of or endorsed by Taylor & Francis. The accuracy of the Content

should not be relied upon and should be independently verified with primary sources of information. Taylor and Francis shall not be liable for any losses, actions, claims, proceedings, demands, costs, expenses, damages, and other liabilities whatsoever or howsoever caused arising directly or indirectly in connection with, in relation to or arising out of the use of the Content.

This article may be used for research, teaching, and private study purposes. Any substantial or systematic reproduction, redistribution, reselling, loan, sub-licensing, systematic supply, or distribution in any form to anyone is expressly forbidden. Terms & Conditions of access and use can be found at <http://www.tandfonline.com/page/terms-and-conditions>

So far as I have yet been able to determine, the development of the crocodile closely resembles that of the bird. A noticeable feature is the tail, which is of great length at a very early stage, and is at first rolled up in a spiral form, and afterwards, when the embryo is more strongly bent, twisted round the neck.

That the rudiment of the egg-tooth appears very early has already been mentioned.

Rudiments of the genital protuberance are already to be seen in embryos which are about 10 millim. long (measured in the bent position). A rod-shaped structure may then be observed between the posterior legs; it is about 1 millim. in length and protrudes from the cloaca, with the anterior wall of which it is fused. It at first lies parallel to the median line of the abdomen, is subsequently erected, and finally completely retracted within the opening of the cloaca. It is not until the embryos are almost full-grown, after about two and a half months, that the genital protuberance begins to disappear altogether, and is then only to be seen by separating the lips of the cloaca.

IX.—*On newly-discovered East-African Chameleons, with Remarks on some other Reptiles described by Dr. Steindachner.* By G. A. BOULENGER.

JUDGING from the number of descriptions recently published* it would seem that the Chameleon fauna of East Africa is likely soon to rank next to that of Madagascar with regard to variety of species. I have, however, no doubt that the list of species has to be reduced by three, of which two have just been described by Dr. Steindachner and the third by Mr. Stejneger. On comparing the descriptions and figures of *Chamaeleon Höhnelii* and *leikipiensis* with the late J. G. Fischer's account of *Ch. bitanaiatus* and the specimens in the British Museum, I cannot come to any other conclusion but that they all belong to one and the same species, Dr. Steindachner's specimens being fully-developed males. The

* *Chamaeleon sphaerophilis* (Victoria Nyanza) and *Fischeri* (Usagara ?), Reichenow, Zool. Anz. 1887, pp. 370 and 371; *Ch. Roperi* (Kilifi), Boulenger, P. Z. S. 1890, p. 85, pl. viii. fig. 4; *Ch. Höhnelii* (Leikipia, west of Kenia, 6000 feet), *leikipiensis* (Leikipia), and *tavetensis* (Taveta, foot of Kilimandjaro), Steindachner, Anz. Ak. Wien, 1891, pp. 141 and 142, and Sitzb. C. 1891, pp. 307, 309, 310, pl. i.; *Ch. Abbotti* (Kilimandjaro), Stejneger, Bull. U. S. Nat. Mus. xiv. 1891, p. 353; *Rhampholeon Robecchii* (Somaliland), Boulenger, Ann. Mus. Genova (2) xii. 1891, p. 11, pl. i. fig. 3.

Downloaded by [North Carolina State University] at 20:24 09 March 2015

Downloaded by [North Carolina State University] at 20:24 09 March 2015

Downloaded by [North Carolina State University] at 20:24 09 March 2015

Downloaded by [North Carolina State University] at 20:24 09 March 2015

Downloaded by [North Carolina State University] at 20:24 09 March 2015

Downloaded by [North Carolina State University] at 20:24 09 March 2015

Downloaded by [North Carolina State University] at 20:24 09 March 2015

- Downloaded by [North Carolina State University] at 20:24 09 March 2015

purpureus, Cant., to be the same as *L. effrenis*; the latter has three labials in contact with the eye, the former only two. The type of *Lycodon ophiteoides*, Blkr., is preserved in the British Museum.

2. *Simotes Meyerinkii*, Steind., is, I suggest, only a variety of *S. octolineatus*, Schn.
3. *Chalcides Simonyi*, Steind., from Fuertaventura, I regard as a variety of *C. viridanus*, and as there is a *C. viridanus*, var. *Simonyi*, either of the names will have to be changed. My reason for not accepting *C. Simonyi* as a valid species, although some of its characters do not fit into the diagnosis I have given of *C. viridanus* (Cat. Liz. iii. p. 402) is that a female specimen from Grand Canary, preserved in the British Museum, falling, as regards coloration, into Steindachner's var. *bistriata*, β , and with 36 scales round the middle of the body, agrees precisely in its proportions with *C. Simonyi*, as may be seen from the following measurements:—

	millim.
From snout to vent	87
From snout to fore limb	25
Head (to ear-opening)	13
Width of head	10
Fore limb	15
Hind limb	22
Tail (reproduced)	78

4. *Molge Luschani*, Steind.—I have no doubt this is a *Salamandra*. There seems to be less difference between *Salamandra Luschani* and *S. caucasica* than between the latter and *S. maculosa*.

X.—Description of a new Snake from Nubia.

By G. A. BOULENGER.

Gongylophis Muelleri.

Rostral large and broad, with angular horizontal edge; upper surface of snout and crown with small smooth shields, the largest of which is an azygous shield behind and wedged in between the internasals, which form a short suture behind the rostral; five shields from eye to eye across the forehead; nine or ten scales round the eye, which is separated from the labials by a single series of scales; nine upper labials. Scales