The Journal of the American Medical Association

Published under the Auspices of the Board of Trustees.

VOLUME LI

CHICAGO, ILLINOIS, NOVEMBER 7, 1908

NUMBER 19

Original Articles

TECHNIC OF ABDOMINAL HYSTERECTOMY.

ARTHUR E. HERTZLER, M.D.
KANSAS CITY, MO.

New methods for the removal of the uterus in whole or in part by the abdominal route have become so numerous that it would seem impossible to present anything that is new. The recent modifications in technic have been, however, for the most part, designed to shorten the time of operation at the expense of exactness, in the hope of thereby lessening shock. Surgeons still seem to believe that the duration of an operation can be measured in minutes and hours. The condition of the patient at the conclusion of an operation is the only index as to whether the operation has been a short or a long one. Ten minutes is a long time for an operation if the patient is shocked; two hours is a short time if the patient wakes up promptly with everything in perfect condition.

It is a great fallacy to believe that time is the only or even the chief element in the production of shock. It is the degree of injury to nervous tissue and loss of blood which are the primary elements, and a rough operator will shock a patient more in a short time than a gentle one will in a much longer time.

Those who advocate a technic that has for its chief end the abbreviation of the time of the operation and who purchase this speed by mass ligatures and crushing forceps seem to overlook the fact that a crushed stump of a ligament wound up in a mass of peritoneum and all held together by a tightly drawn ligature may continue its shocking influence long after the surgeon has gone home. For him the time of operation has been shortened it is true, but for the patient it has been indefinitely prolonged.

With these preliminary points in mind I do not hesitate to present a technic that has grown in my hands in the course of years which aims at anatomic exactness rather than at speed. I have aimed to treat the pelvic stumps with the same consideration that is given the stump of an amputated limb. The points to be observed in both instances are the same: 1. Remove what is diseased. 2. Tie that which bleeds and nothing more. 3. Arrange the remaining parts so as to preserve the best possible function. 4. Cover the whole with the natural covering of the parts.

TECHNIC OF THE OPERATION.

The technic is as follows: The incision is made in the median line and the uterus or tumor delivered, if possible. The broad ligament is then grasped on either the mesial or lateral side of the ovary, depending on whether this organ is to be removed or not. The fingers

of the left hand are placed behind the ligament (Fig. 1) and the blade of the peritoneal scissors¹ is passed below the anterior layer of the peritoneum and above the vessels. The peritoneum is cut to the median line or entirely across to the opposite side as desired. The peritoneum is pushed gently back with gauze or with the tip of the scissors. The ovarian vessels are then located and ligated by passing eye first a needle threaded with No. 1 catgut. The artery is isolated and tied separately, or the ovarian veins may be included in the same ligature with the artery, since they lie close together and are both small (Fig. 2). In case of varicosities of the broad ligament, however, the veins should have a separate ligature. The uterine veins should next be tied. They lie above the uterine arteries and may conveniently be tied first. The point of ligation of the uterine artery varies with the conditions to be met. If the cervix is to remain I prefer to tie it close to the uterus, but if the cervix is to be removed, certainly in operating for cancer, I prefer to tie it lateral to the point of bifurcation. It is necessary in this instance to expose the ureter and tie lateral to it. The number of veins requiring separate ligation varies, usually not more than two separate ligations need be made on either side, but in large vascular fibroids three or more may be applied with advantage. Many of the cuts in standard textbooks show the arteries alone ligated. The veins, too, require ligation quite as much as the arteries.

After the ligatures have all been placed the broad ligament is cut down to, but not through, the peritoneum. The fibrous tissue with the vessels now retract and the posterior peritoneal layer may be cut through, some distance from the plane of severance of the fibers, forming in this way a peritoneal flap which is to be used later to cover the raw surfaces (Figs. 3 and 6). If the cervix is allowed to remain it is cut wedge shaped (Fig. 3). The cut surfaces so made are then united, after treating the canal as one sees fit, with a figure-of-eight suture drawn tight enough to hold the surface in apposition (Fig. 4). These sutures it should be noted do not penetrate the peritoneum. The depth of the sutures may depend on the disposition of the stump to ooze. When the suture is tied the cervical canal is effectually closed off above (Fig. 5). The peritoneum is now united over the raw edges of the broad ligament and the stump of the cervix (Fig. 5). A Lembert suture or a mattress, using No. 0 plain catgut is best. This suture penetrates the peritoneum and nothing else and is drawn just tight enough to bring the surfaces gently together.

When the cervix is removed the vault of the vagina is caught with forceps as it is being cut in order to pre-

1. The scissors used in this operation are made by grinding off 3 mm. of the upper blade of the Greg Smith's peritoneal scissors and rounding off the lower blade to a probe point. This rounded point can be easily thrust through the peritoneum and it then follows the space between the peritoneum and the blood vessels without injuring either (Fig. 1).

vent retraction. The stumps of the broad ligaments are grasped and are included in the first several sutures that close the vault of the vagina. Figure 6 makes this plain without further comment. The purpose of this maneuver is to prevent the sagging of the vagina. In extensive operations for cancer this procedure is not feasible, for too much of the broad ligament and vagina are removed to permit approximation.

After the broad ligaments are fastened, as above indicated, a gauze drain is passed into the vagina (Fig. 7) or the entire vagina may be closed without drainage. In

Fig. 1.—Cutting the anterior peritoneal layer of the broad ligament; the blood vessels are not cut with this. A, uterine; B, bladder; C, anterior layer of broad ligament; D, ovarian vessels; E, uterine vessels; F, Fallopian tube; G, ovary; H, tumor.

either event the operator may, if he wishes, place several subperitoneal sutures; this appears desirable sometimes to prevent oozing if there is much loose tissue. I usually prefer to omit it.

After the gauze drain has been put in place the peritoneum is closed over it (Fig. 8). This closes off the

Fig. 2.—Ligating the ovarian and uterine vessels. Each vessel, artery or vein is tied off separately. The peritoneum has been pushed back to expose the vessels before ligating.

peritoneal cavity from the site of operation and precludes the possibility of omental or intestinal adhesions to the stump left from the amputation.

I first used this peritoneal closure with independent drainage below, in cases of hysterectomy for cancer in which injury to the bladder actually occurred or was feared.

By this means in a few hours the general abdominal

cavity is shut off from the bladder region, and, should that organ leak from faulty suture or from traumatic necrosis during the operation, the general abdominal cavity is protected from the possibility of urinary infection.

Dudley illustrates a similar procedure but nowhere mentions it in his text, so that I am not certain whether the point was intended. At any rate the procedure is worthy of repetition.

Fig. 3.—After all the vessels are tied off the uterus and the posterior layer of the peritoneum are cut across.

ADVANTAGES OF THE OPERATION.

An amputation of the uterus done in this manner follows the general rules ordinarily accepted for all other amputations.² Its merit consists in the fact that no clamps are used, hence there is no crushing. Tissue not vascular is not subjected to ligation, therefore there is

Fig. 4.—Suture placed to close the stump of the cervix.

no tissue to necrose and no nerves are compressed and no raw surfaces are left exposed. The peritoneum is included only in delicate coaptation sutures which hold the layers together without causing pressure, consequently there is a minimum amount of pain.

2. A similar technic is used in salpingo-oöphorectomy. Peritoneal flaps are made and the vessels are ligated separately and the peritoneum closed over the stumps. In ovarian tumors, too, the pedicle is treated in like manner. A cuff of peritoneum is made as in appendicectomy, the vessels ligated and the cuff united over the stump of the vessels with fine sutures.

The objections to this operation are that it is time consuming, that ligatures on the isolated vessels are apt to slip, that such a technic can not be used in all cases, and, finally, that the operator must be an expert anatomist. In my hands the operation requires from forty-five minutes to an hour and a quarter (with twenty-five minutes as a minimum time). I have not seen shock worthy of note follow in any instance.

Whether ligatures are apt to slip or not depends on

Fig. 5.—Stump of the cervix closed and the peritoneum being sutured.

how they are put on. Ligatures are more apt to slip when tissues of varying contractility are included in a single ligature because the more elastic tissue tends to withdraw itself from the ligature. The cause of a ligature slipping is the faulty application or the use of unsuitable material.

Hard catgut, chromicized, or dry catgut can not

Fig. 6.—Closure of the vagina when the cervix has been removed. The infrauterine ligament is here seen sutured to the upper end of the vagina.

be made to tie securely about a vessel the size of those in question. I prefer Ramsey County or plain catgut previously soaked in water to make it pliable. The use of the surgeon's knot, too, makes it impossible for so small a vessel to be compressed securely, a point which my colleague, Dr. J. N. Jackson, has particularly emphasized. The ligature must be placed under the eye

so that the operator may know when the vessel is securely compressed. If these precautions are used mass ligatures, interlocking and backstitching will not be required to satisfy the operator of the security of his work, any more than they are in ligations in the amputation of an extremity.

That the operation can not be used in all cases is admitted. In suppurating fibroids and in acute infective

Fig. 7.—Vaginal opening now completely closed; gauze drain inserted to drain peritoneal cavity.

processes it is inexpedient. It has been suggested that large fibroids would make its use impossible. My experience has been that in large tumors the vessels are brought nearer the surface and are more readily reached than in smaller tumors.

The truth of the last objection I admit: The operator must be secure in his knowledge of the anatomy of the region involved. Without such knowledge neither this, nor indeed any other, operation should be undertaken.

Fig. 8.—Sagittal section of complete hysterectomy. Subperitoneal space drained by gauze (G. D.) into the vagina.

Adjustment of Spectacle Frames for Infants.—Mr. Bishop Harriman (*Brit. Med. Jour.*, Sept. 12, 1908, p. 745) demonstrated in the ophthalmic section of the British Medical Association an ingenious mode of fastening spectacle frames in infants, which he has used for over a year with excellent results. Children require strong frames, while curl sides must necessarily be frail. He used the ordinary straight arms with a loop at the end. A piece of tape, elastic or not, was looped under the nucha, its ends threaded through the eyes of the spectacle bows, then carried on to the vertex, where they are tied together.