TABLE II.
Differential Count of Leucocytes.

No.	Leucocytes. Per cmm.	Lymphocytes.		Large Mononuclear.		Polynuclear.		Eosinophiles.		Mast Cells.	
		Per cent.	No.	Per cent.	No.	Per cent.	No.	Per cent.	No.	Per cent.	No.
1.	18,000	30	5,400	6·4	1,152	60	10,800	2·2	396	1·4	255
2.	22,000	32	7,040	8	1,760	57	12,540	2	440	1	220
3.	18,300	52·6	9,626	6	1,098	38	6,954	1·2	219·5	2·2	402·5
4.	10,000	37	3,700	10	1,000	50	5,000	1·8	180	1·2	120
5.	20,000	46	9,200	2·8	560	45·6	9,120	5	1,000	·6	120
6.	14,500	45·4	6,583	5	725	44	6,380	4	580	1·6	232
7.	19,000	44·4	8,436	3·6	684	44	8,360	6·6	1,254	1·4	266
8.	22,000	48·6	10,692	4·6	1,012	38	8,360	8	1,760	·8	176
9.	18,000	43	7,740	9·8	1,764	21	3,780	25·8	4,644	·4	72
10.	16,000	60	9,600	4·2	672	21	3,360	9·2	1,472	5·6	896
11.	21,500	44	9,460	4·8	1,033	34·8	7,480	11	2,365	5·4	1,162
12.	17,000	47	7,990	5·4	918	39·2	6,664	6·4	1,088	2	340
13.	22,000	60	13,200	2·6	572	34·2	7,524	3	660	·2	44
14.	12,000	65·8	7,896	1·6	197	26·2	3,144	3·8	466	·8	96
15.	16,000	51·4	8,224	4	640	40	6,400	4	640	·6	96
16.	20,000	79·8	15,960	2	400	13	2,600	4·8	960	·4	80
17.	21,500	57·4	12,341	2	430	35	7,525	5·4	1,161	·2	43
18.	17,000	45	7,650	2·4	400	47·4	8,058	5	850	·2	34
19.	9,500	68·8	6,536	1·6	152	25	2,375	4	380	·6	57
20.	25,000	54	13,500	4	1,000	33	8,250	8	2,000	1	250
21.	19,600	67·4	13,210·4	·8	156·8	23·6	4,625·6	7·8	1,528·8	·4	78·4
22.	16,500	49	8,085	2·6	429	40	6,660	6·6	1,089	1·8	297
23.	23,500	53	12,455	3	705	36·4	8,554	7	1,645	·6	141
24.	24,500	50·6	12,397	2·8	686	38·8	9,506	—	1,568	1·4	343
Max.	25,000	79·8	15,960	10	1,760	60	12,540	11	2,360	5·6	1,162
Min.	9,500	30	3,700	·8	152	13	2,375	1·2	180	·2	34
Av.	19,000	51·6	9,400	4·6	715	37	6,820	5·2	1,050	1·3	245
2nd Av.	—	—	9,890	—	874	—	7,030	—	988	—	247

INDIAN EQUINE PIROPLASMOSIS.

By CAPT. A. J. WILLIAMS, F.R.C.V.S., A.V.C.

History.

THIS disease has been known in India for a considerable time under the name of bilious fever, and in South Africa an identical disease was commonly known as biliary fever. According to Hutcheon, it was described by veterinary surgeons in Natal, as far back as 1883, as the biliary form of horse-sickness and as anthrax fever; Hutcheon also states that, although it strongly resembled horse-sickness, he was convinced that it was of a distinct origin and amenable to different treatment.

In the same article Bowhill states that, as far as he was able to ascertain, Hutcheon's conclusions are accepted by all recent writers except Edington, who divides horse-sickness into two forms, one of which he describes as a non-virulent form with a malarial type of fever and in which parasites are found.

Theiler admits no co-relation between horse-sickness and equine

malaria; he considers that the diseases are distinct in origin, that equine malaria is a typical piroplasmosis, and that piroplasmosis equi is represented by a similar variety of forms to that seen in redwater.

Bowhill mentions a case showing typical lesions of horse-sickness with the presence of *piroplasma equi* in the blood, pointing to a double infection, and suggests that this may explain how Edington was able to infect clean horses with the blood of horses salted against horse-sickness, and find parasites exceedingly like some of the *pirosoma bigeminum* parasites found in Texas fever, and how that author was led to consider the so-called biliary fever and the bilious form of horse-sickness to be one and the same disease.

The work carried on by Theiler, which resulted in the discovery of an endoglobular parasite in all cases of so-called biliary fever of South Africa, has placed the classification of the disease on a more secure footing, and there is no doubt that the disease known as biliary fever is caused by a special endoglobular parasite which, according to Laveran and Theiler, belongs to the genus *piroplasma*, being closely related to the *piroplasma bigeminum* and *piroplasma canis*. In publications previous to June 1903 Theiler opined that the parasite belonged to the genus *hæmamœba*, and he called the disease equine malaria.

The disease being of the red blood corpuscles, and due to a specific parasite infecting them, the name indicates its nature. Although certain forms of the parasite simulate those seen in æstivo-autumnal fever in man, they differ from the parasite of human malaria in the method of reproduction, and up to the present a pigmented parasite, as seen in human malaria, has not been described. I therefore think the term *piroplasmosis equi* the better name for the disease.

During the South African campaign I had considerable experience with biliary fever, extending over a period of two and a half years. The symptoms (which I will describe later when comparing the disease with the Indian form) were typical in character, and deaths were usually the result of extreme exhaustion following work in a tropical sun under conditions in which even healthy horses suffered to a certain extent.

It is easy to explain why the percentage of deaths from biliary fever during the South African war was high. The animals affected were principally imported ones, often sent into the field within a short time of landing. Being unacclimatised, they were in a state which rendered them very susceptible to disease, and, once contracted, the disease had every chance of making rapid progress upon a system which was in no way prepared to fight against it.

Soon after my arrival in Rajputana, India, from South Africa, in November 1902, cases of the so-called bilious fever appeared among the artillery remounts, which were Australian horses. The cases presented symptoms similar in every way to those of the biliary fever cases which I saw during the South African campaign, and I was therefore led to believe that the diseases were identical.

It is my intention to confine myself as far as possible to the Indian type, as it occurs in Central India, Rajputana, and Gugerat, comparing it with the South African form as I met with it, and as described in various articles written since the campaign.

Prevalence, Climate, etc.

My experience of the disease has been gained principally in Central India and Rajputana, but it also appears to be prevalent in Bengal, Punjab, Bombay, and probably also in Burmah.

Regarding the disease in Burmah, Major Eassie, D.S.O., A.V.D., in an article on "Some Observations on Tropical Biliary Fever" (*Journal of Comparative Pathology and Therapeutics*, June 1905), states: "During over three years' service in Burmah I had a large experience of a disease, of the nature of which I was at the time ignorant, both in indigenous ponies and in mules imported from the Punjab and from Western China. Looking back on it now, I can come to no other conclusion but that it was a form of equine piroplasmosis which I have described as recurrent."

In the districts mentioned as the parts in which my experience has been gained, the disease was believed to be due to climatic changes, and to an unhealthy state of the country following the rainy season. In the autumn the sudden changes and cold winds were blamed, these being supposed to cause chills which affected the liver. Too rich a diet during the hot weather, insufficient exercise, etc., were also thought to take some part in the production of the disease, and the name bilious fever suggests that the affection is due to errors in diet, or to a chill affecting the liver and producing a jaundiced febrile condition.

The disease chiefly occurs during the summer and early autumn months, May, June, July, and August, but sometimes, as in 1905, cases continue up till the end of October. In that year there was a failure of the rain in Rajputana; only 5 inches fell during the twelve months, the result being a continuation of the hot weather much later on in the year than is usual when a normal rainfall occurs.

It is interesting to note that in districts in which I have met with the disease, human malaria is very prevalent, hardly anyone in the stations escaping an attack during a particularly bad year. This was the case at Deesa, Rajputana, in November 1902, the rainy season that year being the first good one for about five years, with the result that malaria was extremely prevalent among the Europeans and natives. I think I am correct in saying that nearly 50 per cent. of the men of the battery and a large number of the natives were ill from malaria. A season of that kind is bad for man and beast, and as a rule a higher percentage of equine malaria cases then occur.

I wish to point out that the disease is more especially one affecting Australian remounts; cases among Arabs and country bred are not common. I can recall very few cases among the latter. In the case of Australian remounts 50 to 70 per cent. become affected within a few months of their arrival in the stations.

Etiology.

The disease is due to an invasion by an intra-corpuscular parasite, identical with the piroplasma equi discovered by Theiler during his researches in connection with biliary fever of horses in South Africa.

In a very interesting article by Lieut. Pallin, A.V.D., on "Biliary Fever of Horses in India" (*Veterinary Journal*, June 1905), there

is a Plate which illustrates very well the various forms of the piroplasm found in the red blood corpuscles. He states that the smears were taken at different stages from a number of typical biliary fever cases, that in all cases some of the red corpuscles contained an organism of the nature of a piroplasm, and that the micro-organism cannot always be found during the whole course of the attack. They were more numerous on some days than others, and there were never more than three or four affected blood corpuscles in a field.

The smears I have taken from typical cases show signet-ring forms, pear-shaped forms, and distinct spherical forms in which the protoplasm is a pale blue tinge, and the chromatin dot stained a deep red and lying in the centre of the sphere. This last form is not described by Pallin, and Theiler states that the karyosome is never in the centre but always close to the edge. These spherical forms with a central karyosome appear very similar to the spherical forms described as occurring in piroplasmosis of the donkey (Dale).

In blood films stained with Romanowsky or Leishman's stain the parasite can be demonstrated fairly easily, using a one-twelfth objective and No. 2 or No. 3 eye-piece.

The number of parasites to be met with in the peripheral blood is proportional to the severity of the disease, and from 1 to 10 or up to 30 per cent. of the corpuscles may be affected.

In the spleen the number of affected corpuscles may be greater, and in the liver still more, while in the kidney 60 per cent. of the corpuscles may be infected.

Free parasites are found in the blood in the latter stages of the disease, especially in the kidneys (Dr Annett, Liverpool Lectures).

With the above-mentioned stains the protoplasm of the parasite is stained a pale blue, and the chromatin a deep red colour. Numerous forms have been described. Theiler mentions stave-like, leaf-like, club-shaped, oval, pyriform, round-shaped, and rosette forms. Bowhill mentions large and small spherical forms, large and small pyriform parasites singly and in pairs, large and small rod-like bodies, rosette and flagellated forms. Pallin mentions flask-shaped, ring-shaped, baton-shaped, and small round bodies resembling cocci.

Theiler states that the round and ring forms are very common, that in fresh unstained specimens the parasites appear commonly as the round form; he considers that the real form is a sphere probably, and that the non-spherical forms depend on fixation, staining of the blood, or other influences. He states that the rosette form with division into four parts is the one commonly found, but division into two can also be observed.

With reference to the presence of the parasites at various stages of the disease and the action of quinine and ammon. chlor. on them, he states that the parasite can always be found in the early stages of the attack, that it is not constantly present during the whole course of the disease, and that it is quite possible that one may diagnose a case as one of typical biliary fever and subsequent microscopical investigation fail to reveal the parasite. The administration of quinine and ammon. chlor. results in rapid disappearance of the parasite from the peripheral blood.

In smears which I have taken I have found the following forms:

ring-shaped, pyriform, flask-shaped, oval, and the so-called reproductive forms, which consisted of four coccus-like bodies in one corpuscle.

CASE I.—In this case (*see* the Temperature Chart I.) smears were taken on the second, third, sixth, eleventh, and twelfth days of the disease; they were positive on the second, third, eleventh, and twelfth days, and negative on the sixth day.

The first smear taken on the second day of the disease showed pear-shaped and ring-shaped forms; one of the pear-shaped forms showed three distinct chromatin dots. A pear-shaped parasite free in the plasma was also seen. The smear taken on the third day showed few parasites, chiefly the ring form; two were seen in one corpuscle, and a parasite in which the karyosome was unstained was also observed.

With regard to the parasites showing non-staining of the karyosome, Bowhill states that the colourability of this is an indication of the parasite being alive.

On the eleventh day the smear showed very few ring forms, a few

CHART I.

parasites free in the plasma, a number of intra-corpuscular ones showing no sign of staining, and a large pear-shaped form showing a chromatin dot at either end. On the twelfth day a large number of parasites were free in the plasma, and there were very few corpuscles affected. A very well-marked dividing form was seen, in the shape of a pear-shaped body with a distinct stained mass across the centre and a chromatin dot at each end; this appeared to be in the stage immediately prior to breaking up into two parasites by central division of the protoplasm, and it is the only one I have seen showing distinct signs of division into two parts.

CASE II.—In this case smears were taken on the first, second, third, fifth, eighth, twelfth, and thirteenth days of the disease. The smears were positive on the first and third days only.

In this case the temperature dropped to normal on the eighth day (*see* Chart II.), and remained at normal throughout with the exception of a rise on the fourteenth day, gradually dropping to normal again on the seventeenth day. This rise was associated with constipation, which I concluded was the cause; no smear was taken.

Towards the end of an attack the number of affected cells decreases, but there are more free forms present; parasites free in the plasma are exceptionally seen in the early stages of the disease.

A number of non-staining parasites were noticed, and the piroplasm varied much in size, some being very large and others very small and easily overlooked, especially when they had not taken the stain well.

The blood cells show alterations as follows:—

Red Cells.—Variation in size and altered shape (megalocytes, microcytes, and poikilocytes). Some of them stained much darker than usual, pointing to anæmic degeneration of the protoplasm (polychromatophilia).

Leucocytes.—Mononuclear and polymorphonuclear increase, also a marked increase in eosinophiles in some smears.

Symptoms.

The attack is sudden and manifested by a rise of temperature, usually 104° to 106° F., associated with the usual train of symptoms

CHART II.

occurring with a high temperature, such as loss of appetite, dulness, pulse and respiration increased, constipation, fæces coated with mucus, thirst, etc.

In the majority of cases the animal appears quite well to within a few hours of admission for treatment, the first symptom being complete loss of appetite and dulness.

In a typical case (see Chart II.) I was asked to see the pony because the owner thought he had something wrong with his feet, as he had appeared quite fit when he started out in the morning, but before mid-day was stumbling badly. I saw the animal at 3 P.M., when the temperature was 106.2° , respiration 60, typical appearance of the conjunctival mucous membrane, etc.

The temperature is always high, 105° or 106° F. in a large percentage of cases, and the conjunctival mucous membrane shows a typical appearance. It is reddish-brown in colour, with a tinge of yellow, and a few bright red petechial spots are scattered over the membrana nictitans. These petechiæ gradually increase in size and alter in colour considerably during the first few days of the disease;

they become more of the nature of blotches, and may coalesce to form comparatively large patches, the colour changing through various shades of red to a deep claret; this latter appearance is reached about the fourth or fifth day.

CHART III.

During the first few days the icteric condition increases, in some cases tinging the lips and skin. This is best seen in white or grey animals. This marked icterus appears simultaneously with the attack or within forty-eight hours.

CHART III.—continued.

I wish to lay special stress upon this peculiar condition of the conjunctival mucous membrane, as I have never seen such a condition in any disease but this one and the South African form; it is always present in *piroplasmosis equi* in India and is to my mind characteristic of the disease.

In some cases of catarrhal jaundice a few bright red petechial spots appear on the conjunctiva, but they do not show the marked changes in colour and extension over the mucous membrane that is seen in piroplasmosis. The characteristic reddish-brown colour is absent in jaundice, and there is not such a marked rise of temperature. Neither is there a very marked fall within about forty-eight hours, as usually occurs in piroplasmosis.

In piroplasmosis the temperature in the majority of cases is at its highest point at the commencement of the attack, and it is only in very rare cases that it again reaches to within 2° of the initial temperature during any period of the illness.

In many cases the temperature drops a degree or two within twenty-four hours; the third day frequently shows a well-marked fall, often two or more degrees, and in some cases a fall to within a

CHART IV.

few points of normal. The latter condition I have noticed after treating with large doses of quinine.

A big fall is followed by a well-marked rise, an intermission for a few days, then a gradual drop to normal, which is reached about the sixth or tenth day in mild cases.

In severe cases the fever is of an irregular intermittent character, and may not disappear within fourteen to twenty-one days.

In very severe cases the temperature occasionally drops to normal, or it may be subnormal during the period of intermissions, and these cases are also occasionally marked by distinct rises of temperature at a few days' interval; for example, a rise on the second, fifth, tenth, and thirteenth days, as seen in Chart III.

To summarise the character of the fever, I may say that it is of an irregular intermittent type, higher at the onset than at any other period of the disease. In mild cases there are few intermissions, and the normal is reached in from six to ten days; in severe cases remissions are fairly frequent, and the normal condition is not reached till about the fourteenth to twenty-first day.

The pulse is usually 60 to 80 per minute, and of a soft character,

improving in tone and showing a marked decrease in frequency corresponding to a fall of the temperature.

The respiration is always much increased ; at the onset it may be 60 or more, decreasing rapidly with a fall of temperature and signs of general improvement.

The urine is dark in colour, but I have never seen a case showing hæmoglobinuria.

Course and Complications.

Recovery is the rule, and convalescence is short, loss of condition not being very marked except in severe and complicated cases.

The disease occasionally recurs in the following year. In these instances the initial temperature is not usually as high as in primary attacks, there are fewer intermissions, and the case progresses in a similar way to a mild primary attack. The convalescent period is short, loss of condition is slight, and the animal is able to resume work within a few days after the temperature reaches normal.

One attack appears to confer a certain degree of immunity.

Pneumonia is the commonest complication, and it usually occurs during the first few days of the disease. It is of the lobar form and presents the usual clinical symptoms ; it is marked by a very distinct rise of temperature and increased respiration, auscultation and percussion showing the usual clinical signs. The right lung is affected oftener than the left, and the lesions are more frequently unilateral than bilateral. The disease is often localised and confined to the middle and lower regions of the lung.

Intestinal complications are less frequent ; they are manifested by sub-acute abdominal pain, constipation, an increased icteric condition, fæces thickly coated with mucus and of a yellow colour—a muco-enteritis in fact.

In a few cases I have noticed a peculiar condition with the general symptoms of intestinal affection as described above, but in which the sub-acute abdominal pain continued for forty-eight hours or more, the animal looking to the sides and placing the muzzle close behind the elbow, pointing to an affection chiefly of the small intestines. These symptoms I put down to a muco-enteritis affecting the small intestines, probably with duodenal obstruction due to disturbance of the liver functions.

In one case I saw well-marked enteritis associated with a dysenteric diarrhœa of an extremely fœtid character ; the temperature rose steadily from 101·2° F. on the ninth day to 106·4° F. on the thirteenth day, the animal dying on the latter date, thirty-six hours from the time enteric symptoms first appeared.

Having had very few deaths, I have little to say regarding the *post-mortem* appearances, and think I cannot do better than give a short outline of the case just mentioned, in which death occurred on the thirteenth day. I include this case under the heading "complicated cases," but I am unable to decide whether the condition leading to death was really an exceptionally acute piroplasmosis or piroplasmosis complicated by a dysenteric condition.

Subject.—Chestnut Australian mare, polo pony, five years old (see Chart I.).

History.—The pony appeared quite well on the 5th August 1905 ;

on the 6th August at 7.30 P.M. she was noticed to be extremely dull, with drooping head and breathing very rapidly.

I saw the case at 9.30 P.M. The symptoms were as follows: Temperature 105.6° F., pulse 60, respiration 60, icterus, extreme dullness, loss of appetite, etc. Petechial spots appeared on the conjunctiva in forty-eight hours, and the usual symptoms developed.

The jaundice gradually cleared off, conjunctivæ on the eighth day clean, fæces normal, feeding well. On the twelfth day diarrhœa set in, watery, very fœtid, temperature rose from 102° to 104.6° , diarrhœa continued all day, extreme dullness, refused all food, tossing head up and down. On the thirteenth day much worse, temperature 106.4° F., diarrhœa very bad, blackish and extremely fœtid, pulse 80 and very weak, subacute abdominal pain, anxious expression, eyes glassy and staring. The fœtid fæces poured from her, but there was no tenesmus; condition grave, prognosis death.

I administered anodynes, astringents, starch and opium enemas, morphia, etc., but the patient gradually grew worse, running down pulse, collapse, quivering of the muscles of the hind quarters and hind legs, pupils dilated, delirium, wandering about the box pressing head against the walls and pawing. Died at 1.30 P.M.

Post-mortem appearances.—The large and small colon and part of the rectum were the seat of an exceptionally acute enteritis, the mucous membrane being blackish-green in colour and easily removed with the fingers, and the submucous coat about an inch thick and filled with a jelly-like straw-coloured exudation, blood-stained in parts. No ulceration of the mucous membrane. Some peritonitis, intestinal lymphatics congested and enlarged, diseased portion of intestines filled with extremely fœtid watery ingesta. Other parts of the intestinal tract appeared healthy and contained normal fæces. Liver congested and fatty, capsule easily stripped off; the liver substance could be easily pulped between the fingers.

This was an exceptional case. Up to the tenth or eleventh day there was no reason to suppose that the animal would not recover, although the attack was a very severe one from the onset. The very sudden appearance of the enteritis, the steady rise of temperature notwithstanding the weak pulse and other symptoms of collapse, pointed to a very severe toxæmia. The extreme fœtor of the evacuations, the abdominal symptoms, low delirium, etc., and the intestinal lesions confined to the large intestines, all point to a condition similar to gangrenous dysentery in man.

Treatment.

Very important factors in the treatment are, the prevention of affected animals doing any work during the early stages of the attack, protection from the sun, and good nursing.

With reference to the first-mentioned; in South Africa where I treated a large number of cases under active service conditions I noticed that if one could get animals under treatment within a short time of the first symptoms appearing recovery was the rule, but cases which had been worked all day in the sun, with a temperature of 106° F. probably, were in such an exhausted condition that treatment was of no avail, and the animal would drop from exhaustion even though moving at a very slow pace in rear of the column.

The disease appears suddenly ; the rider would not notice anything wrong until he had ridden the horse some hours, and then the animal would stumble along and reach camp in an exhausted state.

I have often seen men spurring along an animal barely able to crawl, have stopped them, examined the animal, and found the temperature 105° or 106° F. ; in those cases the death rate was high.

Cases amenable to treatment were those which as soon as symptoms of exhaustion appeared were off-saddled, led quietly into camp, put under treatment as early as possible, and marched along quietly in rear of the column.

Eassie, in the article already referred to, states that he is convinced that exposure, without the option of shade, to a tropical sun over a sufficient period, is a determining factor both in the appearance and intensity of primary equine piroplasmosis and to even a greater degree in its recurrence.

There is no doubt that protection from the sun is a very important matter in the treatment, as it is in other fever cases. An animal exposed to the sun and exhausted suffers more severely than an acclimatised well-looking-after animal ; when exposed to the sun and in an exhausted condition, the state of the system is such as to favour the development of any disease.

Cases occur on active service and at other times when it is impossible to do as much as one would wish towards treatment, but one can often fix up a temporary shelter, and measures should be adopted which would lead to the early reporting of cases.

When an animal is off feed the temperature should be taken, and thus avoid the working of an animal in the very early stages ; when possible, affected animals should be protected from the direct rays of the sun. Good nursing, careful dieting, plenty of water, and good hygienic surroundings are important items, as in the treatment of any fever case.

I keep the animals on bran mash diet until the period of convalescence is reached, gradually bringing them back again to their ordinary food. If the animal dislikes bran I give a small amount of steamed oats with it. Fresh lucerne is an excellent food, easily digested, and animals will usually eat it in preference to anything else.

Medicinal Treatment.

Quinine in large doses at the onset, decreasing the dose after the first few days, salines and tonics during the convalescent stage, is the line of treatment I have found successful. I give quinine in 2 drachm doses two or three times a day during the first few days ; mag. sulph. 4 oz. three times a day, and 4 drachms ammon. chlor. once daily, in the drinking water. The following are useful drinks, viz :—

R.

Quiniæ. Sulph.
Ammon. Carb. aa ʒij.
Mag. Sulph. ʒiv.
Aquam ad Oj.

M. ft. haust.

Sig. One three times a day.

R

Quiniae. Sulph. $\bar{z}$ ij.
 Spts. Aeth. Nit.
 Spts. Ammon. Aromat. aa $\bar{z}$ j.
 Mag. Sulph. $\bar{z}$ iv.
 Aquam ad Oj.

M. ft. haust.

Sig. One three times a day.

I continue the mag. sulph. in 4 oz. doses three times a day until an aperient action is obtained. When the fever shows signs of abating I give quinine 1 drachm and ammon. carb. 2 drachms in bolus twice daily, and continue the ammon. chlor. in the drinking water.

During convalescence I give the following balls, and in addition liq. arsenicalis $\frac{1}{2}$ oz. once daily for a week or so, in cases where the attack has been a severe one.

R

Quiniae. Sulph. $\bar{z}$ fs.
 Ferri. Sulph.
 Ammon. Carb. aa $\bar{z}$ j.
 Common Mass. q.s.

M. ft. bolus.

Sig. One morning and evening.

I have tried calomel, but found it uncertain in its action, and consider mag. sulph. the best aperient in these cases.

In cases complicated with pneumonia I treat medicinally as above, and apply mustard to the sides immediately lung symptoms appear.

I treated intestinal cases according to the symptoms shown, using ol. lini. as an aperient and cannabis indica where a sedative was required, care being taken to avoid purgation. In severe cases showing a relapse I have often applied a mustard plaster over the region of the liver with excellent results, the temperature showing a decided fall, and the animal improving generally within twenty-four hours.

Very little appears to have been done regarding the investigation of the disease in India; the method of transmission is not known. The period up to which parasites can be found in the blood after symptoms of the disease have disappeared, and the action of quinine on the parasites with regard to sporulation, etc., would be valuable information, while the discovery of the method of transmission might give results which would lead to the solving of an important question, viz., the development of the parasite.

It is interesting to note that in India the disease occurs among a class of animals which are, as a rule, free from ticks. There are ticks in the districts, but the animals are never tick infested as one understands the term. A tick or two are occasionally found on the animals, but they are so well cared for that when any ticks do find their way on to them they are quickly removed. This refers to army remounts and polo ponies, among which my experience has chiefly been obtained.

There is no condition of tick infestation as is seen in South Africa, where the animals become thickly covered with them, and, reasoning from this, it is possible that the disease in India is transmitted by blood-sucking flies or mosquitoes.

In all other piroplasmoses the transmission of the disease has been traced to various ticks, and in all cases the animals are infested with ticks.

Theiler has discovered that the South African form of the disease is transmitted by the *rhhipicephalus evertsi* in its nymph and adult stage, and it is presumed that there are two stages of development, viz., the schizogonic stage in the host, and the sporogonic stage in the tick.

The discovery of the method of infection in India would be of great interest and a valuable addition to our knowledge of a group of diseases which, during the last few years, have attracted considerable attention on the part of veterinary surgeons, especially those in tropical countries.

REFERENCES.

Theiler. Equine Malaria and its Sequelæ. *Journal of Comparative Pathology and Therapeutics*, June 1903.

Dale. Piroplasmosis of the Donkey in South Africa. *Journal of Comparative Pathology and Therapeutics*, December 1903, and *Veterinary Journal*, June 1904.

Bowhill. Contribution to the Study of Equine Malaria. *Veterinary Journal*, July 1905.

Eassie. Some Observations on Tropical Biliary Fever. *Journal of Comparative Pathology and Therapeutics*, June 1905.

Pallin. Biliary Fever of Horses in India. *Veterinary Journal*, January 1905.

Bowhill. *The Journal of Hygiene*, January 1905.

Annett. Lectures, Liverpool University, 1906.

OSTEOPOROSIS IN ANIMALS.

By HERBERT INGLE, B.Sc., F.I.C., Chief Chemist, Transvaal
Department of Agriculture.

IN this paper a new theory as to the predisposing cause of this disease is enunciated, solely from chemical evidence, and, while the writer regrets that he is not able to furnish any experimental proofs of the success which might attend the treatment which the theory would recommend, he trusts that by the time this paper is published he may, by the co-operation of Dr Theiler, be in a position to furnish such facts as will lead to its confirmation.

The disease is only too well known in some districts of the Transvaal, and therefore perhaps needs no description; but, in any case, it is not in the writer's province to attempt to describe its symptoms or effects, even were he sufficiently familiar with them to do so.

Neither is it in the domain of the chemist to discuss the actual direct cause of the disease, or the character of the micro-organism, if any, to which it may be due. These points must be left to the veterinary bacteriologists and veterinary surgeons. The present communication deals exclusively with the chemical aspect of the question.

The disease, which is very destructive to horses, mules, and