INHERITANCE IN SWINE

JAY L. LUSH

Department of Genetics, University of Wisconsin, Madison

ONE among the many casualties of the Great War, which will not be found listed in the files of the War Department, was the stopping of a certain extensive experiment in swine breeding which was conducted at the Kansas Experiment Station under the direction of Mr. E. N. Wentworth until the spring of 1918 when the last of the persons in immediate charge of the work entered military service. That ended the taking of detailed records and the subjects of the experiment continued their growth, matured, and were marketed, so that it was impossible to resume the experiment at the point at which it stopped.

SOME VALUABLE CONCLUSIONS REACHED

However the experiment was not a complete casualty for the results already secured in four years of breeding make a few conclusions almost inescapable and point to others which can perhaps be verified, one at a time, by a few well-chosen crosses. This is particularly fortunate because there have been so few genetic experiments on swine which have gone farther than the production of a very few F_2 litters, and yet swine are very suitable for studies in heredity, not only because of their economic importance, but also because of the rapidity of their multiplication and the wide diversity of true-breeding forms and colors which they offer to the investigators.

The experiment was undertaken with the idea of securing information about the inheritance of all the well-defined characters which differed in the breeds used, but conclusions have been reached only in regard to the shape of face, set of ears, color, growth factors, and litter size. The most of the work was done on crosses of a registered

Berkshire boar on registered Duroc-Jersey and Tamworth sows, but the problem of litter size was separately attacked by mating European wild boars of the Schwarzwald type to Tamworth and Berkshire sows.

LITTER SIZE

The use of the wild boar for investigating this economically important character was originally suggested by an investigation by Wentworth and Aubel¹ in which it appeared that the distribution of the various litter sizes among ordinary domestic swine did not conform to the normal frequency curve. Since the wild hog ordinarily produces four pigs per litter, it was decided to cross a boar of this kind upon sows of one of the more prolific of the domestic breeds, such as the Tamworth which ordinarily produces about eleven pigs per litter. There were thirty-eight F_1 pigs produced in this cross, but only one of the females among them had reproduced before the experiment ended. The fact that her litter consisted of but four pigs, taken in connection with the fact that Simpson² reports litters of four, six, and four, respectively, from three sows of the same kind of breeding, gives an indication that the wild litter size is dominant, but tells us no more about its inheritance.

SET OF EARS

The Berkshire, Tamworth and wild hog all have erect ears of fine or moderately fine texture, and since all the pigs produced by crossing these three breeds possessed a very similar sort of ear, they throw no light upon its inheritance. The Duroc-Jersey, however, possesses a distinctly different sort of ear which is not so pointed, is of medium size, and breaks over at

¹ WENTWORTH, E. N. and AUBEL, C. E. Inheritance of Fertility in Swine. *Journal of Agricultural Research*, Vol. 5, No. 25, pp. 1145-1160.

² SIMPSON, Q. I. 1912. Fecundity in Swine. *Am. Breeders' Association*, Vol. 7, pp. 261-275.

SOME WEANLING PIGS FROM A WILD BOAR FATHER AND POLAND CHINA MOTHERS

The swine breeding experiments described in this article were conducted to determine the inheritance of certain well-defined characters which differed in the several breeds used, such as color, shape of face, set of ears and litter size. The economic importance of some definite knowledge regarding rate of growth as well as the greater size and vigor of the hybrids for market purposes was recognized throughout the experiments and some significant data secured. Photograph by courtesy of the Iowa Experiment Station. (Fig. 6.)

about one-third of the distance from the tip to the base, so that the outer third droops downward. There is quite a bit of variation within the breed but this is closely associated with variation in the general quality of the animal. That is, the coarser ear more than one-third broken over is usually found on the coarser, rougher animals, while the finer, more pointed ear, broken over less than one-third, is found more often on a lighter-limbed, rangy type of hog. Whether this correlation is genetic (linkage) or due simply to the mechanics of growth is not known.

The twenty-nine F_1 pigs of the Berkshire $\times$ Duroc-Jersey cross, except for one boar and one sow in the first litter (which was sired by a boar not used subsequently, and which was discarded for breeding purposes), all possessed ears which were perfectly erect but slightly larger than on Berkshires of equal size.

Only forty-two F_2 individuals were mature enough to determine definitely the shape of the ear when the work was discontinued. Of these, only one showed a typical Duroc-Jersey ear, several were intermediate, but the great majority possessed ears as erect as the Berkshire.

The F_1 boar when mated back to his dam, sired three litters totalling thirty-five pigs, nine of which had lived to be as much as six months old when the experiment ended. Of these, four had typical Duroc-Jersey ears, one was still undetermined, and four had erect ears.

The following conclusions in regard to the inheritance of the set of ear seem warranted by the data: The typical erect ear of the Berkshire is dominant by at least one and probably not more than three (F_2 ratio of forty-one erect or intermediate to one of Duroc-Jersey type) principal factors. There may be a number of minor modifying factors for size and quality as well as for the amount of breaking over. Neither breed is homozygous throughout for all the factors concerned in the

A WILD BOAR AND ONE OF HIS PROGENY FROM A DOMESTIC TAMWORTH SOW

European wild boars were used in these breeding experiments to determine the inheritance of litter size. Wild sows ordinarily produce only about four pigs per litter which is much less than is produced by sows of the domestic breeds. Tamworths farrow on the average about eleven pigs. The picture shows the wild boar used (at the right) with one of his crossbred sons. (Fig. 7.)

production of its own peculiar ear shape; i.e., being purebred is not equivalent to being homozygous in this respect.

SHAPE OF FACE

This character, like the set of ear, does not complete its development until the animal is mature, and is fairly constant within each of the breeds of swine. Both the Tamworth and the wild hog are characterized by long narrow straight faces with no forehead prominence. The Wild face is the more extreme in each of these three respects, and the seven mature F_1 pigs all approach very closely to the face shape of their wild parent.

Among the common breeds of swine in America the Berkshire represents the opposite extreme to the wild hog in

face shape, having a very short, extremely dished face and a forehead so broad and prominent as to give it a sort of pompadour appearance. Only two pigs from the Wild $\times$ Berkshire cross matured and both resembled the wild parent so closely that, except for color and wider foreheads, they would have been indistinguishable from the Wild $\times$ Tamworth F_1 's.

Seven pigs of the F_1 generation of the Berkshire $\times$ Tamworth cross matured and could not be distinguished from the purebred Tamworths as far as face shape was concerned. This is especially surprising because Simpson³ who crossed Tamworths and Large Yorkshires (which have a face shape similar to the Berkshire in outline although longer), reported that the result was an intermediate face shape. He does

³ SIMPSON, Q. I. AND J. P. 1909. *Am. Breeders' Association*, Vol. 5, pp. 250-5.

A TYPICAL BERKSHIRE BOAR

In addition to the wild hog, the Berkshire, Duroc-Jersey and Tamworth breeds were used. A photograph of the registered Berkshire boar used is not now available, but the above illustration shows a typical boar of this breed. Its ears are erect like those of the Tamworth and wild breeds. Photograph from U. S. Dept. of Agriculture. (Fig. 8.)

not give the numbers for an F_2 generation and none of the F_1 individuals in this experiment reproduced, hence there is not even an indication of the number of factors responsible for the face differences in these three breeds. However the data clearly indicate this much: Tamworth long straight face is dominant over Berkshire short dished face; Yorkshire dished face forms an intermediate F_1 with Tamworth long straight face; therefore Berkshire short dished face and Yorkshire dished face must be similar phenotypes produced by somewhat different factor complexes.

The face shape of the Duroc-Jersey is intermediate to that of the breeds just discussed in that it is moderate in length and in dish, while the forehead and the hair which grows upon it usually slope smoothly upward and backward. In the F_1 generation of the Berkshire $\times$ Duroc-Jersey cross,

the forehead in every case approached closely that of the Berkshire type and the face was strongly dished but not quite to the degree of the Berkshire ideal. There was some variation in dish and in length of face among the F_1 s but they were all pronounced enough to be more Berkshire than Duroc-Jersey in type. In the F_2 generation very much wider variation was observed. Since no practicable means of measuring these three elements of face shape was discovered, they were simply classified according to their resemblance to one or the other of the grandparental types. Table I shows the result:

TABLE I

Character	Similar to Berkshire	Intermediate	Similar to Duroc-Jersey
Forehead Shape	37	2	3
Dish of Face	17	9	16
Length of Face	21	7	14

THE PUREBRED DUROC-JERSEY WHICH WAS THE MOTHER OF THE FIRST F₁ AND BACKCROSS GENERATIONS

The Duroc-Jersey breed possesses drooping ears in contrast to the erect ears of the other breeds used. All except two of the 29 pigs of the first generation from the Berkshire $\times$ Duroc-Jersey cross had erect ears. (Fig. 9.)

Of course it is likely that some of these groups may contain more than one phenotype, since the characters are not definitely measurable ones, but the variation from one extreme to the other was by no means continuous and the investigators believe that the number of phenotypes which result from this cross is quite limited. There was some evidence of correlation between length of face and dish of face because hogs with both extremely dished faces and extremely long faces did not occur, but on the other hand, short-faced hogs occurred with both extremely dished and extremely straight faces. This probably means that such correlation as may exist is developmental or anatomical rather than genetic (linkage).

Other conclusions which may be drawn with respect to face shape are these: Several factors differing in importance and in degree of dominance are responsible for the anatomical differences which have been considered

under the title of face shape. Certain of these factors affect certain parts of the face much more than they do other parts. The number of these factors can not be so very large, else the parental types would not have been recovered among such limited numbers.

SWINE COLOR

The carefully controlled experiments on swine which throw any light upon the inheritance of the various colors are not numerous and the resulting interpretations which are established beyond any scientific doubt are few indeed. The whole subject has been reviewed recently in the *Journal of Heredity* by Wright⁴ and, as far as the author knows, since then there has been only one notable addition to the literature upon the subject;⁵ therefore what seems most likely to be the truth about the inheritance of the different colors found in swine will be very briefly summed up as follows:

⁴WRIGHT, SEWALL. 1918. Color Inheritance in Swine. *JOURNAL OF HEREDITY*, Vol. 9, pp. 33-38.
⁵LLOYD-JONES AND EVVARD. Iowa Agricultural Experiment Station, *Research Bulletin*, No. 53.

A CROSSBRED FROM THE BERKSHIRE $\times$ DUROC-JERSEY CROSS

This boar was the father of the second and backcross generations. Compare it with its parent breeds shown in Figs. 8 and 9 and note its color, erect ears and intermediate face shape. (Fig. 10.)

White: There are at least three kinds of white which are genetically distinct. Self white is dominant over all other colors with which it has been crossed and is probably dependent upon only one factor. White belt (as seen in the American Hampshire or in the German Hanoverian swine) is usually dominant but must depend upon at least two factors since belted offspring sometimes result from non-belted parents. The white which is best illustrated in the "six white points" of the Berkshire or Poland-China, seems to be an extremely diluted red or the absence of all shades of red. The author has found no record of true albinism in swine.

Black: There are certainly two and very probably three kinds of black. Self

black (as in the English Large Black swine) is reported dominant to Tamworth red. The black which is associated with the white belt in the American Hampshire is dominant to Duroc-Jersey red and depends upon a single factor. There is no evidence as to whether these two kinds of black are genetically distinct, or the latter is merely self black plus the additional unknown factors for white belt. The black of the Berkshire and Poland-China is certainly distinct from both the preceding kinds since it always breaks up, when crossed with red, into numerous black spots scattered over a lighter background, usually red or sandy but sometimes white.⁶ This black, as will appear from the results of this experiment, is caused by a

*SEVERSON, B. O. 1917. Color Inheritance in Swine. JOURNAL OF HEREDITY, Vol. 8, p. 379.

A BERKSHIRE X DUROC-JERSEY CROSS SHOWING A SHORT STRAIGHT FACE (Fig. 11)

ANOTHER BERKSHIRE X DUROC-JERSEY CROSS

This is a first generation boar with drooping ears—one of the two out of the 29 first generation pigs which did not have erect ears. (Fig. 12.)

A SECOND GENERATION BERKSHIRE X DUROC-JERSEY CROSS

An illustration of the appearance of the Berkshire type of forehead in the second generation. Contrast with the different type shown in the following figure. (Fig. 13.)

single factor which alone produces black spots irregularly scattered over a lighter ground color and, when extended by the action of various modifying factors, it covers all the body except the "six white points." Reasoning from the similarity in appearance and ancestry rather than from any definite experimental data, the Spotted Poland-China possesses the same fundamental factor for black spots but many of the factors which influence the size of those spots are absent.

Red: Two kinds of self red are genetically distinguishable by their behavior toward self white. Possibly the factors which affect the white are distinct from the factors for red. Differences in the intensity of the red seem to be inherited but there are no experimental data to back up that popular opinion.

Roaning: Appears quite frequently among the descendants of self whites

by self reds, or of self whites by belted blacks. Often it is not apparent at birth but develops later.

Immature Striping and Adult Agouti of the Wild Hog: This pattern is dominant over all other colors except self white and the white belt, but the dominance is not always complete.

COLOR RESULTS FROM THIS EXPERIMENT

The European wild hog is of a color corresponding very closely to the agouti or gray pattern common among many other wild mammals, except that in this case it is an uncommonly dark slaty shade. Another interesting feature is that the young are born with longitudinal stripes about a centimeter wide running the full length of their bodies and extending over all the back and sides, but with very light bellies. This is quite evidently a protective coloration for at a short distance they blend in very well with the sticks and trash on the ground. The contrasting

A BERKSHIRE X DUROC-JERSEY CROSS

An example of the re-appearance in the second generation (Berkshire $\times$ Duroc-Jersey) of a long straight face similar to that of the Duroc-Jersey. The first cross was closer to the Berkshire than to the Duroc-Jersey in length and dish of face and shape of forehead. Various combinations segregated out in the second generation. (Fig. 14.)

stripes were brown and sandy on all the thirty-eight pigs produced by crossing the wild boars upon purebred Tamworth sows. They were quite uniform in color. However, on the seventeen pigs produced by crossing the wild boars on purebred Berkshire sows, the bellies and light stripes were a very light sandy, almost white, and the darker stripes were a darker brown; that is, there was less of a reddish tinge throughout and, in addition, there were large black spots scattered all along the whole under surface. This is most readily interpreted as meaning that the wild pattern is not completely epistatic but lets the Berkshire black spots develop along the underline and allows the Tamworth red to show up in the reddish tinge of the offspring from that cross.

Four F_2 pigs, all striped but of different belly colors, were produced in the Wild $\times$ Tamworth cross. One had a very light, almost white, belly (presumably lacking the Tamworth factor or factors for red), two were like the F_1 s, and one was like the F_1 s except that it had some large black spots in both hair and skin scattered along its belly. There was one backcross litter by an F_1 boar of a purebred Tamworth sow. Three of these were faintly striped and possessed small black spots on their bodies, two were like the F_1 s, two were red with small black spots, and one was self red. This seems to mean at least two independent factors—one for black spots and one for the striping pattern—and another factor for intensity of the striping which may be linked to the factor for black spots.

THE SECOND GENERATION PIG WHICH SHOWED THE MOST BLACK (BERKSHIRE X DUROC-JERSEY CROSS)

The first generation was composed wholly of red pigs with black spots. There was much variation in the second generation of which the case illustrated above is an extreme example. (Fig. 15.)

Ten pigs were produced in the F_1 of the Berkshire $\times$ Tamworth cross. Eight were red with one or more small black spots and two were self red. The spots were very small and much fewer than in the pigs from the Berkshire $\times$ Duroc-Jersey cross. One had but a single black spot and it seems possible that the restriction factors for black are so strong in the Tamworth breed that the two self red pigs are merely somatic variations in the amount of black and not genetically different from their litter mates.

In the Berkshire $\times$ Duroc-Jersey cross there were produced an F_1 generation of twenty-nine pigs, an F_2 generation of one hundred fifty-one pigs, a backcross generation ($F_1\sigma \times \text{Duroc}\phi$) of thirty-five pigs, one F_3 litter of eleven pigs, and a litter ($F_1\sigma \times F_2\phi$) of five pigs. The F_1 pigs were all red with black spots, but the red became lighter in color as they grew older. The colors

of the F_2 and backcross pigs are given in Table II.

TABLE II

Color	F_2	Backcross
Black-and-red.....	47	19
Black-red-and-sandy....	10	
Black-red-and-white....	10	1
Black-and-sandy.....	11	
Black-sandy-and-white...	6	
Black-and-white.....	27	
Self Red.....	14	14
Red-and-sandy.....	3	1
Red-and-white.....	3	
Self Sandy.....	7	
Sandy-and-white.....	7	
White.....	3	
Totals.....	148	35

The red and sandy are quite distinct at birth but as the pigs grow older they may become darker or lighter until at maturity almost all shades of red are represented. Taking the classification at birth, it appears that black con-

A RED BOAR OF THE SECOND GENERATION OF BERKSHIRE X DUROC-JERSEY CROSS

This boar was lighter colored on the belly than on its sides. The presence of two shades of red on the same animal was characteristic of nearly one-fourth of the second generation. Compare with the nearly solid black pig shown on the opposite page. (Fig. 16.)

trasted with the absence of black gives a perfect 3 : 1 ratio in F_2 and a close approximation to a 1 : 1 ratio in the backcross. Therefore it seems proven beyond doubt that the Berkshire possesses a single factor for black spots which is absent from the Duroc-Jersey and the Tamworth. The size and

TABLE III

Description	F_2		Backcross	
	Calculated	Observed	Calculated	Observed
Black-and-red.....	62.4	67	17.5	20
Black-red-and-sandy.....				
Black-red-and-white.....				
Black-and-sandy.....	20.8	17		
Black-sandy-and-white.....				
Black-and-white.....	27.7	27		
Red.....	20.8	20	17.5	15
Red-and-sandy.....				
Red-and-white.....				
Sandy.....	13.9	14		
Sandy-and-white.....				
White.....	2.3	3		
Total.....	148	148	35	35

A LITTER OF THE SECOND GENERATION SHOWING VARIATION IN SIZE

The second generation was distinctly more variable than the first. (See the graphs in Fig. 19.) This would be expected if there is segregation of growth factors. The weights of these hogs, from left to right, were 221, 293, 90, and 170 pounds (Fig. 17.)

number of the spots is largely determined by modifying factors which must be numerous since no pig in F_2 possessed as much black as a pure bred Berkshire, although one had about ninety-five per cent as much. Those which have red color upon them as contrasted with those which have no darker shade of red than sandy, constitute almost exactly nine-sixteenths of F_2 and fully 100% of the backcross. Therefore it is reasonably clear that the Duroc-Jersey possesses two factors, absent in the Berkshire, which are both necessary for the production of red. If we assume that either factor by itself produces some shade of sandy, everything is satisfactorily explained except that the pigs carrying black have a much higher proportion of those which lack both red and sandy than do the pigs which do not possess the black. If we assume a factor linked to black which can dilute to white the sandy produced by one of the factors for red when the other factor for red is absent, the actual results are very close to the theoretical ones. This is shown in Table III.

The author has no theory to account for the presence of two shades of red on the same animal. The fact that these "bicolor reds" constituted almost exactly one-fourth of the F_2 generation seems to point to a simple explanation, but there were three easily distinguishable types of this bicolor character, as follows: first, roaning, of which the genetic foundation is not known; second, small irregular spots of red, usually around the face, particularly on the eyelids, while the rest of the animal was some lighter color; and third, a lighter colored belly than sides or back. This last feature is always found as an integral part of the agouti pattern where that occurs among other mammals and seems to be connected with that pattern here also, for it is very prominent on the young of the wild hog, and when an F_2 boar which was red with a sandy belly was mated to an F_2 sow (both from the Berkshire $\times$ Duroc-Jersey cross), which was white but had

A LITTER OF THE THIRD GENERATION (BERKSHIRE X DUROC-JERSEY CROSS)

Showing striped pigs with lighter colored bellies. This striping is a curious atavistic recurrence of an ancestral trait present in the wild boar but not in the domestic breeds. The recurrence here is doubtless due to a recombination of Mendelian factors. (Fig. 18.)

black pigment patches in her skin, there resulted an F_2 litter of eleven pigs *every one of which* had a lighter belly than back and was striped in a fashion similar to that of the young of the wild boar but differing from the latter in the shade of the contrasting stripes. They were not uniform in shade themselves. This type of "reversion" in swine has been reported frequently, but the remarkable fact that so large a litter as this should so uniformly show both the lighter belly and the striping pattern makes it seem certain that the factorial basis for this "reversion" is simple, and that both lighter belly and striping are produced by almost identical factor complexes. Two individuals of the F_2 generation showed stripes similar in pattern but very faint.

DIFFERENCES IN GROWTH

The economic importance of some definite knowledge in regard to factors for rapid growth may be surmised from the extent of the practice of cross-breeding hogs in order to secure the

greater size and vigor of the F_1 hogs for market purposes. The Berkshire $\times$ Duroc-Jersey cross was extensive enough to furnish significant data upon this point. Since they were not all born at the same season nor in the same year it is obvious that they were not all exposed to identical weather conditions nor fed rations as identical as would be desirable in a nutrition experiment. However it was the aim always to feed the best possible ration to produce the maximum practical gains and it is believed that the data are, on the whole, reasonably comparable. The pigs were weighed at birth and at monthly intervals afterward and the coefficients of variation of the weights of the F_1 individuals for each month were figured separately from those of the F_2 individuals. The resulting curves are shown in the accompanying figure.

The outstanding fact of genetic importance is that the variability of the F_2 generation is distinctly greater than that of the F_1 . That, to be sure, is

The Coefficients of Variability for the Monthly Weights of the F₁ and F₂ Generations of the Berkshire X Duroc-Jersey Cross

what was to be expected from the genetic standpoint and it is in agreement with the experience of practical breeders. It is definite proof of two things: First, that some of the factors which have the power to stimulate growth are not identical in the Duroc-Jersey and the Berkshire breeds; second, that there is some degree of homozygosis for these growth-stimulating factors within the limits of each breed.

To what extent being purebred from the standpoint of the breed registry society indicates homozygosis for these factors; whether the number of factors involved can be determined; whether the differences in growth indicate differences in the identity of these factors or merely differences in the combinations; and whether these factors are so linked in groups as to make recombination of the desirable ones impossible or at least impracticable; these

are all interesting questions which these data raise but are too meager to answer.

Other interesting questions, not primarily genetic, relate to the shape of the curve. The maximum variability seems to be reached at or shortly following weaning time and coincides with the period of the lowest mortality. Does this mean that the best time to select breeding stock for size and early maturity is when they are from three to six months old? Of course it was impractical to keep the hogs which were not desired for breeding purposes until they were two or three years old and therefore the question is still left open as to whether the hogs finally would not have become more uniform in size; that is, whether the differences were in potential size or in earliness of maturity.

Figure 17 is a photograph of one of the F_2 litters most extreme in its range of variation in growthiness. These four individuals had received identical treatment and none of them had ever been noticed to be sick, yet when this picture was taken when they were being marketed at the age of eleven months the weights were, from left to right: 221, 293, 90, and 170 lbs.! Possibly more animals are born runts than is commonly believed.

CONCLUSION

It is not believed that all the hypotheses which have been tentatively advanced in this article will stand in every detail without amendment the test of further research. However some of them will and if the others awaken enough interest to stimulate

further research with this economically important animal, they will have served a useful purpose.

POSTSCRIPT

Since the above was written a report⁷ of the results of a cross between mule-foot and Duroc-Jersey swine has been published which agrees closely with two of the main conclusions reached in this experiment: Namely, that the presence of black as contrasted to its absence is dependent upon a single factor and that the different shades of red are due to the interaction of a few independent factors which do not affect black. The black of the mule-foot swine behaves like that of the American Hampshire but whether the two are genetically identical is still undetermined.

Castle's Book Revised

GENETICS AND EUGENICS: A Text-book for Students of Biology and a Reference Book for Animal and Plant Breeders, by W. E. Castle, Professor of Zoology in Harvard University and Research Associate of the Carnegie Institution of Washington. Harvard University Press 1920. 395 pages, 154 figures.

Its clearness and pleasing style as well as its thoroughness of treatment have made Professor Castle's "Genetics and Eugenics" one of the most extensively used books on its subject since its appearance in 1916. In the second edition, which has recently appeared, several chapters have been rewritten and the total length has been increased by 42 pages in order to take account of the rapid progress of our knowledge. The subjects of linkage and the nature of genetic change are dealt with at considerably greater length.

The greatest change in viewpoint is with respect to the hereditary basis of fluctuating variation. In his earlier work, Professor Castle assumed that adult characteristics depend on rela-

tively few mendelizing unit factors, and accounted for his success in bringing about great changes, by means of long continued selection of minute variations, by holding that the hereditary factors are themselves subject to fluctuating variation. His own experiments, however, as well as others, stimulated to a large extent by his championship of the above view, have now demonstrated beyond question the remarkably high degree of constancy of most Mendelian factors, and have made it clear that each characteristic is in general affected by many factors. Fluctuating variation is thus interpreted in the present edition as depending largely on the recombinations possible in a system of many unit factors, each insignificant in its own effects, and to a much less extent upon mutation of the units themselves.

Taken as a whole, the book can be heartily recommended as giving a reliable and very readable account of the present state of knowledge of genetics together with a sane discussion of the application to eugenics.—S. W.

⁷ Detlefsen, J. A. and Carmichael, W. J. Inheritance of Syndactylism, Black, and Dilution in Swine. *Journal of Agricultural Research*, Vol. 20, pp. 595-604.