adhering by soft adhesions, which were easily broken down, to the lining membrane of the neck of the sac, which was continuous with the peritoneum. The rest of the appendix and the cæcum were close to the opening of the sac in the abdominal cavity. At the point of junction of the appendix to the cæcum was a ring of discoloration, and here the tube was compressed and slightly ulcerated. The appendix was thickened and covered by flakes of lymph, but otherwise appeared tolerably healthy. The peritoneum in the neighbourhood of the sac was dark-coloured, showing a little soft lymph on its surface. There was no general peritonitis. The lungs were very emphysematous and congested. There was puriform fluid in the bronchi. The heart weighed ten ounces; it was uncontracted. There was a little partially decolorised clot in the right ventricle. The muscular tissue was very soft and friable, and had undergone considerable fatty change. The valves were natural. The liver weighed $3\frac{1}{2}$ lb.; it was soft and pale, and infiltrated with fat. The other organs were natural.

Mr. Pick remarked in connexion with the above case that it was one of great rarity. He had never certainly seen a similar case. Nor did he even remember to have read of one where the vermiform appendix was alone strangulated. Of course cases where a portion of the bowel, probably cæcum, and the vermiform appendix together were found in a hernial sac were not uncommon. He had looked through the tables of operations for strangulated hernia in St. George's Hospital for the last twenty-eight years, and had not been able to find an instance of the same thing mentioned in these records, nor had he found the subject alluded to in any of the authors he had consulted. It seemed difficult to account for the way in which the vermiform process had become strangulated. How a tube of such small dimensions as the appendix should have been so tightly constricted in a canal which on previous occasions had allowed the passage of a much larger body, as a portion of the gut, or, at all events, omentum. For it seemed perfectly clear to his mind that the hernial sac must on a former occasion have contained either gut or omentum, for he could scarcely conceive how a soft, freely movable tube like the appendix could produce a hernia by forcing before it the peritoneum along the inguinal canal, though it was easy to understand how it might find its way into a hernial sac when it was once formed. He could come to no other conclusion than this, that the sac had been originally formed by the protrusion of a portion of intestine or omentum in the usual way, and into the sac thus formed the appendix has descended. It seemed strange under these circumstances that it should have become strangulated, and at the operation he had expected, when he saw what was in the sac, to find a small knuckle of intestine high up by the side of the appendix, but that this was not the case was abundantly proved by the post-mortem examination, which showed that the whole of the intestine was natural, except this little process, and that this had been tightly constricted was proved by the ring of ulceration which surrounded it, at the point of constriction, close to its junction with the cæcum. He believed that the strangulation must be explained by supposing that some inflammatory exudation had taken place in the tissues external to the neck of the sac, and that this was the cause of the stricture. From whatever cause it had arisen it was very tight, for it was with the greatest difficulty that the extremity of the hernia knife could be got beneath it to divide it.

Another point of interest was, he said, the acute symptoms which were produced. One would scarcely have imagined that constriction of the appendix would have produced such urgent symptoms, and such early vomiting, for it must be borne in mind that this was a very different thing to strangulation of a portion of the intestine, in which the perfect condition of the tube was essential to life. With regard to the appendix cæci it was *not* so; this tube might become obliterated without any absolute detriment to the patient, and Mr. Pick supposed that if the operation could be performed with safety it might be removed without in any way interfering with the natural functions of the individual. The symptoms of strangulation were due, no doubt, to arrested peristaltic action; but it was difficult to understand why this arrested peristaltic action should have taken place so suddenly from constriction of a small tube like the vermiform appendix, which was not essential to life. Mr. Pick further stated that at the time of the operation he was in considerable doubt as to how he should deal with the appendix: whether he should return it into the cavity of the abdomen; or whether, having freed the

constriction, he should leave it in the sac. On the one hand, it seemed a hazardous proceeding to return it, acutely inflamed as it was, and covered over by shaggy lymph, into the abdominal cavity; and on the other hand, if it were allowed to remain, it would undoubtedly contract adhesions to the tissues with which it was in contact, and would therefore always remain as an irreducible hernia, and might possibly become again strangulated by the contraction of the tissue during cicatrisation. Under these circumstances he decided upon pushing it back just within the cavity of the abdomen, and leaving it there in the expectation that it would contract adhesions to the peritoneum in the neighbourhood of the abdominal ring. And that this did occur was shown by the post-mortem examination, when it was found that it was already adherent in this situation.

ROYAL HOSPITAL FOR DISEASES OF THE CHEST.

CASE OF PHTHISIS, PROBABLY OF CONTAGIOUS ORIGIN.

(Under the case of Dr. MURRELL.)

THE chief point of interest in the case was the history, the symptoms having appeared immediately after nursing her husband, who died of phthisis.

Lizzie E—, aged twenty-six, was admitted on Sept. 1st, 1879, with phthisis pulmonalis. Her family history was good. Her father, a captain in the navy, was always a very strong man, and died of fever abroad. He was not subject to cough, and never, to the best of her knowledge, suffered from any chest complaint. Her mother was living, aged sixty-four years, and had been "asthmatical" for the last twenty years. She also had a cough, more or less, all the year round, and was very short of breath if she went out in the fog. The patient had three sisters, all well and healthy. There was no apparent trace of consumption at all in patient's own family.

She was quite well in every way up to the time of her marriage. Her husband, as far as she knew, was a healthy man when she married him, but he came of a consumptive stock. His father died of bronchitis, and his mother from phthisis. One of his brothers was, at this time, dying from consumption, and another was far gone with it. The patient was married in April, 1878, and in the second week of the following July her husband broke a bloodvessel and lost a great deal of blood. After that he had a cough, grew weaker and weaker, and could hardly do anything. In August he got too weak to go out, and from that time till the day of his death in February, 1879, he was assiduously nursed by his wife. He expectorated a great deal, a spitting-cup was of no use to him, and he had to have a hand-basin with water to spit into. Patient hardly ever left him, and slept with him. At night he perspired very much, but not enough to wet patient's things, or make her cold. She was in and out of bed pretty well all the night, through the long cold winter of 1878-9, and he refused to have a fire in the room.

After her husband's death the patient was worried and pressed for money, but did not suffer in any way from want of food or comforts. She went to live with her mother, and was as well off there as she would have been at home, or, perhaps, better. She had no cough up to the day of her husband's death; she was then completely knocked up, was seized with cold shivers, which continued for three days. She had a bad cough, and that gradually got worse. Nine months after she had coarse crepitation at both apices, was losing flesh, and sweating profusely at night—was in fact, consumptive.

It is certainly a question worth considering whether in cases of advanced consumption with copious expectoration there is not some danger of infection, however slight.

BIRMINGHAM HOSPITAL FOR WOMEN

LARGE INTRA-MURAL MYOMA REMOVED BY ENUCLEATION.

(Under the care of Mr. LAWSON TAIT.)

A. H—, aged twenty-eight, was admitted into the hospital on May 15th, 1879, suffering from profuse menorrhagia. The fundus was found to be retroflected to an extreme degree, and of very large size. The diagnosis of myoma was made, and an attempt to dilate the cervix by Mr. Tait's plugs was made, but it was unsuccessful owing to the extreme retro-

flexion. The cervix was therefore completely divided by the hysterotome, the thick capsule of the tumour opened, and the tumour separated as far as the fingers could reach. The patient was so extremely anæmic that everything done to her was done in the constant fear of the production of heart-clot, and even the administration of ether in her case seemed risky. Nearly eight weeks elapsed after this preliminary proceeding before it was deemed safe to do anything more, when it was found that the opening into the capsule had closed, though the division of the cervix had remained permanent. On July 26th the dilating plugs were applied, and next day the cervix was so open that a large free incision over the tumour was made, and the tumour freely separated. On August 4th the tumour was found presenting through the os, and a large piece of it was removed by the écraseur. On August 12th the rest of it was found extruded and hanging by a thick pedicle, which was divided, and the mass removed. For many weeks after this operation she hung between life and death, and made a convalescence protracted over nearly eight months.

Mr. Tait is of opinion that in such a case removal of the ovaries is a much safer operation than enucleation of the tumour, and in his hands it has given much better results. Out of five cases where he has performed vaginal enucleation in the last three years there have been only two recoveries, whilst of seven similar cases in which he has performed oophorectomy there has been only one death, and that death was due solely to the operation having been too long delayed.

SOUTH STAFFORDSHIRE GENERAL HOSPITAL.

POPLITEAL ANEURISM; TREATMENT BY FLEXION; CURE.
(Under the care of Mr. NEWNHAM.)

GEORGE L—, aged forty-four, a goods guard, was admitted with popliteal aneurism. His previous history was excellent: he had never had syphilis, and had always been abstemious. Several months before he noticed the aneurism he wrenched his knee, and had to lay up for a short time. He could remember no other injury to the leg, nor could he recollect having made a forcible or sudden muscular effort to lift weights.

The history of the aneurism was as follows: Three weeks before admission (since Jan. 10th), the patient noticed a pulsation in the left popliteal space. His attention was directed to the seat of disease by pain in the calf when walking, the pain passing off if he stood still even for a moment. He thought that the swelling when he first noticed it was nearly as large as on admission.

On January 31st he was taken in under the care of Mr. Newnham. In the left popliteal space there was a tumour, about the size of a hen's egg. It had an extensile pulsation which could be arrested by pressure on the femoral artery. The area of pulsation extended over the popliteal space. There was a distinct bruit, and a thrill was communicated to the hand. On examining the other arteries of the body the left radial and temporal arteries were found feebler than the corresponding arteries on the opposite side, but no intra-thoracic tumour could be detected. The superficial arteries had not an atheromatous feel, the heart was healthy, and no visceral disease could be discovered. The patient had a cataract of the right eye, which did not result from injury. He was a corpulent man, and somewhat aged in appearance, but his health, he said, was exceedingly good.

The patient was put on meat diet, and given five grains of the iodide of potassium and fifteen of the bicarbonate of potash. He was kept in bed and had an occasional purge. On Feb. 4th, he was directed to keep his thigh flexed on the abdomen, and the leg on the thigh; the iodide was increased to ten grains. On the 6th, he was put on a modified Bellingham's diet. Breakfast: bread and butter, two ounces; milk, two ounces. Dinner: meat, two ounces; bread, two ounces; beer, two ounces. Supper: bread and butter, two ounces; milk, two ounces. On the 8th, the extensile pulsation had ceased in the tumour, although the pulsation in the popliteal and posterior tibial arteries could be felt. On the 9th, the angle of flexion was increased; daily allowance of fluid fourteen ounces. On the 11th, flexion was discontinued; patient put on meat diet, with one pint of fluid. The aneurismal sac was felt to the inner side of the popliteal artery; it was a little larger than

a walnut, and felt solid to the touch. The patient was left in bed for ten days after pulsation had ceased, and was discharged on the 9th of March, when the tumour could be scarcely felt.

Medical Societies.

ROYAL MEDICAL & CHIRURGICAL SOCIETY.

Extra-uterine Gestation treated by Abdominal Section.—The Treatment of Pelvic Suppuration by Abdominal Section and Drainage. — Sub-coracoid Dislocation of the Humerus.

THE ordinary meeting of this Society was held on the 11th inst., J. E. Erichsen, Esq., F.R.S., President, in the chair. The subject of abdominal section was under discussion, introduced again by Mr. Lawson Tait, who, it will be remembered, contributed papers dealing with abdominal surgery earlier in the session. This time he related a third case of extra-uterine gestation, and in his second paper advocated the rather bold treatment of pelvic abscess by abdominal section. Both papers gave rise to some discussion. A paper by Mr. Eve, describing a rare variety of dislocation of the humerus, was also read.

A paper was read on "A Third Case of Extra-uterine Gestation, treated by Abdominal Section," by Mr. LAWSON TAIT. The case differed from the two previously read before the Society in that the operation was performed before the death of the child, whose life was saved, but the mother succumbed to the operation. Mrs. S—, aged thirty-three, had been married fourteen years, and had had six children. Since her last menstruation in May, 1879, she had been in constant pain, of which a severe attack occurred in July. In October she became satisfied she was pregnant, and Mr. Lattey, suspecting the nature of the case, asked Mr. Tait to see her. A large abdominal swelling was evidently a pregnancy, and to the left of the umbilicus was a prominent mass, which, from its rhythmic contraction, was clearly the uterus, and was attached to the larger mass behind it. The head of the child was detected between the rectum and the vagina. The pain becoming agonising, and being unrelieved by opiates, an operation was determined on. Ether was administered on the morning of Jan. 31st, and on making the median incision the empty uterus was found, and the cyst, which was the right broad ligament, was opened, and the child removed without difficulty, the edges in the opening of the cyst being then secured to those of the parietal wound, leaving an opening for the discharge of the placenta, and the peritoneal cavity was closed. The child was at first profoundly narcotised, but is now, three weeks after, thriving and well. The death of the mother, which occurred on the fourth day, seemed to be from protracted shock. At the post mortem there were found some morbid appearances in the liver and ante-mortem clots in the heart, but no peritonitis; the cyst proved to be the broad ligament as diagnosed, and there was a deciduous lining in the uterus in process of separation. The Fallopian tube had been ruptured at its lower aspects, the ovum thus escaping into the tissue of the broad ligament; and the case confirmed the author's opinion that all extra-uterine pregnancies are originally tubal, that the tube bursts in between eight and twelve weeks, and when not fatal an extra-uterine pregnancy is completed. The unfavourable result to the mother was due to the delay in the operation; but the case is of interest from the saving of the child. The child might have been removed by vaginal section, but large venous sinuses in every direction made it likely that any injury to one of them would be fatal, and in vaginal section it is impossible to see what to avoid.—Mr. DORAN had recently dissected three cases of advanced tubal pregnancy, and had come to the conclusion that operating from the vagina was unjustifiable. The cyst in each of the three cases had pushed down the upper part of the rectum, which had become adherent to it. An incision of any length sufficient for the extraction of a fœtus could not have failed to wound the gut (and possibly a loop of small intestine as well pushed under the cyst); this would entail the most disastrous results.