phosphorised diet than Ireland. Where England consumes cereals, a phosphorised diet, Ireland consumes the potato, a non-phosphorised. Hence the sum total of the causation would stand as the product of a predisposing cause and a predisposed nervous system, in both of which elements of causation—if they are what I have here propounded—Ireland has the advantage of England. And I confess I am not able to discover, either by statistics or reflection, any other elements of causation in which the two countries differ.

CLINICAL NOTES AND CASES.

The Treatment of Hæmatoma Auris. By GEORGE J. HEARDER, M.D., Medical Superintendent of Joint Counties Asylum, Carmarthen.

The “Insane Ear” has recently had a large share of attention accorded to it by various writers, who at least agree in this that they abstain from recommending any procedure whereby we may hope to check the disease, or prevent the withering and shrivelling which generally remain as a permanent local record of grave cerebral disturbance.

Five of the cases here recorded occurred in males; one, in a female who had both ears affected. In each case the tumour involved the left ear; and the right auricle became diseased in the woman after the left othæmatoma had terminated. The form of mental disease was:—

In 3 cases—Melancholia.

In 1 case—Mania chronic.

In 1 case—General paralysis.

In 1 case—Dementia.

In two instances the development of the tumour was probably assisted, or perhaps caused, by local injury.

1.—S. D., female, aged 33. Congenitally weak. For 12 months restless, violent, melancholic and suicidal. On the 27th August, 1874, a hæmatoma auris of the left side was observed; it increased steadily, became very tense and painful, and was evacuated by free incision eleven days after it was first noticed. Result—ear shrivelled, thickened, and its normal markings obliterated. On the 1st October, a hæmatoma on the right side was progressing rapidly, the effusion filling up the concha and fossa of antihelix. The inner surface of the pinna was painted over with a few drops of the acetum cantharidis.

The swelling was immediately checked, and by the 6th was subsiding markedly. Result—the auricle perfectly preserves its form, and appears unaltered from its healthy state; only by touch it is evident that the concha is permanently thickened to a small extent.

2.—D. J., male, aged 53. Mania chronic. Always merry and mischievous. On the 3rd October, 1874, an othæmatoma of the left side was noticed. The parts affected were the concha, the antihelix, and its fossa; the tumour was half an inch thick, and steadily growing. The blistering fluid was applied to the inner surface of the pinna. Result—the markings of the ear are well preserved without any shrinking; but with thickening and increased rigidity of the parts involved.

3.—D. S., male, aged 25. General paralytic, very restless and excitable. On the 10th October, 1874, an othæmatoma of the left side filled the concha and the fossæ of the helix and antihelix, the ear being quite half an inch in thickness. Acetum cantharidis was applied to its inner surface. Result—within three weeks, absorption of the fluid had taken place; the outline of ear and its markings were preserved; there was slight permanent thickening, noticeable only to the touch, and perfect opacity when placed between the eye and the light. The propriety of blistering this case was discussed and doubted, owing to the tendency to sloughing in general paralysis, and the local gangrenes the patient had already developed. The blister, however, acted effectually on the cuticle, without producing any unfavourable complication. The patient died on the 28th November, 1874.

4.—J. P., male, aged 56. Melancholic, acutely depressed, and incessantly and loudly bemoaning his lost condition; suffering also from cardiac and arterial disease. On the 6th November, 1874, he stated that he stumbled and fell, while getting out of bed during the night, and bruised his left ear. Two days later the ear was still discolored, and there was then evidently a layer of fluid within it, and a hæmatoma forming. The inner surface of the ear was blistered, as in the previous cases. Result—recovery, excepting slight thickening still observable on examination. The auricle preserves its size and shape.

5.—T. J., male, aged 25. Demented and scrofulous. On the 5th March, 1875, a left hæmatoma was forming, affecting the antihelix, its fossa, and helix. The swelling was about three-eighths of an inch in thickness. The inner surface of the pinna was blistered. On the 19th the fluid was not absorbed, though the morbid action was checked; and the blister was repeated. Result—size and markings of the ear well preserved; the parts which had been affected remaining slightly thickened. The hæmatoma in this case might have had a traumatic origin, as the patient was in the habit of beating his face with his fists.

6.—D. L., male, aged 54. Melancholic, quiet, intensely pre-

occupied; has abstained from food; suffers from cardiac disease. On the 1st February, 1876, a hæmatoma was noticed, affecting the upper portion of the left auricle. On the following day the tumour was much larger, involving the entire upper half of the pinna. The blistering fluid was applied to the inner surface of the diseased part. All morbid action was at once checked, and by the 16th the effused fluid had been absorbed, and, save for a slight thickening of the upper portion of the helix, no malformation remained. This patient has been able to leave his bed for only a few hours daily for several months past; he reposes invariably on his right side; he is under most careful supervision, and there is no suspicion of the ear having been in any way injured.

We have had records of six consecutive cases of othæmatoma similarly treated. In each the diseased action was arrested on the application of a blistering fluid to the inner surface of the pinna. In all, we can trace the results of the morbid agency in permanent thickening of those parts which had been subject to it; but there is none of the shrivelling and obtrusive distortion which ensues when the affection has run its course uncontrolled, as it did in the left ear of the first case.

Cases with Mental Symptoms following the Administration of Chloral Hydrate. By G. HUNTER MACKENZIE, M.B. Edin., Resident Medical Officer, District Infirmary, Ashton-under-Lyne.

CASE I.—Mrs. A., æt. 55, married. Admitted into the Infirmary 14th April, 1875, suffering from aortic valvular disease, with sequelæ. Ordered iron, digitalis, and gin, generous diet, and an unstinted supply of fresh milk. On account of the sleeplessness and restlessness, on the 30th April, there was prescribed for her 15 grs. of chloral hydrate, to be taken at bedtime. A week afterwards this was doubled. About the middle of May she was observed to be getting decidedly "strange;" the mental aberration gradually increased, and towards the end of the month culminated in subacute mania, with hallucinations and illusions of sight. No improvement took place in the physical symptoms.

CASE II.—Mrs. B., married, æt. 25. Admitted 7th July, 1875, suffering from mitral valvular disease, chronic articular and muscular rheumatism, and sciatica. Treated by alkalies and a non-nitrogenous diet. Opium by the stomach and mouth being badly borne, on the evening of the 20th July she was ordered 20 grs. of chloral hydrate. This, she stated, made her feel "drunk." Two nights subsequently the dose was increased to 30 grs. with the following results:—The medicine was given at 9 p.m., and at 10.15 p.m. she was seen by re-