THE
BOSTON MEDICAL AND SURGICAL JOURNAL.

VOL. LXXXVIII.]

THURSDAY, JUNE 19, 1873.

[No. 25.

Original Communications.

THE MODERN OPERATION FOR CATARACT, AS MODIFIED
BY M. DE WECKER. WITH A PARTIAL TRANSLATION
FROM HIS ANNUAL HOSPITAL REPORT.

By HASKET DERBY, M.D., Boston.

SINCE the flap extraction of cataract has been so largely superseded by the peripheric linear method of von Graefe, Dr. Liebreich, of London, has brought prominently forward another operation, which now generally bears his name. He makes a cut throughout in the cornea, curved, about midway between its horizontal meridian and its upper or lower edge, and, having ruptured the capsule, extracts the lens without removing any portion of the iris.

The operation of von Graefe consists, as is well known, in a nearly linear incision at the upper sclero-corneal junction, in the excision of a piece of the iris, and the subsequent extraction of the lens. Dr. de Wecker makes a larger and more curved incision, which, "instead of being linear, forms a small flap two millimètres high." This done he follows, step for step, the method of von Graefe; and may be said to do his operation with a comparatively slight variation as well as equal results. These last words are justified by the following quotation from de Wecker's annual report.

"M. de Wecker's section does not afford, as regards the number of cures obtained, any larger figures than those obtained by the method of von Graefe. In spite of its curve, the section made use of by our master (it is the assistant of de Wecker who writes) allows a coaptation of the lips of the wound as close as that observed in the method of the celebrated ophthalmologist of Berlin, and, like the latter, results, in a great majority of the cases, in a union by first intention and an immediate and total cure, as may be seen by a comparison of the statistics which represent the complete success of the two operators."

The methods of von Graefe and de Wecker belong to the same family and may, for all practical purposes, be considered as the same. The operation of Liebreich is entirely distinct. The lines between these opposing systems are fairly drawn. On the one hand, we have a cataract made to emerge through an artificial pupil and a peripheric linear or curved incision; on the other, an extraction through the

VOL. LXXXVIII. No. 25

natural contracted pupil and a median corneal cut. The relative advantages of these two methods are now engrossing the attention of ophthalmic surgeons. On the side of Liebreich, most plausible assertions have been made. On that of von Graefe and de Wecker, numerous carefully prepared statistical tables have been compiled, showing a larger percentage of success than has hitherto been known to be attained. Until answering and more favorable figures have been brought forward by the upholders of the new operation, its value must, of course, be considered as undecided.

Meanwhile I desire to present, for the benefit of those interested in such matters, a portion of the excellent annual report* of de Wecker's clinique in Paris, giving his reasons for modifying the peripheral linear extraction, as well as some practical and novel hints regarding preparation for the operation and after treatment.

(Translation.)

The best means to employ in order not to provoke a prolonged irritation of the portion of the ciliary body adjoining the wound, is to keep farther away from it than von Graefe did. Prof. Horner, a particularly experienced and prudent operator, wrote recently to Dr. de Wecker:

"Nothing appears to me to more surely avoid the dangers of healing-in of the iris (irido-cyclitis and even sympathetic iritis, which I have been unfortunate enough to observe twice) *than a less peripheral incision* and the non-use of atropine before the operation; I have no doubt but that your experience has been the same."

Our first care, then, should be to avoid all healing-in of the extremities of the sphincter of the iris at the corner of the two commissures. The iridectomy once performed, every portion of this membrane ought to occupy the same position that it did before the creation of the artificial pupil.

Moreover, numerous observations have proved to us the advantage, alluded to by the Zurich professor, of not dilating the sphincter of the iris before proceeding to extraction. If the instillation of atropine be avoided, it can be seen that this membrane returns much more readily into position after the excision of one of its parts. M. de Wecker, accordingly, makes a more sparing use of this mydriatic. During the year 1871, he began to give it up after the operation in the majority of cases, in order to avoid the irritation of the conjunctiva that comes on in a certain number of individuals.

For nearly a year, all extractions at the clinique have been performed without the previous production of mydriasis. In fact, as has been noted in the foregoing tables, we only use this agent when the aqueous humor is somewhat cloudy, when it contains masses of corticalis, and when the iris threatens to inflame.

* Ophthalmological Clinique of Dr. de Wecker at Paris. Statistics of the operations performed during the year 1872. By Dr. Georges Martin, Chef de Clinique. Paris, Adrien Delahaye.

A combined extraction* is often accused of being a complicated method, and involving for the patient an artificial pupil. But in reality such an operation makes no unreasonable demand on the skill of the surgeon. It is unjust to attribute to an operative method those drawbacks which are not inherent in it, but are solely due to a want of accuracy in the performance.

If a coloboma of the iris interferes with vision, it is in a far smaller proportion of cases than is generally supposed; the majority of patients easily learn, at the end of a certain time, to disregard it by slightly drooping the upper lid.

And, as the escape from the slight inconveniences entailed by a solution of continuity of the fibres of the iris is purchased at the price of several suppurations, no operator should hesitate to choose that method which the interest of the patient demands. It is in establishments like those ruled by M. de Wecker, where hundreds of extractions occur in a year, that the choice of that method that leads most certainly to a cure is most imperatively demanded.

In a simple extraction, the larger portion of the cataractous mass can emerge through the natural pupillary opening, but it is almost impossible for all the cortical layers imprisoned behind the iris to follow the same road. Moreover, setting aside clinical experience, anatomy and mechanics alike demonstrate that a *complete* delivery of the crystalline is only possible by a combined method, unless the cataract be extracted in its capsule, which would be an unhappy reversion to a process considered dangerous, under many circumstances, by nearly all the operators who have employed it.

The present tables give seven entire failures, figures that cause just pain to an operator, even when met with among 217 extractions. How great, however, would have been nearly every week the burden of care if the twenty-two suppurations, that would have resulted from an equal number of extractions by the old flap method, had to be undertaken. Does not, moreover, the surplus of fifteen successes attending the method of curved combined extraction more than compensate for the slight amount of dazzling experienced by a few stray patients?

Remark that the proportion of ten per cent. of failure does not attend flap extraction solely; the new methods without iridectomy, elaborated though they be, lead to an equally inferior result, as the most recent statistics of the method of Weber prove.

Despite all the above-mentioned precautions, the most dexterous operator will always happen on a certain number of patients in whom suppuration is inevitable.

This complication, in very many cases, is impossible to foresee, and cannot find its explanation in a morbid state of the fundus oculi. The most serious affections of the choroid and vitreous humor (pos-

* I. e., extraction prefaced by an iridectomy.

terior staphyloma, disseminate choroiditis, floating bodies in the vitreous, partial detachment of the retina) seem to have no unfavorable influence on immediate accidents, but rather on those which occur after a certain lapse of time.

It is safe to state that the majority of suppurations which occur the second or third day after the operation, are due to an irritation of the conjunctiva.

When the mucous membrane is inflamed, either primarily or in consequence of chronic lachrymation, of slight dacryocystitis, or of blepharitis, the pus corpuscles emigrate towards the lips of the sclerocorneal wound, and the adhesive process is interrupted and fails to induce a union by first intention. In another series of cases, the direct cause eludes observation; the organ is probably the seat of a congestive condition which, stimulated by operative interference, encourages the exit of white blood corpuscles; these latter, placed between the edges of the section, interfere with the developement of the phenomena of cicatrization.

Governed by this theory at the clinique, whenever there is the least reason for fearing a suppuration of the lips of the wound, we have recourse to a remedy which has been adjudged useful in preventing the passage of leucocytes, and incapable of interfering with the phenomena of proliferation. We speak of the chlorhydrate of quinine, recommended by Prof. Nagel several years ago to meet the inflammatory complications experienced by some patients.

If, at the time of the second dressing, which generally takes place twenty-four hours after the operation, the conjunctiva is observed to be the seat of a considerable secretion, the aqueous humor to have lost its transparency, the segment of the cornea adjoining the wound to be invaded by a grayish tint (differing from the opaline color which follows a simple contusion), and if, finally, the conjunctiva presents a slight tumefaction, we prescribe immediately a solution of chlorhydrate of quinine.* The wound is once an hour carefully bathed with this preparation, a certain quantity of which is allowed to trickle down, drop by drop, from a slight distance above it. After this application, a drop of atropine (about gr. i. to 3 iij.) is instilled between the lids, in contact with which is constantly kept a compress soaked with the solution of quinine. This treatment is only suspended at night, during about six hours. The patient's sleep is favored by the administration of hydrate of chloral in doses of from gr. xxx. to gr. xlv. During this time, unless the suppuration is excessive, the eye is protected by a slightly compressive bandage.

By means of the quinine and the atropine, an eye threatened with panophthalmitis can be saved one time in three, and in nearly every case suppuration can be limited to the cornea. A fact worthy of notice is that the salt of quinine induces a nearly immediate cessation of the acute pain experienced by the patient, and reduces it to a sim-

* R. Quininae chlorhydrat., gr. vi.;
Aqua destill., dr. xv. M.

ple sensation of warmth. We see, therefore, that, though the solution of chlorhydrate of quinine does not invariably arrest the disease, it offers the immense advantage of disembarassing in many cases the loss of the eye, of acute suffering, and of nearly complete atrophy (phthisis) of the operated organ.

As has been demonstrated, this system of medication and the operative process, on which we have so earnestly dwelt, grant an amount of security procured up to to-day by no other method of treatment. This operation allows the surgeon to quit his patient, divested of uneasiness as to the result of the extraction. Entire assurance may be felt when the following principles have been complied with:—

Easy and complete evacuation of the nucleus and corticalis.

Restoration to its position of the internal sphincter of the iris.

Perfect coaptation of the wound as well as of the conjunctival flap.

(End of translation.)

I will add a brief *résumé* of the 217 extractions included in de Wecker's report of this year.

Twenty-five cataracts were not fully ripe. Five patients had diabetes. Vitreous was lost in twelve cases, all of which recovered with a vision of at least $\frac{1}{4}$. Iritis occurred six times. Seven eyes were entirely lost. The percentage of success was thus 96.88. The year previous it was 96.52.

Vision amounted in 19 cases to $\frac{2}{3}$

" 58 " " $\frac{2}{3}$

" 49 " " $\frac{1}{2}$

" 69 " " $\frac{2}{3}$

" 17 " " $\frac{2}{3}$

" 10 " " $\frac{1}{4}$

" 10 " was unrecorded;

these patients having their examination deferred until they should return for the operation on the second eye.

"The figures obtained by testing the vision, joined to those representing the proportion of cure, offer in themselves convincing testimony as to the superiority of extraction by the small peripheric flap, combined with iridectomy, over every other method."*

Such cases give no uncertain sound. Their results challenge comparison, and are not to be met by vague and unsupported laudation of a new and different method. That the pure or modified operation of von Graefe is never to be improved upon, I do not for an instant maintain. But what is claimed to be an improvement upon it must rest, in this day and generation, upon a basis of solid facts, and be known by its fruits.

* Quotation from Report.