already referred to under the head of general disinfectants, is often considered to be devoid of antiseptic properties because of the readiness with which it oxidizes all organic matter. It is true that potassium permanganate is readily decomposed by the organic matter with which it first comes in contact, and that it would be unusual indeed to find bacteria the only form of organic matter present in any one case. When we remember, however, that a very large number of bacteria represent but an infinitesimally small amount of organic matter, we will appreciate why potassium permanganate is capable of destroying vast numbers of micro-organisms in those cases in which its action can be confined to them.

This substance finds its greatest usefulness as a local application in rendering foul ulcers odorless. For this purpose a fairly strong solution, 2 per cent., is freely used as a wash. Much weaker solutions are sometimes used with great benefit as urethral irrigations in gonorrhea and as vaginal douches. A hot 1 per cent. solution of potassium permanganate will be found extremely useful as a deodorizer for instruments after use in the vagina while awaiting sterilization. Potassium permanganate destroys some alkaloids much more quickly than others, e. g., morphin more readily than strychnin. It can not enter the circulation, hence when swallowed it can act only on the substances found in the stomach. Obviously it is not suited for hypodermic injections.

Clinical Reports

SARCOMA IN A YOUNG CHILD.

FRED S. SPEARMAN, M.D., C.M.

WHITING, IOWA.

A number of interesting points are to be noted in connection with the following case:

1. The extreme youth of the patient, so that the malignant growths or the causes favoring their production must have been present almost, if not quite, from the time of birth.

2. The occurrence of tumors in several generations of the patient's family, showing an apparent hereditary tendency to neoplasms.

3. The marked displacement of the heart and spleen from their normal locations as the result of pressure by the tumor in the lung.

Patient.—E. S., born Nov. 15, 1902, male.

Family History.—Parents are both healthy; their occupation farming. A brother and sister are also in good health. Father's father and grandmother died from carcinoma. Both died years before the birth of my patient. Father's mother had an epulis removed about six years ago. Another growth came on another portion of the gums a year or so later, but disappeared of itself.

Previous History.—The child was never very robust, but had had no serious illness before the present one.

Present Illness.—I saw the patient for the first time on July 10, 1905. The parents stated that a little more than a year previously they had noticed an enlargement in the right side of the boy's scrotum. This had grown steadily but slowly for the first eight or nine months, but since then more rapidly. It did not seem to be painful at any time, but lately, on account of its increasing bulk, had caused some inconvenience, and the child appeared slightly nauseated at times.

Examination.—This showed a tumor of an ovoid shape about half the size of one's fist in the right side of the scrotum. It was solid, smooth, not translucent, no fluctuation was present or nodules felt, and there was no tenderness on pressure. The skin over the tumor was normal, there was no redness or sign of ulceration. The left testicle could be felt in the upper part of the scrotum and was of normal size. The child was bright and intelligent. He was of rather slender build and his complexion was tanned. Evidences of rachitis were present, as shown by a rather large head, somewhat box-shaped, and a

slightly developed rosary, which became more pronounced as the case progressed. I advised operation.

Operation.—A trocar was introduced, but no fluid obtained. An incision about one and one-half inches long was made through the superficial tissues into the growth. This was of a reddish-gray color, homogeneous, was friable, of about the consistency of rather old bologna sausage, and was adherent to the tunica, which were considerably thickened. I got out the growth piecemeal, removing portions of the tunica also. A small quantity of clear fluid escaped when the upper part of the tumor was removed. Chloroform was the anesthetic used, and I noticed that the child did not breathe well toward the last, although the heart's action was fairly good throughout. Recovery was uneventful, and I let the parents do the dressing after the first week. There was a slight serous discharge for a time from the wound, but it had practically healed by July 26.

I sent the tumor to Dr. G. S. Browning of Sioux City for examination, who pronounced it a fibrosarcoma containing a large proportion of fibrous tissue.

After the operation the boy's general health improved for a time, his appetite gained and he played more with the other children.

Further History.—August 10 the parents brought him to me again. They said that he had seemed listless for a few days, caring neither to eat nor play and showing some nausea at times. His skin and conjunctivæ were slightly jaundiced. As the weather was very hot at the time, I attributed his condition to this and did not examine him closely, but gave him a calomel purge and a bitter stomachic mixture. August 13 he was again brought in, as he was no better. Jaundice was more marked, and the mother informed me that the nausea was increased and that he had complained at times of pain in the left side in the region of the lower ribs.

Examination.—Percussion showed the area of liver dullness extended nearly an inch below the costal margin. On the left side there was an area of dullness reaching from about the lower level of the fourth rib in the anterior axillary line to an inch below the border of the ribs. This dullness extended to the spine behind and nearly to the median line in front. On palpation a firm mass could be felt below the ribs. Auscultation revealed an absence of breath sounds over the lower area of pulmonary dullness and faintness in the upper portion. There was less expansion of the left side of the chest. The apex beat of the heart was displaced nearly to the left border of the ensiform cartilage. Considerable meteorism was present. Pulse was about 140 and temperature 99. I suggested the possibility of an internal growth similar in nature to that occurring in the testicle, and judged from the physical signs present that the liver and spleen were probably involved, as well as the lung. An examination of the urine made a few days afterward was negative. In view of the autopsy findings, later examinations, which I omitted to make, would no doubt have revealed renal involvement.

The patient was put to bed on liquid diet, and under fair-sized doses of calomel the jaundice disappeared to a great extent in a short time. At first the stools were bile-stained, but afterward became slate-colored, remaining so until the end. There was considerable trouble throughout on account of meteorism, requiring constant use of purgatives and enemata. The dullness extended on both sides quite rapidly, and the child's respiration became continually more embarrassed. An aspirating needle, introduced into the left chest on August 21, brought only a small amount of bloody fluid. The following day Dr. R. E. Conniff of Sioux City saw the case with me and agreed in the probable diagnosis of an internal sarcoma. A blood count made by him showed the white cells to number 35,000 to the c.mm.

At this time there was a marked bulging out of the ribs on the left side behind and laterally, although the intercostal spaces did not bulge. The dull area on both sides was nearly two inches below the costal margin and the liver was plainly palpable. The outlines, both of it and the tumor on the left side, could be seen through the abdominal walls. On deep pressure on the left side of the abdomen at the level of the umbilicus a hard body could be felt, which I thought to be a nodules in the large intestine. The apex beat was displaced to

within one inch of the right nipple and about one-half inch below. To and fro murmurs were present. They were harsh, prolonged, and of a somewhat ringing or humming quality, giving the impression of tenseness, like the sudden stretching of an elastic band. Pulse rate was about 160 and very weak, respirations over 40, no rise in temperature. There was slight puffiness of the eyelids, especially the right one, and the superficial veins on the right side of the neck and over the liver were distended. Priapism was present throughout the latter part of the case, being more marked after micturition. A condition of phimosis present may have had something to do with this, as well as the splenic involvement. There was no urinary suppression at any time, although the quantity was decreased. The child became continually more emaciated, his breathing more and more difficult, as toward the last air entered only the very apex of the left lung, accessory muscles of respiration were brought into play, the heart's action became faster and weaker, and on September 1 he died.

Autopsy.—This was performed the next day. An incision was made from the ensiform cartilage to the umbilicus through the abdominal wall, which was very thin. The diaphragm could be seen bulging downward on the left side and was very tense on account of pressure of the growth from above. On incision the tumor was found closely adherent to the diaphragm, which formed its base. It was of a spongy, stringy consistency and nearly all of the left lung was involved. The mass was softer in the center, being almost semi-fluid, and portions removed were of a grayish color, slightly tinged with pink. The spleen was not much enlarged and was shoved down to the level of the umbilicus. The liver extended down to the same level, was very friable, and of a lighter color than normal, and yellow patches and streaks were present throughout. A whitish fluid escaped from the cut or broken surfaces on squeezing. The kidneys were enlarged and engorged, being of about the same color as the spleen.

Specimens of the different organs were sent to Dr. Henry Albert, state pathologist, of Iowa City, for examination. He reported that the pulmonary growth belonged to the class of endothelial sarcomata, although the arrangement of the cells was such that many pathologists would call the growth a carcinoma. The liver showed subacute inflammation. The spleen had distinct areas of inflammatory changes, with infiltration of round cells, and the kidneys manifested a distinct acute diffuse nephritis. Neither liver, spleen nor kidneys revealed any evidences of malignancy, however.

PNEUMONIA COMPLICATED BY MENINGITIS.

W. H. AXTELL, M.D.

Surgeon Northern Pacific Railway.

BELLINGHAM, WASH.

I report the following because of its infrequent occurrence:

History.—Mrs. W., aged 32, family history negative, very sparely built, has one son 5 years of age, and has had no illness since childhood. The present illness began Nov. 28, 1904. In the morning she had a sharp chill, followed in the afternoon by high temperature and a slight pain in the right lower chest. During that night and the forenoon of the next day the pain became very much augmented.

I saw her at 3 p. m. on November 29. Her temperature then was 105 1/5; pulse, 140; respiration, 36 and panting in character; she was cyanotic. Auscultation revealed crepitant râles in the lower lobe of the right lung in the axillary line. The area of dullness on percussion was about the size of a silver dollar. The pleurisy extended over a greater part of the right lateral chest. There was slight delirium when partially asleep.

Diagnosis.—Lobar pneumonia, with pleurisy.

Course of Disease.—She was moved to the hospital the same evening. The course of the disease was tempestuous, but followed that of a typical pneumonia, the crisis occurring on the

eighth day. On the third day slight meningeal symptoms developed. These began with a very sharp frontal headache, the pain of which was very distracting. The pupils, at first contracted, were dilated after the fourth day. There was some rigidity of the muscles at the back of the neck, but no convulsive movements. The heart showed no signs of depression or of inflammation. The delirium, at first mild, increased in severity until, on the day of the crisis, she became a raving maniac.

Osler says: "Meningitis is usually associated with endocarditis and occurs in 8 per cent. of fatal cases and endocarditis in 25 per cent. of all cases."

This woman had no heart complications whatever. Two days before the crisis, when the temperature was the highest, the patient became so violent that it required the combined effort of two nurses, night and day, to restrain her. On the twenty-first day of the disease she was discharged from the hospital, thoroughly convalescent. Up to the present time her mental condition has remained normal and there has been no evidence of insanity. The treatment was simply supportive. The ice cap was applied for the relief of the head symptoms and cool spongings for the temperature.

POINTS OF INTEREST.

1. Complication of meningitis without endocarditis.
2. The violent nature of the disease itself. (The meningitis in no way retarded or altered the usual course of the disease or lengthened its duration.)
3. The heart during the whole course of the disease and without special treatment showed very little depression.
4. A study of the chart shows a marked similarity in the variation and fluctuation of the delirium, temperature, pulse and respiration.
5. Of the many authorities consulted, Osler is the only one who particularly mentions meningitis as a complication of pneumonia. Tyson in his recent work simply calls attention to Osler.
6. Complete recovery.