to map out the dangerous zones and the safe ones, and exercise a little gentle massage. At this period the education of the venous circulation can begin. The patient can be placed with her legs hanging outside the bed, and in this position they will of course get congested and edematous. A little gentle massage will be found of service afterward. A few days later she can be placed on an inclined plane and lifted up to an angle of 45 degrees, then perfectly upright. During the fourth week some passive movements of flexion of the thigh are carried out; the patient is allowed to make a few steps; the latter should be of the same length. No sticks nor crutches should be allowed, so as to prevent any bad habits being acquired. A certain amount of redness and edema may appear, but will vanish after rest in bed. As for trying to sit up, it is found useful to use a number of cushions to cause variations in the angle allowed. When this has once been carried out, the patient can be told to carry out the treatment by herself. Dr. Dagron would seem to have derived good results from this form of treatment.

SYPHILIS OF THE INTESTINES.

At the last meeting held by the Academy of Medicine, Dr. Lereboullet and Dr. Fournier spoke on a very rare affection: syphilis of the intestine. Dr. Lereboullet spoke of a case where he had found symptoms of enterocolitis, abdominal pains, and discharges resembling the expectoration in pneumonia. There was also noticed pronounced swelling of the cervical, axillary and inguinal glands. The patient had a history of syphilis. Dr. Fournier advised a syphilitic treatment which was carried out by using injections of mercurial oil and iodid of potassium. After eight days the patient had improved very much, and after a certain lapse of time he was quite transformed. Dr. Fournier said he had only seen twelve cases of this disease in his practice, and added that it was often enough not recognized. Dr. Fournier also spoke of infantile diarrhea, which was sometimes due to hereditary syphilis, and would be readily cured by the use of Van Swieten's solution. Mercury was the best preparation, as it had a fundamental and eradicating action, iodid of potassium could only efface temporarily the lesions that had been formed. External treatment should be used so as to spare the intestines and stomach, and frictions or injections employed.

IS SYPHILIS A PRODUCT OF THE OLD WORLD?

It has been a question often brought up to know whether syphilis was of European or American origin. Dr. Zambaco Pacha, who has become quite an authority on such subjects as leprosy, Morvan's disease and kindred affections, presented some photographs of bones found in Upper Egypt, and dating from eighty centuries back. The bones showed indications of suppurative osteitis, and enlargement of the femora, tibiae and peronea was plainly perceptible. The disposition of the lesions was such that Professor Lannelongue admitted that they seemed to be of syphilitic origin. This would seem to prove that syphilis has existed in the Old World from primeval times. Dr. Fournier examined the photographs and declared that there was no sure sign that the lesions were of syphilitic origin. The simple examination of a bone is not sufficient to demonstrate the existence of syphilis. The common varicose ulcer produces sometimes hyperostosed tibiae which are similar to those caused by syphilis; the same can be said of typhoid fever. Dr. Zambaco insisted on the great number of lesions observed, but Dr. Fournier said that tuberculosis would sometimes produce similar results.

Correspondence.

The Plague.

NEW YORK, July 18, 1900.

To the Editor:—May I presume to commend entirely your most timely and well considered article upon the plague situation in San Francisco? The infection with plague of San Francisco's Chinese quarter; the destruction by fire of such a shameful aggregation somewhere in American territory—and the consequent scattering of infective material broadcast—and the tedious and hampered efforts to control the infection, which can not be assured and can only succeed as the result of the most patient attention and inconceivable expense, were matters of confident prediction by me as long ago as the summer of

1896. The present justification of my then statements is not a matter for rejoicing: but I do take unusual pleasure in stating that the only persons who interested themselves in studying the bacillus of plague, which I brought to the United States with me then, were Dr. Douglas W. Montgomery, of San Francisco, and then P. A. Surgeon J. J. Kinyoun, United States Marine-Hospital Service, who was at that time in charge of the bacteriologic laboratory of his service in Washington. The singleness of purpose and the devotion of these gentlemen to true professional ideals are beyond any commendation of mine; but I am endlessly pleased at your thorough recognition of their intents.

In conclusion, I trust that my interest in the matter of plague and my slight acquaintance with it may justify these protests against current errors: 1. That humbleness of station in life and disregard of occidental ideas of personal or of domiciliary cleanliness are essential factors of predisposition to plague. It might, I think, quite as reasonably be contended that a thoroughbred horse is not as liable to glanders as a cart-horse, nor so likely to die of the disease, should it nevertheless have been contracted. I should not consider it a mistake to repeat to the laity, with the iteration and insistence of a machine-gun, that plague, like glanders, and the various specific acute infections to which nearly all animals in their turns are subject, seems to have been designed to *kill*, and that there is much evidence to show that it will do its work fairly well under almost any conditions. Its unusual danger lies in the fact that it may be spread by many more diverse agents—such as the rodents, blood-sucking insects and vermin—than most other acute infectious diseases. 2. The weak, not to say maudlin, wails that there is no specific against its ravages. Haffkine's inoculation is a reasonable protection during active exposure; and Yersin's serum affords more chances of recovery to a victim seized by it than is usually believed, so it seems to me. I hold it greatly to the shame of the hygienists of the United States that the work of this single-minded man of science should have been almost entirely neglected in this country. 3. That street-cleaning and the like is more important than the insuring of correct death certification. By knowing absolutely what is being dealt with and by the correct management of cases as they occur, and *by these means only*, in the humble opinion of the undersigned, may plague be successfully dealt with. And the cost will be large—very!

Yours most sincerely,

W. F. ARNOLD, P. A. Surgeon, U. S. N.

The Paris Exposition.

To the Editor:—I send you this, thinking that a letter relative to medical and surgical matters at the Exposition might be of interest to readers of THE JOURNAL.

The exhibits in this line constitute Class 16 in Group 3 of the catalogue, and include not only everything pertaining to medicine and surgery but also all objects dental and the instruments of the veterinary surgeon. The jury for this class is composed of fifteen members, only three of whom are not Parisians and one of these three is a Frenchman, viz., the noted surgeon, Reverdin, of Geneva. Only one member is not a physician, Leclerc, our secretary, and also a leading local manufacturer of surgical dressings, sutures, etc. But one of the resident members is not a native-born Frenchman, though he has lived in Paris for the past fifteen years. I refer to Dr. de Christmas, who is a Dane, and is physician to the Danish legation here. For the rest, they are men of high standing, being, for the most part, professors to the Faculty of Medicine and members of the Legion of Honor, and all are Fellows of the Academy of Medicine. The president of the jury is Professor Pinard, the obstetrician. Then there are Galippe, chief of laboratory to the Faculty of Medicine; Pozzi, the gynecologist, at present also senator of France; Professor Berger and Dr. Tuffier, the well-known surgeons; Nocard, the head of the Agricultural and Veterinary School at Alfort; Professors Dentu, Hartmann, etc. The two other members are Eschbaum, of the orthopedic firm at Bonn, for Germany, and myself for the United States. Our work, which is now almost completed, has been most interesting, agreeable and instructive.