regions. Patient admitted having taken two pills of diachylon four months before because her menses were late. A week after taking the pills she began to feel dizzy and weak. Examination: A pale, thin girl, with a distinct blue line on gums. Patient is drowsy and at times slightly delirious. She can sit up alone, but cannot stand alone. Knee-jerks absent; plantar reflexes well marked. Her arms are rather weak, but not paralyzed. There is a well marked double optic neuritis. Pupils, eye and face movements normal. No anesthesia. Patient's vision was not examined at this time, but the writer thinks she could recognize people and read large type. During the first month in the hospital her condition remained about the same, except that she became more apathetic and drowsy, and a paresis of the left external rectus developed. Her intense headache and the vomiting continued. Then her symptoms gradually disappeared until she was in perfect health. But when the optic neuritis subsided it was found that the discs were atrophied and the patient was blind. The girl has since remained blind. The treatment was the same as in the previous case.

These three cases illustrate most of the symptoms of lead encephalopathy, and show how closely this disorder resembles cerebral tumor. The fact that the severe symptoms did not appear in these cases for such a long time after the taking of the diachylon is interesting as illustrating the slowness of elimination of the lead from the system.

Ransom thinks it curious that so much reliance should be put upon potassium iodide in the treatment of lead-poisoning, when researches seem to prove that it does not increase the elimination of the metal, and he calls attention to the fact that case I. did not begin to improve on potassium iodide until digitalis was added. He thinks that Epsom salts, with light diet and general tonic conditions, the most useful elements of treatment. The writer cites some cases in which diachylon taken to procure abortion produced toxic symptoms other than cerebral, and he discusses the increasing use of diachylon as an abortifacient.

BONAR.

UEBER DEN DIAGNOSTISCHEN WERTH DES GRÄFE'SCHEN SYMPTOMS UND SEINE ERKLÄRUNG (The Diagnostic Value of Gräfe's Symptom and its Explanation). G. Flatau (Deutsche Zeitschrift für Nervenheilkunde, Vol. 17, Nos. 1 and 2, p. 109).

Numerous observers have found that Gräfe's sign (the appearance of the sclera between the upper portion of the iris and the upper eyelid when the eyeball is directed slowly downward) is not pathognomonic of Graves' disease, and may be absent in pronounced cases of this disease and present in other conditions. Flatau has made a study of the cases in Oppenheim's clinic and has obtained the sign in bulbar paralysis, brain tumor, neurasthenia, hysteria, etc., and even in healthy persons. He was able to produce it in his own person at will.

SPILLER.

SEASICKNESS. H. Partsch (Medical Record, June 9, 1900).

A long experience on ship-board has taught Dr. Partsch that previous preparation is unnecessary and useless. Rooms amidships are preferable and good ventilation is extremely important. For the first few days, lie down most of the time, keeping the eyes closed and avoid excitement or tiring conversation. Whenever the slightest sensation of illness is felt lie down at once and close the eyes. Usually only one pillow should be used, and if very ill no pillow at all. Seasickness is the result of the disturbance of the circulatory apparatus and the accu-