were for the removal of sponges. In most of these thirteen cases the patients came from rural districts, and had been operated on for acute conditions by general practitioners who probably had had very mediocre assistance and were further handicapped by the confusion attendant at such operations. The only wonder is that such accidents do not happen

Fig. 2.—Abdominal sponge with ring attached.

more often in this kind of surgery. After a more or less stormy convalescence, the patient recovers from the operation; but he is left with a persistent sinus, and while he has the original operator to thank for having saved his life, if he learns that a sponge has been left in the wound and is the cause of the sinus, he will seldom hesitate to enter action for unlimited damages.

With a view to the special marking of all abdominal sponges, I have them made with a metal ring placed around the base of the tape and firmly sewed to both sponge and tape with strong thread. If a sponge is reported missing, the abdomen may be roentgenographed before it is reopened; the presence of the sponge will be revealed by the metal band. In hospital practice this procedure should be especially applicable, since if after a morning's work a sponge is reported

Fig. 3.—Roentgenogram showing sponge through a 200-pound patient.

missing and any one of two or three patients may be the unfortunate retainer, roentgenograms may be taken of all of them, and thus a difficult situation is cleared up with a minimum amount of trouble.

The Sin of Ignorance.—Every mind was made for growth, for knowledge, and in its nature is sinned against, when it is doomed to ignorance.—Channing.

USE OF SUPERFICIAL JUGULAR VEINS OF NECK FOR INTRAVENOUS INJECTIONS

DAVID J. KALISKI, M.D., NEW YORK

Not infrequently the practitioner is confronted with the difficult problem of finding superficial veins for the injection of arsphenamin or other arsenical preparations, saline solution or therapeutic serums, and for use in blood transfusion. To overcome this difficulty, I propose the more extended use of the external jugular veins.

In the majority of thin or muscular persons, especially men, the veins at the bend of the elbow in the cubital fossa are easily visible or palpable and should be chosen. Occasionally after numerous injections of arsphenamin these veins become occluded by thrombosis and are no longer available. In some cases these veins are mere threads and unsuitable for repeated injections. In obese persons, especially women, and occasionally in men and children with poorly developed veins, the operator is at a loss to find a suitable vessel into which to inject. Even the most skilful workers encounter these difficulties in hospital and private practice, and for this reason I have been prompted to describe my method of using the external jugular veins.

This method has been in use in selected cases in the wards of the Mount Sinai Hospital and in private practice, and hundreds of injections have been given without a bad result. The procedure is quite simple, presenting no greater difficulties than puncture of the veins of the arm, is not more painful, and is not attended with any greater danger to the patient. Indeed, up to the present time, I have not seen a single instance of thrombosis of

Fig. 1.—Method of making injection into external jugular vein (painted white), and apparatus used by the author.

the external jugular after injection of arsphenamin; and in a few instances the same vessel has been punctured repeatedly over a period of more than a year.

The external jugular veins arise in the region of the parotid gland, on a level with the angle of the lower jaw, and runs down in the direction of the clavicle. They are not infrequently double, and even in children are good sized vessels. They cross the sternocleidomastoid muscle in its upper part, and run parallel to its posterior border; but there are frequent anomalies in size and position. The vessels terminate behind the clavicle in the subclavian veins. In most instances the vessels are superficial, being covered by the skin, superficial fascia and the thin platysma muscle, and stand out prominently especially on forced expiration or inspiration with compression above the clavicle, or on straining or crying. The anterior jugular veins are also occasionally quite large, especially at their lower part, where

they approach and join the external jugulars. They are situated along the anterior border of the sternomastoid muscle.

TECHNIC

The patient reclines on a level table. No pillow is used, the head is turned slightly to one side. Compression may be made over the clavicle with the finger of the left hand in order to make the vein fill up and stand out more prominently while the right hand grasps the needle and pushes it through the skin and into the vein in a downward course in the direction of the vein axis. After practice in the method, almost any type needle may be used; but for greater ease of introduction, a needle with a shield like the writer's needle¹ is preferable. For arsphenamin and other therapeutic injections, a needle with a fairly sharp point, not too large bevel, and about 18 to 20 gage B. & S., about 1¼ to 1½ inches long, is most suitable.

In introducing the needle, the skin over the vein should be gently drawn upward toward the chin. This tends to fix the vein and to allow the needle to pierce the anterior wall of the vessel more easily. The skin of the neck is sometimes quite tough, and therefore the needle should not be too large or too blunt. As the needle enters the vein there is a perceptible "give" as the point enters the lumen. It should be advanced in the vein for about 1 cm. If the needle is properly

Fig. 2.—The external jugular vein stands out prominently because of fluid that is being injected.

in the lumen of the vessel, there is observed a slow dropping of blood from the needle, or a drop of blood comes into the hub of the needle (pressure above the clavicle having been removed) and is seen to advance and recede slightly on inspiratory efforts. It should be borne in mind that there is little bleeding from the jugulars compared with the amount of bleeding observed after puncture of a vein in the arm.

The apparatus best suited for injection is the gravity apparatus, and care should be taken to free the tubing of all air bubbles before beginning the infusion. Saline or water should precede the medicament. If the needle is in the vein properly, the injection will proceed without any pain or infiltration at the site of the puncture. Care should be taken after the fluid has been introduced to remove the needle before the tubing is entirely empty to prevent the entrance of air bubbles. A small glass window made by the insertion of an inch of glass tubing in the rubber tubing near the coupler at the hub of the needle renders this impossible.

The most convenient site for introduction of the needle is in the lower third of the neck, but occasionally the vein is more prominent in the middle third, and the puncture should then be done there.

1070 Madison Avenue.

CROSS-RACIAL TRANSPLANTATION OF TESTES

FURTHER REPORT OF CASE

G. FRANK LYDSTON, M.D., CHICAGO

Through the courtesy of Dr. L. L. Stanley, surgeon to the state penitentiary at San Quentin, Calif., I am privileged to report further on Case 9, previously reported,¹ and to present a photograph of the condition of the implanted testes taken three months after the operation.

It will be recalled that the glands were taken from the body of a negro who had been hanged for murder, and implanted in the scrotum of a white moron, apparently with remarkable results. Supplementing his previous report, Dr. Stanley writes under the date of February 11, about five months after the operation:

"The testicles in the case, a report of which I gave you, have since atrophied very little, and the patient has improved to a great extent. Several days ago he went out to our 'honor camp,' where prisoners are building highways. This camp is about 200 miles distant from the prison, and I shall not have an opportunity to see him until he returns. He is, however, so far improved physically that he is able to do heavy ordinary labor."

Condition of implanted testes three months after operation.

A COFFERDAM INTESTINAL RETRACTOR

R. M. HARBIN, M.D., ROME, GA.

This inexpensive retractor is made in the form of a truncated cone by sewing a section of stocking around pliable copper rings. A dozen or more of different sizes will be found

Cofferdam intestinal retractor.

useful in the isolation of infected areas, in reaching inaccessible sources of hemorrhage and facilitating suturing without heavy packing and in searching for lesions causing intestinal obstruction.

1. Kaliski, D. J.: A Cannula and Needle for Blood Transfusion and Intravenous Infusions, *M. Rec.* 87:482 (March 20) 1915.

1. Lydston, G. F.: Further Observations on Sex Gland Implantation, *J. A. M. A.* 72:396 (Feb. 8) 1919.