CALORIMETRY MADE CONCRETE.

By F. M. GREGG,

State Normal School, Peru, Neb.

In dealing with problems in calorimetry in a high school where apparatus was limited, the writer devised a set of transportation problems which were taken up before the subject of calorimetry was considered and served so well in the after clearing up of that subject that he ventures to present the device for the benefit of the readers of SCHOOL SCIENCE AND MATHEMATICS.

These problems were introduced as a simple diversion from the regular physics lessons, and nothing was said about any after application. Indeed, the student was allowed to make his own discovery and application later. Generally but one lesson was given to the transportation problems. To begin with, a rough sketch of the scene that heads this article was put upon the blackboard, the conditions of transportation explained, and the problems then given out.

In the solution of the problems it is understood that a traveler in passing toward A from D, or any intermediate point, *pays* fare as follows: the distance from D to S is 1,000 miles and the fare on the airship going toward S is $\frac{1}{4}c$ per mile; to be transferred from the airship to the steamboat costs \$5.37 per passenger; the distance from S to Z is 100 miles, and to go any distance on the steamboat toward Z costs $1c$ per mile; it costs 80c to be transferred from the steamboat to the sledge, and the fare by sledge anywhere between Z toward S is $\frac{1}{2}c$ per mile, the total possible distance being 273 miles.

It is further to be understood that in this strange country the traveler *receives* pay for traveling from A or intermediate points toward or to D just what he would have to *pay out* if he were to go his same journey in the opposite direction. Typical problems follow.

PROBLEM 1.—What would be the total cost of a trip from D to A? What would one receive for going from A to D?

PROBLEM 2.—What would one receive for going from A to S but without being transferred to the airship? How does this sum compare with the cost of being transferred from the airship to the steamboat?

PROBLEM 3.
—What would five passengers receive in going from a point 20 miles to the left of Z to a point 20 miles to the right of Z?

PROBLEM 4.
—How much would it cost one to be transferred from the airship to the steamboat, conveyed to Z, and be transferred from the boat to the sledge?

PROBLEM 5.
—A passenger leaves a point 20 miles to the right of S and goes to a point 58 miles to the left of S. How many passengers would have to leave a point 56 miles to the left of Z and go to meet the one passenger and receive for it just what the one passenger had paid out?

PROBLEM 6.
—Five passen-

gers leave a point 112 miles to the right of S and go to meet, somewhere between S and Z, twenty-five passengers coming from a point 10 miles to the left of Z. How many miles from the right of Z must they meet so that the one party spends just as much as the other party receives?

COOLING THROUGH A CHANGE OF STATE.

BY C. K. STUDLEY,

State Normal School, Chico, Cal.

This experiment is neither new nor original but follows the same lines as Exp. 21 in Millikan & Gale's Lab. Course in Physics, but so far as I know the material used is new in this capacity, and is so simple and inexpensive that it ought to be of use to teachers of physics. The experiment referred to above calls for acetamide which is neither very cheap nor easy to obtain.

The apparatus necessary for this experiment is a thin walled test tube, a bunsen burner or alcohol lamp, a test tube holder, a thermometer and some sodium thiosulphate or ordinary hypo such as is used in fixing photos.

Fill the test tube nearly full of the crystals of hypo and carefully melt the crystals by holding over the gas jet or alcohol