The child's expression was one of extreme terror. She died on the third day of cardiac and respiratory paralysis. There was in this case a paralysis of the muscles of deglutition as well as of the muscles of the face and eyes.

CASE 17.—The patient was a boy, aged 6, white. This was a very slow case in development; the child had been sick for six days, complaining of general malaise, with a moderate fever, before any signs of paralysis appeared, which was on the sixth day. This consisted of a partial paralysis of one leg. This case was so insidious in its development that the diagnosis of the consultant and myself was typhoid fever. This proved to be a mild case, with complete recovery.

CASE 18.—The patient, a boy, aged 3, white, had experienced none of the infectious diseases of childhood. When first visited, the patient showed moderate fever and complained of general aching. Paralysis of one leg was discovered on the third day. Beyond this the trouble did not extend. A gradual improvement began in two and one-half weeks from the onset, and in a few days the child was able to walk, but even at the present time there is a decided dragging of the affected leg.

REPORTED BY DR. E. W. DAVIS, SALEM, VA.

CASE 19.—The patient was a boy, aged 3, white. This case presented no points of special interest. Both legs were completely paralyzed. The child has recovered from the paralysis with the exception of a slight dragging of the left foot.

CASE 20.—The patient, a boy, aged 3½, colored, affected with partial paralysis of the left leg and right arm, recovered completely.

REPORTED BY DR. W. P. NORRIS, SALEM, VA.

CASE 21.—The patient, a boy, aged 4, colored, had had roseola, but no history of other disease. When first seen he was constipated and languid; his temperature was 102, his tongue white-coated, and his pulse about 110. On the third day paralysis of left arm and leg developed. Recovery is now nearly complete.

CASE 22.—The patient, a boy, aged 13 months, white, was vomiting, had two or three degrees of fever and very restless when the physician was called. The child seemed to improve after first visit, until the fourth day, when paralysis of the right leg came on. Recovery was almost complete in two weeks; is now complete. The patient had had no previous disease.

CASE 23.—The patient, a boy, aged 3, colored, with negative history, when first seen was paralyzed in the left arm. He could not swallow well for two or three days. Recovery ensued in ten days.

CASE 24.—The patient, a girl, aged 2, white, had no history of previous disease. She vomited, was restless, and had high fever. Her bowels were very loose for several days. Fever continued until the fourth day, when paralysis was first noticed, which affected the left arm and leg. No improvement was noticed for six weeks. The child is now much better, but is not completely recovered, and in all probability will never be. It is feared that the leg will never be normal, as there is now indication of atrophy.

CASE 25.—The patient, a girl aged 15 months, white, had a partial paralysis of the right leg. There was complete recovery in three weeks.

ANALYSIS OF CASE REPORTS

An analysis of these cases shows the following points of interest: The disease was confined to children under 5 years of age, except in one case, a boy of six. The youngest patient was 13 months old. There were 7 cases in which one leg was involved, 1 case in which one arm only was affected, 5 cases in which one arm and one leg were paralyzed, 1 case in which one arm and the leg of the opposite side were affected, 6 cases in which both legs were involved, 4 cases in which all the limbs were paralyzed, 5 cases showing bulbar paralysis, and 3 deaths, which shows a mortality of 12 per cent., higher than is commonly attributed to epidemic infantile paralysis, the authorities stating it to be 6 or 7 per

cent. In the sporadic form of this disease fatal termination is said to be a rare occurrence. In all the fatal cases there was respiratory and cardiac paralysis. A feature in many of these cases worthy of note, because it misleads many of the laity, and possibly some of the profession, who have not had experience with this disease, is the extension of the head, which is due to a paralysis of the neck muscles, in contradistinction to the rigid retraction of the head in meningitis. There were 8 cases of complete recovery, and 14 of incomplete recovery. Each of these incompletely recovered patients has shown some improvement, and it is possible that some of them may yet ultimately recover, as it is stated by M. Allen Starr that "as a rule such improvement is likely to go on for two years." A striking feature in a majority of these cases was pronounced hyperesthesia. Another was profuse sweating. The fact that not more than one case occurred in any house, although in many instances there were other children in the family who associated with the affected child, would seem to prove that the disease is not contagious. Following a decided drop in the temperature of the weather, which occurred in the latter part of July, no new cases developed for three weeks, until a return of the hot wave, which occurred early in August, when two additional cases developed on August 6 and 10, respectively, thus apparently sustaining the belief that hot weather is in some way essential to the development of this disease.

TREATMENT

In regard to the treatment, we regret that our colleagues failed to make any report. We ourselves administered salicylates and applied counter-irritation, either in the form of mustard plasters or of tincture of iodine to the spine during the acute stage. The subsequent treatment consisted of ascending doses of potassium iodide to promote absorption in the inflammatory area in the cord, and strychnin in increasing doses to the point of toleration for its tonic effect. We also employed massage, warm baths, and whenever practicable the galvanic and faradic current. None of these patients have as yet developed deformities, consequently we have nothing to report in the way of orthopedic treatment, although we are well aware of the great importance of orthopedic apparatus and of nerve and tendon transplantation in the later treatment of these cases.

MECHANICAL (DYNAMIC) OBSTRUCTION OF THE BOWEL*

J. SHELTON HORSLEY, M.D.

Professor of Principles of Surgery and Clinical Surgery in the
Medical College of Virginia; Surgeon to Memorial Hospital
RICHMOND, VA.

It is not the purpose of this paper even to touch on all of the various phases of intestinal obstruction, but merely to call attention to a few features of this interesting affection. The first is the abuse of purgatives in this condition. Every surgeon has seen deaths caused by the injudicious use of purgatives. There are few things that require more judgment or skill than the correct exhibition of a suitable purgative at the proper time. Certainly in acute infectious diseases or in any case in which a toxemia is primarily present, a purgative is not only advisable, but absolutely necessary. It is dif-

* Read at the meeting of the Medical Society of Virginia, Richmond, Oct. 23, 1908.

ferent, however, with acute obstruction from any other cause than that which has just been mentioned. A sudden obstruction, not due to nervous affections, always means one of two things: either an inflammatory condition of the peritoneum or else some form of mechanical obstruction, as bands, strangulated hernia, intussusception, volvulus, stricture, or tumor.

When an inflammatory condition is present, purgatives are contraindicated. The great principle of rest is applicable in every form of therapeutics, whether medical or surgical, and inflammation of the outer coat of the intestines is no exception to the rule. This principle is recognized so universally in other instances that it should be superfluous to call attention to it here. A fissure of the anus is best cured by temporarily paralyzing the sphincter, so giving the fissure rest, when it will promptly heal. The object of using splints for broken bones and retentive apparatus in injury of any portion of the body is solely to secure rest for the injured part. In order to give rest we must first find the function of the organ involved and attempt to suspend this function, either completely or partially. Affections of the eye, for instance, would not be benefited by rest in bed, unless light was cut off and the patient prevented from seeing, nor would a darkened room help the patient with a broken leg unless the bones were properly set and retained in a position of rest.

One of the chief functions of the intestines is to propel the food it contains, and this is accomplished by peristalsis. In the inflamed bowel we should apply the principle of rest by restricting this function as much as possible, so we should attempt to inhibit peristalsis rather than accelerate it. If we give purgatives in such conditions, whatever the inflammation is due to appendicitis or to whatever cause, we not only tend to spread the infection, but we fly in the face of the practice we follow when other organs are affected, and instead of lessening the function of the bowel and so giving rest and an opportunity to recuperate we make matters worse by increasing peristalsis with the purgative.

There are some instances in which a purgative apparently does good in inflamed conditions of the peritoneum, because the toxemia from the infection is so great that the purgative causes elimination of the toxins and benefits the patient in a constitutional way, while really making the local condition worse. If such patients were given an abundance of water, and if enemas were used, which empty the lower bowel with a minimum amount of peristalsis, a sufficiently beneficial effect could be obtained in an eliminative way with but a small increase of peristalsis, while at the same time more work would be thrown on those great eliminative organs, the kidneys, by means of the copious supply of water. In this way the excretive function of the bowel would be partially suspended and the rest necessary to repair would be obtained.

When the acute obstruction is due to some mechanical cause, the contraindication for a purgative is even plainer. All purgatives act either by promoting peristalsis in some way or by stimulating the secretion of the intestines. As a matter of fact, while some purgatives act more decidedly in one of these two manners, they all tend to increase peristalsis either directly or indirectly. The more vigorous the purgative in mechanical obstruction the more absolute the obstruction will be. Just as pulling on a string increases the security of the knot, so purgatives by stimulating peristalsis

will make the obstruction from any mechanical cause more complete, whether it be due to volvulus, bands or stricture. Besides this deleterious effect, the increased secretion within the bowel, caused by the purgative, augments the normal contents of the alimentary canal and causes greater distention, which rapidly adds to the seriousness of the condition. Food of any kind in the stomach aggravates the obstruction by giving the bowel more work, thereby stimulating it to greater activity.

The other point to be considered is the surgical method of dealing with the obstruction. The earlier surgical interference is attempted after reasonable efforts have been made with enemas to produce an evacuation, the better for the patient, as the mortality rate increases with each hour's delay. When distention after an obstruction is very marked, frequently an enterostomy is indicated for drainage to carry off the dangerous toxins that have formed. Whether this should be done depends on the findings during the operation, and must necessarily be left to the judgment of the surgeon. When resection is attempted, the method by which the bowel is united is of great importance. A fixed metallic body, with a small lumen for drainage, such as the Murphy button, which by its weight may cause a kinking or volvulus of the portion of intestine in which it has been inserted, is hardly the logical thing to use in such conditions. Intestinal sutures, carefully placed by one who has acquired skill in this line either by clinical or by experimental experience, are much safer and can be inserted in but little longer time than it takes to insert the Murphy button. Of course, when the bowel can not be delivered into the wound and sutures can not be accurately placed, the Murphy button may be indicated; but, judging from my experience, such conditions are quite uncommon.

Since the opening of Memorial Hospital, July 27, 1903, about five years ago, I have done eleven resections of the bowel. These cases will be reported more fully in a subsequent paper, but I will here give a brief summary of them. Of the eleven cases, two were of such a nature that union could not be later accomplished.

In one instance, resection of a large, malignant tumor of the cecum was done, and an artificial anus was made. The patient left the hospital, but soon developed metastasis in the liver, from which he eventually died. There was no time at which a subsequent operation would have been justifiable.

In the other case, which is reported below, the involvement of the rectum by gangrene made union here impossible, so a permanent artificial anus was established.

Of the other nine cases, in one, resection involved the upper rectum, was deep in the pelvis, and on account of the adhesions and raw surface left after removing very adherent pus tubes, suturing was apparently contraindicated. The patient made a satisfactory recovery.

Another patient was a feeble man whose intestine was gangrenous from a strangulated hernia. In order to make assurance doubly sure, a Murphy button was used and sutures were inserted over this. This was one of my early cases, and I believe I made a mistake in using the button, as events subsequently proved, for the patient developed a large fecal fistula evidently due to the presence of the button. The fistula closed rather suddenly two months later.

In seven cases union was effected solely by suturing. Of these patients six recovered uneventfully. One patient died the third day after the operation. He had a strangulated hernia and two different loops of intestines were involved, which made a double resection necessary. The patient was a chronic alcoholic, and considerable chloroform had been administered to him for some time before he came under my care in order to reduce the hernia. A postmortem was held by Dr. E. G. Williams, pathologist to the hospital, and showed a hobnailed liver and other evidences of chronic alcoholism. Both resections were intact, and there was no sign of leakage from either of them, nor was there any indication of peritoneal inflammation. The small bowel contained a few ounces of blood, but hardly enough to affect materially the fatal issue, which was, in all probability, due to the suspension of the functions of the liver and kidneys following the large amount of anesthetic necessary in a man who had been a heavy drinker for many years. Certainly this death can not be attributed to any technical defect in the operation. In none of the cases in which suturing alone was used was there a fecal fistula or subsequent trouble in any way attributable to the resection. The clean and uneventful recovery of these cases so far as the resection is concerned contrasts strongly with the stormy convalescence sometimes seen after a Murphy button has been used, when the formation of fecal fistulas and anxiety about the passage of the button, which may require a subsequent operation for its removal, often afford an occasion for considerable apprehension.

The two cases reported below occurred within three weeks of each other and are selected to illustrate two types of mechanical obstruction in adults.

CASE 1.—Intestinal obstruction from a band. Enterostomy, followed by resection of the bowel and union by suturing.

Patient.—D. T., female, aged 21, colored servant. Admitted to Memorial Hospital, Aug. 2, 1908. Discharged Sept. 9, 1908. Referred by Dr. Angus Nichols, of Richmond, Va. Family history and personal history of no significance.

Present Illness.—Three months ago the patient complained of severe pain in the right side of the abdomen, which was accompanied by vomiting. Three days after this the patient was operated on in Norfolk, but it is impossible to learn the nature of the operation except that drainage was used. She left the hospital after one month and was apparently well until Friday, July 31, when she suffered severe pain in her abdomen, particularly in the right side, with diffuse tenderness and persistent vomiting. On the following Sunday she was referred to me and admitted to Memorial Hospital. After her stomach had been washed out and the administration of everything by mouth discontinued she seemed much better. Soapsud enemas were ordered and a small bowel movement obtained. On the next day, August 3, the patient was considerably worse, with a pulse of 120 and the abdomen becoming much distended. An operation was decided on at once.

Operation.—Under ether anesthesia an incision just to the right of the old median scar was made. A considerable quantity of blood-stained serum was found in the peritoneal cavity. The small intestine was distended and congested. At a point eight inches from the cecum a firm band crossed the ileum and completely occluded its lumen. The band ran from the right iliac fossa to the root of the mesentery. The intestine below this point was collapsed. The band was divided and the bowel underneath it was found to be partly gangrenous; the mesentery for several inches above this point was infiltrated and thrombotic. The patient's condition was becoming rapidly worse, her pulse being 180, and very weak. On this account and because drainage of the distended bowel seemed indicated, an enterostomy was done. The affected loop of intestine was drawn into the wound, sutured to the skin, and surrounded by iodoform gauze. It was then incised and a large

rubber tube inserted in the proximal end. She left the table with a weak pulse of 180.

Postoperative Treatment.—The patient was put in the Fowler position with continuous rectal saline infusion and hypodermoclysis of saline solution. This is, practically, the treatment recommended by Murphy and has proved most efficient, not only in curing peritonitis, but in preventing it. Under this treatment she rapidly improved, and on the following day, August 4, her pulse was 110 and of good volume and there was a profuse discharge of fecal matter through the enterostomy. Her improvement was continuous, and on August 15, twelve days after the enterostomy, she was operated on with a view to re-establishing the continuity of the intestine. For several days before this second operation the patient was given sterile food and careful mouth toilet was done every few hours.

Second Operation.—Under ether, an elliptical incision was made, including the enterostomy, the old scar, and the adjacent skin. The bowel opening had been covered with moist gauze to prevent infection from this source. When the peritoneal cavity was entered very dense adhesions were found around the intestines. They were separated and two clamps were placed on the intestine, one above and another below the enterostomy. All of the bowel between these points, including the enterostomy opening and its adjacent skin, was excised in one mass, the surrounding intestines and peritoneal cavity having been carefully protected with moist gauze. The ends of the bowel were then united by a continuous right-angled suture of linen penetrating all coats. A few reinforcing sutures were placed at the mesenteric border, as the infiltrated mesentery made approximation at this point more difficult than usual. Several inches above the line of union a tear was discovered which had probably occurred during separation of adhesions when the abdomen was first opened. This was closed by a right-angled continuous suture, reinforced by mattress sutures to relieve tension. The appendix was congested and very much enlarged. It was removed, being tied off at its base by a kangaroo tendon which included both the mesentery and the appendix. The stump was disinfected with pure carbolic. The patient's condition had become gradually worse until it was quite desperate. Her pulse was 190 and respirations exceedingly bad, necessitating the use of oxygen and at one time artificial respiration. The abdominal incision was rapidly closed with through-and-through silkworm-gut and a large drainage tube was left in the lower angle of the wound.

Postoperative History.—The patient was put to bed with continuous rectal saline injection and hypodermoclysis of saline solution, and as soon as she had reacted sufficiently was placed in the Fowler position. Under this treatment her pulse gradually came down until by the next day it was less than 120. She made an uninterrupted recovery from this time on, except that she suffered a great deal of pain for the first ten days after the operation. This was to be expected from the large amount of raw surface necessarily left after this operation. At the end of two weeks the pain had practically disappeared. On September 9 she was discharged from the hospital after being up for several days previously. There was a small raw surface about an inch in diameter at the lower corner of the wound where the drainage tube had been inserted. There was practically no discharge from this, as there had been but little from the tube at any time. The patient was comfortable and feeling well and her bowels were moving regularly.

CASE 2.—Volvulus of the sigmoid; gangrene of the sigmoid, rectum and descending colon. Resection of the gangrenous bowel and colostomy.

Patient.—S. F., male, aged about 40, colored. Admitted to Memorial Hospital Aug. 17, 1908; discharged September 11. The patient was a deaf-mute and it was impossible to secure a satisfactory history, but he had evidently suffered from this condition for several days and had not been relieved by purgatives.

Present Illness.—The abdomen was enormously distended and coils of intestine could be distinctly outlined. The patient's pulse was weak and about 140 to 150.

Operation.—The patient's condition was so desperate that an operation under general anesthesia could not be considered. Consequently, cocain was used. I thought that the obstruction

was in the sigmoid and probably due to cancer, so an incision was made over the cecum with the idea of making an artificial anus and doing a radical operation for cancer of the sigmoid later on; an artificial anus in the region of the sigmoid would have made a subsequent radical operation in this region much more difficult. After opening the peritoneum a considerable amount of brownish fluid was evacuated. Toward the median line a large cystic mass was felt. Another incision was made in the median line and disclosed an enormously dilated, slate-colored sigmoid which was at first difficult to identify. The gas it contained was evacuated by a trocar and canula and the intestine drawn into the wound. It was found to be a volvulus of the sigmoid which had turned through 180 degrees. The whole of the sigmoid, the upper part of the rectum, and the descending colon were gangrenous. Along the mesenteric border of the affected intestine were numerous black thrombotic areas. Healthy bowel was found about the splenic flexure of the colon, which was pulled down and the intestine divided at this point. The proximal end was temporarily clamped and all of the gangrenous intestine excised, after ligating the mesentery in sections, down to the rectum. The rectum was gangrenous as far as it could be inspected, and it was not thought proper to leave the rectal stump free in the pelvis, so the intestine was amputated through the affected area and the lower portion, which consisted of the upper part of the gangrenous rectum, was sewed in the lower angle of the wound. A tube was inserted in this stump and was packed around with iodoform gauze. The rest of the wound was approximated with sutures except at the upper angle, where the clamped end of the colon was brought down and sewed to the skin. The end of this bowel was drawn snugly around a tube with a purse-string suture. A large drainage tube was put in the primary opening over the cecum to drain the peritoneal cavity and a liberal amount of gauze surrounded the openings of the intestines in the median line to prevent infection of the wound.

Postoperative History.—The patient stood the operation very well. He was put to bed and the head of the bed elevated about two feet. He was given salt solution by hypodermoclysis. There was an abundant fecal discharge within a few hours. By the next day the patient had improved markedly, though there was some difficulty in respiration, and on September 19, two days after the operation, his pulse was only 100 and of good volume, with a temperature of 100. He made an uninterrupted recovery from this time on. The part of the rectum that was sutured in the wound gradually sloughed away. The patient was up and walking about three weeks after the operation and was discharged in good condition on Sept. 11, 1908, twenty-five days after admission to the hospital. On account of the affection of the rectum and the amount of sigmoid and colon removed it will be impossible to restore the continuity of the intestine and the artificial anus will be permanent.

303 West Grace Street.

THEORETICAL AND PRACTICAL CONSIDERATIONS CONCERNING THE SIGNIFICANCE OF THE CONJUNCTIVAL REACTION (OCULAR TUBERCULIN TEST)*

A. WOLFF-EISNER, M.D.

Bacteriologist to the Krankenhaus Friedrichshain
BERLIN, GERMANY.

The new tuberculin reactions have opened up a wide field in connection with both theory and practice and have revived the discussion of many questions that had come to a standstill. I do not regard it as my task now to take up all of the theoretical questions; the lack of space would preclude this. I shall occasionally refer to the papers in which those interested may find further data that may interest them. I believe it would be best

to pick out a few practical questions and try to come to a conclusion concerning them.

Recently a number of new tuberculin reactions have been placed at our disposal for the diagnosis of tuberculosis. A large amount of literature has followed the first publication about these matters, a great part of which is worthless because of an insufficient knowledge of the fundamental basis of the reactions on the part of the observers. A recital of such examples is not in place. Such publications have caused confusion and dispute, a fact which has made the practitioner avoid this field of contest. But the time of toil and trouble will pass. It was the unavoidable result of an unsubstantiated approval of methods, from which I kept aloof, as is shown in my first communication on the subject.

We have now first the old subcutaneous method of Koch; second, the *Stichreaction*, just as old, but newly applied; third, the cutaneous and the conjunctival reaction; fourth, a series of so-called "new" reactions which I will not discuss.

The subcutaneous reaction is a specific one; it is a very sensitive test. A positive result indicates the presence of tuberculosis, but does not demonstrate necessarily an active or clinically evident tuberculosis. The occurrence of a positive reaction in 50 to 80 per cent. of clinically healthy individuals tested by the subcutaneous method should guard against such an error. I would not mention such a self-evident fact before this assemblage were it not that such errors are made daily in practice and being stated in literature. (Cf. Röpke and others.)

On the whole, the cutaneous and the *Stichreaction* give the same results as the subcutaneous method, which fact establishes their importance for clinical purposes. They are simple, safe methods which can be substituted for the subcutaneous method, demonstrating the same facts. They demonstrate a tuberculous focus, which, however, may be inactive or latent.

The conjunctival reaction differs materially from the cutaneous and the *Stichreaction*, being a reaction on a mucous membrane. It is positive in active tuberculous processes which are not too far advanced, and does not react in those cases in which a clinical examination is negative. It has already been shown that some of the apparently healthy people who gave the reaction had active tuberculosis. The future will tell whether or not tuberculosis is present in the remaining 5 to 8 per cent. But we can say to-day that, in contradistinction to the other tuberculin reactions, the conjunctival reaction is the one which indicates an active tuberculosis. An apparently healthy individual showing a positive conjunctival reaction is to be regarded in a suspicious light as being afflicted with an active tuberculosis.

These facts are proved by the records of reaction in 4,000 cases obtained under all precautions. The diagnostic value can be found only in the positive reaction. I have never stated that the negative conjunctival reaction excluded the presence of active tuberculosis, but it is improbable, except in cases in which clinically an active tuberculosis is present. For this reason the conjunctival reaction has a greater diagnostic value than the subcutaneous reaction. At first we considered them of equal value, but comparative examinations in about 100 cases have convinced us of the greater clinical value of the conjunctival reaction.

The inherent difference between the tuberculin local reaction and the subcutaneous method lies in the pro-

*Read by invitation at the meeting of the New York Pathological Society, Oct. 14, 1908.