testes and cords, a fifth atrophied testicle and perhaps a sixth which is undescended.

It would be interesting to see a section to note where the attachment of the second set joins the vas of the first, and it is not at all improbable that he has also multiple seminal vesicles, although only one prostate is palpable.

A SHORT METHOD OF MAKING NOGUCHI ACETONE-INSOLUBLE ANTIGEN

G. A. MAGNUSSON, M.D., SEATTLE

In order to gain the most information from the Wassermann test, more than one antigen should be used in the "set up." One of the antigens so used should undoubtedly be the

The accompanying chart illustrates the "set up" used in titration.

To each tube, physiologic sodium chlorid solution is added to bring the volume to 1.5 c.c.

Incubation is performed for one hour.

To each tube 0.5 c.c. of sensitized sheep cell suspension (5 per cent. sheep blood cells and 2 units of amboceptor) is added.

Incubation is performed for one hour.

Most of the antigen preparations we have made in this way have had an antigenic unit of 0.02 c.c. of the 1:10 emulsion, while a full cubic centimeter was neither hemolytic nor anticomplementary.

TREATMENT OF CHRONIC GONORRHEAL SKENITIS WITH THE ELECTRIC CAUTERY*

CHESTER B. MOORE, B.S., M.D., SAN FRANCISCO

A not uncommon pelvic focus for the lodgment of the gonococcus is the gland of Skene. In this minute epithelial pocket such an infection may persist indefinitely in spite of local applications. Even if the ducts are exposed and medicinal injections directed into them with a hypodermic needle, the infection may continue without interruption. Often it is possible to express a few or several drops of pus week after week, and month after month, along with such oft-repeated treatment. Furthermore, this minute gland may remain in a chronic state of suppuration without a local symptom, the focus of infection being discovered only in the course of a general pelvic examination. Unless one makes an attempt to express the contents of these ducts, this focus may be entirely overlooked. In this type of gonorrheal infection of Skene's gland nothing less than the complete destruction of the gland will terminate the disease in this locality. This destruction may be easily attained by the use of the thermocautery.

The operation is quite easy and simple, provided one can get a good exposure. For this purpose I take a Kelly endoscope that has been cut off at the proper length so that it will not extend into the bladder, close it with the olive tip, and cut out a small window as shown. I have found that a No. 8 endoscope serves the purpose very well. This makes an instrument, a skenoscope, with a handle that is directed upward out of the way; the entrance is expanded to hold back and protect the labia; the tubal portion protects the urethra except at the point of attack, the gland, the ducts of which protrude into the window cut for the purpose. Under local anesthesia, it is possible to pass a small wire thermocautery

Front, side and top views of instrument devised for use in the treatment of chronic gonorrheal skenitis with the electric cautery.

into the gland without fear of cauterizing any other place than that desired. In order to obtain good light, I have had a small post so placed on the handle that it will carry an E. S. I. light directed toward the aperture. However, any means for obtaining good light will do.

To cauterize the gland, as much discharge as possible is expressed from the ducts. The urethral entrance is cocainized with a 10 per cent. solution for about ten minutes. The instrument is passed, handle upward, until a good exposure is obtained. The entrance of the ducts is sought with a large size hypodermic needle, from which the point has been filed, or with the wire contained in the tubal portion of the instrument. The needle or wire is passed along the duct to the

* From the Division of Obstetrics and Gynecology, Leland Stanford Junior University School of Medicine.

<div> <div>2 units complement</div> <div>0.1 cc. positive serum.</div> </div>		<div>Antigen</div> <div>0.5 cc.</div> <div>1.0 cc.</div>
---	--	--

"Set up" used in titration.

Noguchi acetone-insoluble, as it will give positive reactions with a higher percentage of syphilitic serums than the alcoholic extracts, while it is less likely to give false positive reactions than the more sensitive cholesterinized extracts. But its preparation requires so much time that this becomes quite a factor in a laboratory where considerable quantities are used. With a view of hastening the process of extraction and evaporation we tried this at higher temperatures and find that boiling temperature in no way interferes. In this way we can prepare and titrate an antigen all in one day.

PROCEDURE

To one part of finely minced beef heart, nine parts of commercial ethyl alcohol are added. (We have recently tried the extraction with denatured alcohol and this seems to do as well. As alcohol is denatured according to varying formulas, some may not be usable.)

This is boiled in a water bath for from ten to fifteen minutes, and filtered while hot.

The filtrate is evaporated over a boiling water bath until the alcohol and water are removed.

The residue is extracted with a small amount of ether.

The ethereal extract is poured into a centrifuge tube and centrifuged until the supernatant liquid is perfectly clear. This is added to ten volumes of chemically pure acetone.

The precipitate is allowed to settle. The greater part of the acetone is decanted, and the rest is poured into a clean graduated centrifuge tube and centrifuged for five minutes at moderate speed, the acetone poured off, and drained thoroughly.

To the precipitate, ten volumes of chemically pure methyl alcohol are added, and the mixture is stirred with a glass rod until as much as will dissolve has gone into solution and then centrifuged until the supernatant liquid is perfectly clear.

This is poured off and kept in a clean, well stoppered bottle.

TITRATION OF ANTIGEN

An emulsion is made with one part of this stock alcoholic solution and nine parts of physiologic sodium chlorid solution.

Titration is performed for antigenic, anticomplementary and hemolytic properties.

base of the gland. They will act as a guide to direct the passage of a small, pointed, electric thermocautery until as much tissue as desired has been destroyed. Up to the present, no discomfort or ill effects have followed this procedure. If the first operation has for any reason been unsuccessful, it may be repeated. Also, it is possible to cauterize the glands without passing a wire for a guide.

A NEW PROSTATIC TRACTOR *

A. J. CROWELL, M.D., CHARLOTTE, N. C.

While performing a seminal vesiculectomy some time ago, it occurred to me that the operation could be greatly facilitated by the use of a tractor inserted into the bladder, through

Prostatic Tractor.

the urethra, which has the same curve as that of a sound. For this purpose I therefore designed the one shown in the accompanying illustration.

It consists of two fenestrated blades attached to shafts, the inner one of which has a cable in the curved portion which allows one of the blades to revolve around the other. There are two handles at the outer end of the shaft which regulate the rotation of the blades after insertion into the bladder through the urethra. When inserted, the handles are rotated from each other to open the blades within the bladder. The gland is thus pulled into the operative field and at the same time the handle of the instrument is out of the way of the operator.

A glance will show the superior advantage of the curve in any operation on the perineum when a guide is needed to locate the urethra, prostate, seminal vesicles or bladder. We use it as a routine in perineal prostatectomies and find that by its use it is much easier to expose the prostate, and the danger of getting into the rectum is greatly diminished.

After the tractor is inserted and the blades opened, slight pressure on the handle by an assistant will pull the gland well into the operative field. It is a better guide than the sound ordinarily used for this purpose. It gives not only a urethral guide that is fixed but also a definite location of the prostate gland and membranous urethra, which are firmly held in one position while the gland is exposed.

* From the Crowell Urological Clinic.

Elimination of Inhaled Dust.—In a paper read before the Institute of Mining Engineers on dust inhalation and the health of miners, J. S. Haldane, F. R. S., among other things said that investigations had shown that coal and shale dust were rapidly eliminated from the air passages by special dust collecting cells which phagocyted the dust particles and then carried them away. This, however, does not occur to the same extent with quartz dust, and whereas miners in coal and shale suffer practically no injury from the dust itself, quartz mining results in serious respiratory difficulties, both on account of the sharpness of the dust particles and because of their defective elimination in the manner described. The widespread introduction into coal mines of stone dusting as a means of preventing coal dust explosions has directed attention to the possibility of injury from this cause.

Military Medicine and Surgery

THE TREATMENT OF HYSTERIA

SUCCESSFUL RESULTS OF A RAPID REEDUCATION METHOD

JULIAN MAST WOLFSOHN, M.S., M.D.
(SAN FRANCISCO)

Captain, M. R. C., U. S. Army; Assistant Professor of Medicine,
Leland Stanford Junior University School of Medicine

LIVERPOOL, ENGLAND

The world war has provoked such an immense number of the various types of hysterical disorders that the conception of hysteria can now be studied more easily from every aspect. It has taught us to look on what we ordinarily designate "hysteria" as merely the outward sign, the symptom, of the fundamental causative factor—the hysterical psychic background. So, when we speak of the treatment and cure of these patients, we wrongly say we have cured the hysteria; we have cured only the hysterical symptom. What most physicians in their ardor have left untouched and untreated is the hysterical psyche which gave rise to the symptom which has incapacitated the patient. This is important to remember when the question arises of returning a hysterical subject to the same environment that originally caused the hysterical symptom.

The war has also shown us that hysteria seldom, if ever, occurs during violent emotion, but rather during the contemplative stage succeeding the emotion; also that there is no essential difference, except perhaps in the number and in the intensity of the emotion produced, between the hysteria of warfare and that of civilian life.

Fig. 1.—Patient before and after treatment for hysterical blepharospasm.

Babinski has so far offered the clearest and most concise definition of hysteria: "Hysteria is a pathologic state manifested by disorders which it is possible to reproduce exactly by suggestion in certain subjects and can be made to disappear by the influence of persuasion alone." Instead of "hysteria," which is obviously a misnomer, he has substituted "pithiatism," which means the possibility of being cured by the influence of persuasion.

There have been published volumes on the classification of hysterical disabilities, but very little on their treatment. In Babinski's book on hysteria, for exam-