Postoperative History.—The patient recovered from the anesthetic without vomiting and voided six ounces (urine and boric acid solution), three hours after the operation and without discomfort. The bladder was irrigated with boric acid solution three times daily for the first three days, then twice daily until the sixth day when the urine became almost clear, and by the tenth day the urine voided was all clear normal, and the irrigations of the bladder were stopped. The capacity of the bladder by the tenth day had increased to 8 and 10 ounces, while before the operation it was only from two to four ounces. By the eighth or tenth day all postoperative hemorrhage and irritability of the bladder had ceased. After the third day the patient partook of three regular light meals and was quite comfortable throughout the entire period of convalescence, amusing himself reading and relating funny stories to his attendants. He was up and around on the twelfth day and left the hospital on the seventeenth day. The postoperative temperature never arose above 99.4 F. and the pulse remained normal. After the operation there was no abdominal pain at any time, and no more discomfort at the neck of the bladder than had been present for months previously. The only inconvenience which the patient experienced after the intoxication from the ether had ceased was from the pulling of the abdominal stitches.

The surgeons who have adopted the intraperitoneal cystotomy operation for the removal of malignant tumors of the bladder, all agree in the praise of this method over the perineal and suprapubic routes.

The mortality from the cases reported so far, is less than 10 per cent., whereas by the suprapubic and perineal methods it is from 28 to 47 per cent.

In the hands of the careful operator, by the intraperitoneal route the danger of soiling and infecting the peritoneum is practically nothing. In the case here reported, there was no necessity for drainage and there was no postoperative annoyance to the patient; whereas every surgeon dreads the postoperative annoyance to which the patient is subjected by either the perineal or the suprapubic route, and the surgeon's anxiety relative to these postoperative dangers is practically eliminated by doing the intraperitoneal operation.

This single case, with the ease of approach and the ready access to the growth which the surgeon has at command to the base of the bladder, at once suggests the idea that, under favorable conditions, the enlarged prostate gland may be as easy and safely removed and all hemorrhage controlled by closure of the capsule by suture.

1116 Lake Street.

HEMORRHAGIC EROSIONS OF THE STOMACH WITH THE REPORT OF THE AUTOPSY FINDINGS IN A CASE OF EINHORN'S DISEASE*

N. W. JONES, M.D.
PORTLAND, ORE.

Hemorrhagic erosions of the stomach from the viewpoint of the examination of the fasting organ is not an infrequent clinical finding. In 600 consecutive cases in which this routine examination was made as a part of general diagnosis, there were thirty-five cases, or 5.8 per cent., in which definite erosions of the gastric mucous membrane were found. As shown by the accompanying table, these cases were divided among the following disease conditions: Pernicious anemia, general asthenia, carcinoma of the stomach, carcinoma of the esophagus, cardiospasm, cardiosclerosis, dyspepsia ner-

vosa, Schmidt's fermentation dyspepsia, Einhorn's disease, chronic anacid gastritis, chronic subacid gastritis, chronic acid gastritis, chronic gastrosuccorrhea, chronic gall-bladder disease, chronic intestinal catarrh, spastic obstipation, chronic salpingitis and chronic ulcer of the stomach. Two cases of so-called simple erosion, characterized by sudden severe gastric hemorrhage, afforded no fragments of mucous membrane either by vomiting or later by examination.

From a study of these cases the erosions group themselves logically into, (1) those without clinical significance, and (2) those with positive clinical significance.

Into the first group are placed the fragments of mucous membrane which are, for the most part, accidentally recovered. Chronic gastritis, together with the muscular contractions of the stomach and the trauma from the tube, has afforded thirteen of thirty-four cases. This trauma, together with the contractions of the muscular coats, has seemingly been the entire cause in all of the remaining cases with the exception of the two malignant ones, in which erosions were found. It has been noted also that these erosions were obtained only where there was more or less retching and resistance on the part of the individual. Study of most of the fragments in teased preparations or from sections, in those instances in which the erosions were large enough to be so treated, has not thrown any further light on their etiology. There is always a variable amount of small round cells between the gland acini, but it has been quite impossible to distinguish between round-cell infiltration and normal lymphoid structure. In one large erosion from a case of subacid gastritis "the cuticula and crypts were not broken or disintegrated, but in the deeper portions of the glands the cytoplasm stained faintly and was swollen. In places it seemed to be disintegrating and portions of the glands were missing. There was some small round-cell infiltration, but nowhere was it of marked degree." Other etiologic factors determined by Mintz, Wurtz and Leudet, Dieulafoy, Nauwerk, Gerhardt, Langerhans and E. Fraenkel, were not found.

There are, however, two types of hemorrhagic erosions of the stomach of distinct clinical importance, which present fairly definite clinical pictures. They are (a) the so-called simple erosion of Dieulafoy and (b) the chronic and recurring type of erosion described first by Einhorn and now termed Einhorn's disease.

Of the simple erosion, which is characterized by sudden, severe, at times fatal hemorrhage occurring during normal health, usually without pain and with but little sickness, I do not wish to speak, excepting to emphasize the need of a better recognition of the condition and the employment of rational medical treatment. In one of the four cases seen by me death was hastened, if not actually caused, by surgical measures, and two of the three other patients were strongly urged by competent operators to submit to operation. I think it is now quite agreed among surgical writers as well as among physicians that acute bleeding from the stomach is a condition best handled by medical means and not by surgery. In 1901 Rodman reported thirty-two operations for acute gastric hemorrhage with nine deaths, a mortality of 40.6 per cent. In 1906 Laurent reported 93 per cent. recovery in his patients treated medically. A critical review of the literature since these dates bears out this contention more forcibly, and yet such patients are still being operated on. This fact, in my opinion, justifies the clinical grouping of this class of cases in such a way

*Read in the Section on Practice of Medicine of the American Medical Association, at the Sixty-Second Annual Session, held at Los Angeles, June, 1911.

as will emphasize the importance of their treatment whatever the pathogenesis underlying the condition may be.

According to Einhorn, the disease which was described first by him in 1894, and which now bears his name, is a chronic condition characterized by diffuse stomach pain which comes on shortly after eating and is not influenced by pressure, vomiting or eating again; by a sense of great weakness, a rather rapid loss in weight, and, pathognomically, by the repeated presence of from two to four erosions in the wash water from the fasting stomach. Einhorn's belief that this clinical picture was the expression of a distinct disease has been upheld by Mintz, Berger and others. Pariser, Nauwerk and Elsner, on the other hand, have maintained that the basis of the picture has been a chronic gastritis, the ordinary symptoms of which have been forced into the background by the periodic occurrence of the erosions and the bleeding. Owing to the entire absence of post-mortem examinations of such cases, the following type

Reproduction from drawing showing histologic characteristics of tissue. A, remnants of gland acini; B, blood extravasation in areas of dense, small round-cell infiltration.

ically illustrative case which came to autopsy is given briefly in detail, together with the anatomic findings:

History.—Mr. D., a business man, aged 57 years, began twenty years ago to have, with the coming of each winter, periods of illness characterized by a diffuse sense of burning in the stomach after eating, sometimes on rising, a rapidly developing weakness, paleness and loss of weight. At times there was some nausea and the regurgitation of clear fluid which was not especially sour to taste. All foods produced distress. There was no local tenderness, no fever, chills or sweating. The bowels were regular. Each year the illness lasted several weeks, usually in bed with restricted diet. More or less paleness and weakness would last until warm weather set in, when he would regain his health rapidly, his anemia would disappear and his stomach would again tolerate any kind of food without discomfort. The patient believed a causal relationship existed on the part of the cold and always dreaded the approach of winter. During the illness of the year preceding that of his death occult blood was found in the stools. His last illness came on as they always had before and was of several weeks' duration. Blood was again found in the stools. Dr. Mackenzie tells me that blood examinations showed a very

severe grade of a chlorotic type of anemia. The patient had weighed 202 pounds before his illness began but had lost much flesh during it. His own personal history and that of his family were negative in all respects.

Necropsy.—Through the courtesy of Dr. Mackenzie I made a post-mortem examination on Dec. 20, 1909, six hours after death. Briefly the anatomic findings were as follows: The body was that of a tall, heavily boned, greatly emaciated middle-aged man. The panniculus was pale and scanty; the peritoneum was smooth and glossy. All parenchymatous organs were very pale and dry. The heart and lungs were otherwise negative. Very small scattered atheromatous patches existed in the beginning of the arch of the aorta. They were absent throughout the course of the abdominal aorta. There was no gross evidence of terminal infection. The bone-marrow of the ribs was pale in color. The intestine and colon were more or less filled with dark, tarry content that contained much blood. The stomach was of normal size, blanched, and contained a little turbid fluid. Over the mucosal surface a moderate amount of greyish, somewhat blood-stained, tenacious mucus was found. On the posterior wall about midway between the cardia and the pylorus were seen five small eroded areas with hemorrhagic margins and bases. They were superficial in character, not extending to the submucosa, and varied from the size of a split pea to 1.5 cm. in length and $\frac{1}{3}$ cm. in width. No eroded vessels could be found.

The stomach was given to Dr. Roberg of the medical department of the University of Oregon for histologic examination. He reports as follows:

The mucosal surface of the organ is pale yellowish in color, mottled by several deeply colored and rather circumscribed areas. Sections for examinations were taken through both the pale and the darker portions of the stomach-wall. Microscopically, the dark areas show no difference in appearance from the other. They are pigmented areas of submucous hemorrhages, the coloring of which is derived from dissolved hemoglobin deposited in the interglandular tissue and in the secreting cells of the gastric glands. The interstitial tissue of the epithelial layer of the mucosa is the seat of a dense, small round-cell infiltration. This infiltration is diffuse and more marked in the surface of the mucosa. The deposit of these small round cells is so dense that in many cases the stomach glands have been completely obliterated. Scattered throughout this dense infiltration are islands or portions of glands. Their lumina are distended to two or three times the normal size. The cell cytoplasm is granular and disintegrated and in many places the lumina are filled with a cytoplasmic debris. In the fundus region no differentiation of the gland cells occurs into parietal and chief cells. All gland-cell nuclei stain less deeply than normal. The capillaries in the mucous membrane are markedly dilated along the basement membrane and, in the region of the crypts, show a diffuse extravasation of blood. The blood-vessel plexuses in the submucosal and muscular layers show a thickening of the walls and a slight congestion. The muscular layer, submucosa and muscularis have to a slight degree an infiltration of small round cells.

I submitted also to Dr. Roberg for histologic examination three large erosions from a typical case of Einhorn's disease which I have had under observation for over three years and which I have previously reported. His findings are as follows:

The gastric crypts and cuticular border of the mucosa are quite intact. The neck and fundus portions of the gastric glands are displaced and partly obliterated by a small round-cell infiltration not unlike that described in the previous case but not so dense. The destruction and displacement of the glandular elements, although considerable, are likewise not so marked. Interstitial hemorrhage and congestion are absent.

Thus the microscopic examination of the two cases presents the picture of a form of chronic gastritis characterized (1) by a dense small round-cell infiltration in the interstices of the glandular elements, which when extensive obliterates the gastric glands by pressure; (2) by the disintegration of the cutic-

ular portion of the mucosa as the crypts are displaced by the small round cells; and (3) by the fact that with the disintegration of the cuticula and crypts the congested vessels are exposed and ruptured, which accounts for the gastric hemorrhages occurring during the clinical course of the disease.

An entire absence of proliferation of the connective tissue elements, the engorgement of the superficial blood capillaries and the manner in which the gland cells are destroyed sharply differentiate the process from ordinary chronic gastritis. The claim, therefore, of Elsner and others that the disease is one of ordinary gastritis with the factor of erosion formation added to it is no longer tenable, and neither, on the other hand, is that of Einhorn, for the process must be classified as a type of chronic gastritis. It is, in fact, a chronic, non-proliferative, hemorrhagic gastritis.

In general, I believe it will be well to retain this clinical grouping of hemorrhagic erosions, and to consider them, clinically speaking, as disease entities for the sake of defining their relation to other types of gastric hemorrhage. Simple erosion, on the one hand, probably possesses a manifold etiology. Whether Einhorn's disease likewise does, only further autopsy findings will determine. From the similarity of these two cases, however, it is probable that its pathology will be found to be the same in others.

TABLE OF SIX HUNDRED CASES AND RELATION OF STOMACH EROSIONS TO THEM

Disease.	No. of Cases.	Cases with Erosions.	Erosions Found in Each.
Anemia, pernicious	4	1	2
Anemia, metasyphilitic	1	0	0
Anemia, chlorosis	1	0	0
Aneurysm abdominal aorta	2	0	0
Appendicitis, chronic	3	0	0
Abdominal arteriosclerosis	4	0	0
Arthritis, toxic	3	0	0
Asthma, general	79	1	4
Brain tumor	1	0	0
Carcinoma, stomach	18	1	2
Carcinoma, pancreas	1	0	0
Carcinoma, sigmoid	3	0	0
Carcinoma, liver (see)	1	0	0
Carcinoma, esophagus	1	0	0
Colitis, mucous	1	0	0
Colitis, chronic ulcerative	24	0	0
Cardiospasm	7	2	2, 2
Cardiosclerosis	15	1	1
Cirrhosis stomach	1	0	0
Dyspepsia nervosa	11	1	1
Dyspepsia fermentation (Schmidt)	13	2	2, 1
Erosions of the stomach (Einhorn's disease)	1	1	4
Erosions of stomach (simple hemorrhagic) examined late after bleeding	2	0	0
Epilepsy, petit mal	1	0	0
Fibroid right broad ligament	1	0	0
Gastritis—chronic anacid (inc. achylia gastrica)	47	6	11, 4, 2, 1
Gastritis, chronic subacid	22	3	1, 2, 3
Gastritis, chronic acid	13	2	1, 2
Gastroenterorrhoea, chronic	23	2	1, 1
Gastroptosis (without ulcer)	11	0	0
Gastromyorrhoea	12	0	0
Gall-bladder disease, chronic	1	0	0
Hypochlorhydria	29	3	1, 1, 2
Hypertension, arterial chronic	8	0	0
Intestinal catarrh, chronic	3	0	0
Myotonia congenita	2	0	0
Migraine, chronic	22	1	1
Nephritis, chronic	1	0	0
Neurasthenia	4	0	0
Obstipation, atonic	3	0	0
Obstipation, spastic	31	0	0
Organic heart disease	69	3	1, 1
Polycythemia	11	0	0
Pyloric insufficiency	1	0	0
Urticaria univ.	4	0	0
Rheumatism, chronic	1	0	0
Sacro-iliac joint relaxation	1	1	1
Scoliosis	1	0	0
Tuberculosis, lungs	1	0	0
Tuberculosis, peritoneum	7	0	0
Tubercles	1	0	0
Ulcer of stomach, chronic	2	0	0
Ulcer of stomach, healed	31	3	1, 1
Ulcer of stomach, duodenal	2	0	0
† Many.			
600	35		

ABSTRACT OF DISCUSSION

DR. ROBERT T. MORRIS, New York: In these days when so much credit is given to proper stomach surgery, I wish to emphasize a point made by Dr. Jones that hemorrhagic erosions of the stomach are not cases for the surgeon. Whatever credit comes to surgeons is from wise selection of cases. Surgery does not at all comprehend the question of gastric erosions and does not give relief because the original condition persists. Again, there are the shock and other features which were incidental to operation, features that are not desirable. There was one point, however, that Dr. Jones did not bring out and one which I believe to be very important. We should learn more of the early septic history, and especially of the ordinary medical treatment that was employed at the time the erosions occurred. Of course, the patients must have treatment when these erosions of the stomach occur, but the earlier toxic history should be kept track of in all these cases. We should remember that hyperchlorhydria and hypochlorhydria are symptoms like coughing or sneezing. The hormones send wrong messages to the secreting glands. What reason have we for believing that excreted toxins are basic factors in these erosions? What is found in the peritoneum of the vicinity gives us a clue? We all have found peritoneal adhesions so common in the pyloric region that we must assign some significance to them. We often meet with cases of adhesions, "cobwebs in the attic of the abdomen," which are due apparently to the excretion of toxins by the liver; in some cases the lesion might be due to the actual excretion of bacteria, chiefly of the colon-bacillus group, which pass the liver unharmed at times, but set up trouble in the bile-tract. The endothelium desquamates, lymph is excreted, conglutination occurs and adhesions are then formed and gastric disturbances follow. These hieroglyphics most of you can read and must read. When these hieroglyphics in *bas relief* occur in the peritoneum, the "cobwebs in the attic of the abdomen," we must look for an explanation in the toxic influences preceding erosion. Such toxic influences result in spasm of the terminal arteries situated in this region. Please do not forget the terminal arteries, for we are dealing with them. The toxic spasm of the terminal arteries here results in cell infiltration and other pathologic conditions, such as Dr. Jones has described in his paper.

In the treatment of hemorrhagic erosions of the stomach, surgery should not, I believe, be considered at all; medical treatment alone should be employed; any operation in these cases must result in disaster. If we hark back to the original cause I think we shall find that hemorrhagic erosions of the stomach are the result of excreted toxins.

DR. J. F. HULTGEN, Chicago: Erosions of the stomach are not considered sufficiently in connection with the constant anemia which one meets with in these cases. This fact suggests to me that there might be an anatomic cause for this condition—the symptom-complex might be due to the anemia itself; it might be due, as Dr. Morris has suggested, to some toxic process. The anemia coexistent or coexisting in these cases has not been thoroughly inquired into; more investigations should be made in order to bring out this matter. As a matter of fact, the anatomic basis, so far as was shown by Dr. Jones, was not sufficient to explain the clinical course of the disease. I believe that the gastric hemorrhage from the erosion is nothing but a symptom, and that the underlying morbid process might be of hemolytic nature or of toxic origin.

DR. PHILIP KING BROWN, San Francisco: This condition of hemorrhagic erosions of the stomach is not recognized easily. Hemorrhage occurring in association with pain might be the result of chronic disease of the appendix, so fully described by Moynihan several years ago. He reported twelve cases, the symptoms of which were classical of gastric ulcer; at operation no ulcers were found. In each instance, however, there was found a diseased appendix. When the appendix was removed in these cases all the symptoms of gastric ulcer disappeared. It was my lot to recommend for operation practically the same kind of a case; this patient presented symptoms that made one feel quite positive of his diagnosis. The individual had been under the care of Dr. Boas and Dr. Ewald and other

stomach specialists of note. They all made a similar mistake in the diagnosis. The symptoms were entirely gastric and hemorrhages were quite frequent. Under my care and while on a strict Lenhart diet, with the usual subnormal temperature, there suddenly developed a rise of 4 or 5 degrees above the normal. Evidently there was some pathologic condition in the appendix, as the fever was accompanied by pain, spasm and tenderness at McBurney's point, but it did not seem wise to operate on her at the time. Later at operation a very careful exploration of the gastric region was made, but nothing was found at all. The appendix was examined and removed, because adjacent to it was found a calcified gland and there were other evidences of old disease of this organ in the way of firm adhesions and partial occlusion and thickening. No really satisfactory explanation of the cause of hemorrhage from the stomach in these cases has yet been given. Whatever statements have been made thus far are not at all satisfactory to those who are looking for an explanation, but the point of gastric hemorrhage without ulcer from disease of the appendix is well established and should be borne in mind.

Dr. N. W. Jones, Portland, Ore.: In an exhaustive search of the literature on Einhorn's disease there are found no records of cases coming to autopsy. The pathology of the case reported by me seems to be distinct, but it needs verification by other cases. This condition is entirely distinct from those septic conditions in which stomach bleeding has occurred, as reported by Moynihan and Dieulafoy; and, it would seem to me, can have no relation to them. Its pathogenesis is not known. It does not seem to be that of ordinary chronic gastritis.

BUBONIC PLAGUE

WITH SPECIAL REFERENCE TO THAT OF GROUND
SQUIRREL ORIGIN *

GEORGE W. MCCOY, M.D.

Passed Assistant Surgeon, U. S. Public Health and Marine-Hospital
Service

SAN FRANCISCO

Bubonic plague is a subject with which, fortunately, few of us have to deal. The practitioner in California, however, especially if his lot has been cast in the rural communities, must always keep the disease in mind when any acute febrile case comes under his observation. It is my intention to discuss only the isolated cases that occur in the country districts. Those that have been seen in San Francisco and in Oakland at various times between 1900 and 1908 were clearly associated with the presence of plague among rats, while the cases to which I particularly refer are associated with, and due to, the prevalence of the disease among ground squirrels (*Citellus beecheyi*). While plague in man when contracted from these animals is probably not essentially different from that derived from other sources, usually from rats, the cases that have come under observation are sufficient in number to justify us in examining them with care, especially as they are due to infection from a source that is unusual in other countries.

Two or three years after plague had been recognized in San Francisco, several cases of the disease occurred among persons living in the rural districts in the country across the bay from San Francisco. Surgeon Rupert Blue, of the Public Health and Marine-Hospital Service, noted in one or two instances that the disease occurred in those who had hunted ground squirrels a few days prior to the onset of the illness, and he suggested that since there had been no contact with human plague, and no known opportunity for infection from rats, the dis-

ease might be present in the squirrels, and be communicated from them to man.

During the past three years I have conducted the bacteriologic investigations in eleven cases of plague of ground squirrel origin. Seven of these patients I have seen clinically, and verified the diagnosis by laboratory methods, while in four the diagnosis was made from the bacteriologic examination of tissues taken at autopsy and sent to the laboratory. Six of these patients, all from the known infected zone, gave a clear history of handling ground squirrels within a week prior to the onset of the symptoms. Five gave no such history although they had been in vicinities where plague was known to exist among the ground squirrels at the time, or where it was demonstrated as a result of the investigations set in motion by the discovery of the disease in man. Of these eleven patients, seven were young men between the ages of 13 and 37; one was a boy aged 5; another, a boy aged 7; while two were young women. Six of these proved fatal, while five recovered.

Clinically there were three groups: The severe acute, the mild acute, and the subacute or pyemic. Six belonged to the first group. A typical history of one of these is as follows:

About a week before becoming ill, the man had been squirrel shooting. The first complaint was of pain in the back and in the groin. There was great prostration. The temperature was 101.4 F., pulse 100. The second day, a swelling was made out in the femoral region. The third day, the patient was delirious and remained so until he died. The highest temperature recorded in this case was 103.8 F. but it usually goes to 105 F. or over. Death occurred on the fifth day of the illness.

One of these cases was rather unusual in that recovery occurred after invasion of the blood-stream by the plague bacillus. The evidence of septicemia was the recovery of the bacillus from the blood, and the development of secondary buboes in the regions of the various peripheral lymph glands.

The mild cases were four in number: all ending in recovery, but none was so mild as to justify calling it pestis minor. The history of the most recent case is as follows:

There was no evidence of contact with squirrels, but the patient, a girl, lived in a territory that was known to be heavily infected. The first symptoms were headache and fever. These were followed by pain and tenderness in the right groin. The disturbance was so mild that it was thought sufficient to consult a physician by telephone. However, when improvement did not occur, the patient was visited by the medical attendant, who found a temperature of 104 F. There was a bubo in the femoral region. One week after the onset of the illness the glands were removed under an anesthetic. The patient made an uneventful recovery, never having been regarded as dangerously ill.

The subacute class is represented by only one among the squirrel cases that have come under my observation. This has been reported in detail by former Acting Assistant Surgeon W. B. Wherry and myself.¹ These cases of subacute or pyemic plague are seen occasionally in epidemics of rat origin.

The patient was a youth, aged 13, who had been squirrel shooting about five days prior to the onset of the symptoms. The illness began with headache and vomiting. The next day the temperature was 104 F. Forty-eight hours after the onset of the symptoms, the glands in one axilla were found to be much enlarged and very painful. The nature of the dis-

* Read in the Section on Practice of Medicine of the American Medical Association, at the Sixty-Second Annual Session, held at Los Angeles, June, 1911.

* From the Federal Laboratory, San Francisco.

1. Wherry, W. B., and McCoy, G. W.: Jour. Infect. Dis., November, 1909, vol. 5, p. 670.