III. Zum I. *de Vico'schen* Cometen.

Mittlere Oerter. 1844,0.

Beob.		Beob.	
$\alpha = 0^h 40' 46'' 74$	3	$\delta = -13^\circ 29' 30'' 9$	3
0 45 10,13	3	-12 59 26,4	2
0 58 37,45	1	-10 36 9,7	1
1 9 45,27	2	-5 37 58,6	2
1 12 55,37	2	-2 7 54,1	2
1 13 52,03	1	-2 48 54,3	1
1 18 23,27	1	-1 2 6,4	1
1 18 26,50	2	-1 15 29,7	2
1 22 30,66	2	+0 33 5,5	2
1 27 19,67	3	+0 28 48,7	2
1 27 28,27	1	+4 12 38,7	1
1 27 54,53	1	+0 28 21,2	1
1 35 30,75	2	+4 57 26,7	2

10 Gr. unsich.

IV. Zum II. *Mauvais'schen* Cometen.

Mittlere Oerter. 1845,0

Beob.		Beob.	
$\alpha = 2^h 56' 19'' 76$	1	$\delta = -20^\circ 49' 57'' 6$	1
2 58 8,73	3	-19 40 35,3	3
2 59 4,46	1	-10 51 19,4	1
3 2 4,12	3	-25 29 22,1	3
3 2 11,40	1	-10 10 12,4	1
3 2 47,72	1	-9 43 8,4	1

unsicher.

unsicher.

unsicher.

V. Zum großen Cometen vom Juni 1845.

Mittlere Oerter. 1845,0.

Beob.		Beob.	
$\alpha = 4^h 55' 22'' 82$	2	$\delta = +44^\circ 30' 46'' 3$	2
6 14 49,40	1	+44 48 35,6	1

VI. Zum *Biela'schen* Cometen.

Mittlere Oerter. 1846,0.

Beob.		Beob.	
$\alpha = 6^h 17' 25'' 10$	1	$\delta = -14^\circ 45' 26'' 1$	1
6 18 1,37	1	-14 25 22,5	1
6 41 58,68	1	-14 58 33,8	1
7 18 4,34	2	-14 35 0,0	2

VII. Zum III. *de Vico'schen* Cometen.

Mittlere Oerter. 1846,0.

Beob.		Beob.	
$\alpha = 4^h 44' 17'' 17$	1	$\delta = +23^\circ 3' 11'' 3$	1
4 48 48,69	1	+25 50 35,3	1
5 7 4,00	1	+31 21 46,1	1
5 41 13,66	2	+39 30 53,4	2
5 41 57,00	3	+39 31 40,4	2
5 48 30,71	3	+40 0 39,4	3
5 55 5,08	2	+40 41 26,2	2
6 2 46,47	1	+41 43 41,0	1
6 3 55,76	1	+42 9 39,3	1

VIII. Zum II. *Brorsen'schen* Cometen.

Mittlere Oerter. 1846,0

Beob.		Beob.	
$\alpha = 6^h 5' 38'' 98$	1	$\delta = +57^\circ 11' 19'' 9$	1
6 24 45,82	1	+53 35 28,9	1
6 28 36,29	1	+53 31 40,5	1
6 34 22,18	1	+50 31 50,4	1
6 42 22,21	3	+48 45 32,1	2
6 42 38,41	1	+50 13 32,3	1
6 45 8,17	1	+46 27 45,3	1
6 45 30,58	1	+46 53 51,8	1
6 53 35,72	3	+47 59 39,7	3
6 58 10,91	1	+37 40 45,2	1
6 58 25,44	1	+37 46 7,3	1
6 58 41,00	1	+38 50 51,5	1

IX. Zu dem Cometen von *Hind*, entdeckt im Juli 1846.

Mittlere Oerter. 1846,0.

Beob.		Beob.	
$\alpha = 2^h 14' 11'' 95$	2	$\delta = +60^\circ 50' 51'' 1$	2
2 21 1,89	2	+61 2 49,9	2

v. Littrow.

On the new Planet Neptune.

Von Herrn *Everett* mir eingesandt. S.

Professor *Peirce* communicated to the Academy the following notice of the computations of Mr. *Sears C. Walker*, who had detected a missing star in the *Histoire Céleste Française*, observed by *Lalande* on the 10th of May, 1795, near the path of the planet Neptune, at that date, which may possibly have been this planet.

Shortly after the arrival of the news of the physical discovery of Neptune at Berlin, on a suggestion by Mr. *E. C. Herrick* of its probable identity with the Wartman planet of

1831, Mr. *Walker* engaged in the study of the orbit of the former, and soon concluded that they could not have been the same, and that no set of elements could be found, with a mean distance at all probable, which would represent the four places of Wartman's planet, as published in the *Contes Rendus* for 1836.

His first examination of the orbit of Neptune led to the presumption that the orbit is nearly circular. Also, the large planets lead by analogy to the same conclusion.

The eccentricity of Jupiter is	0.048
" " " Saturn "	0.056
" " " Uranus "	0.047
" " " Neptune "	<0.060, conjectured.

With a small eccentricity, it was impossible for the sun's mass at that distance to impress much daily variation of the radius vector. Accordingly, an approximate solution was made from the places observed on the 26th of September, 26th of October, and 21st of November, on the supposition of a constant radius vector. The concluded true sidereal orbital motions n' , n , and n'' , together with the mean daily sidereal motion μ , for the radius vector r = the semiaxis major = a , are here given.

r	n' First thirty days.	n Average motion.	n'' Last 28 days.	μ
34	12 ^h 8	16 ^h 7	19 ^h 7	17 ^h 90
33	14,6	17,7	20,3	18,71
32	16,6	18,8	20,8	19,60
31	19,4	20,1	21,2	20,56
30	21,7	21,6	21,6	21,58
29	24,1	23,4	22,0	22,67

The most plausible value of r from this table is that in which $(n - n')^2 + (n - n'')^2$ is a minimum. This value by the table is 30, nearly, and for this value we have very nearly $n = n' = n'' = \mu$. Hence the orbit comes out nearly a circle, unless we suppose the planet now to present the possible, but still improbable, case of a great eccentricity and true anomaly nearly 90° .

Accordingly, he selected for the next trial the circular hypothesis, for which two places of the planet sufficed, those of the 26th of September, from the mean of nine European observations, and the 26th of December, from the mean of 33 transits and 11 measures in declination of Neptune (compared with the same two stars used in September) by himself with the Washington equatorial. All the small corrections were taken into account. In this manner he obtained Elements I. in the table below. These elements enabled him to compute an ephemeris of Neptune for the six months following August 1st, 1846, with which he compared 116 nights' works, 70 of the European and 46 of the Washington Observatory, and derived from them 16 normal places, which indicated the following corrections of the geocentric longitude computed from Elements I.

Normal places of Neptune.

t Date, 1846 yrs.		hh		$k'h'$	
No. of place.	Mean Time, Greenwich.	α Obs. Geo. lon.	No. of obs.	δ Obs. Geo. lat.	No. of obs.
1	215 ^d 5670	327° 9' 49"34	1	-0°31' 36"24	1
2	223,5441	326 57 9,04	1	44,09	1
3	270,5	325 46 25,82	16	57,99	16
4	276,5	39 54,23	13	56,94	13
5	282,5	34 16,11	13	56,09	13
6	290,5	28 21,99	12	53,16	12
7	298,5	24 25,25	18	51,13	19
8	306,5	22 32,46	6	47,61	6
9	313,5	22 40,00	4	45,15	3
10	319,5	24 6,40	4	41,51	6
11	325,5	26 50,59?	4	37,30?	4
12	334,5	33 9,44	7	33,92	6
13	345,5	44 26,93	4	30,79	4
14	353,5	54 58,01	2	27,10	2
15	359,5	326 4 2,52	3	26,04	3
16	372,5	326 26 39,11	3	23,60	3
		$\Delta\alpha$		$\Delta\delta$	
		Obs. — Eph.		Obs. — Eph.	
		-16 ^h 75		-0 ^h 63	
		- 7,27		-1,03	
		- 1,02		+0,84	
		+ 0,27		+1,51	
		+ 1,12		+0,03	
		+ 3,13		+0,80	
		+ 4,19		+0,56	
		+ 3,02		+0,23	
		+ 2,40		-0,68	
		+ 1,95		+0,51	
		+ 3,77?		+2,21?	
		+ 2,46		-1,13	
		+ 0,96		-0,03	
		- 0,72		+1,51	
		- 0,23		+0,77	
		- 4,40		+1,28	

The above places are referred to the mean equinox of January 1st, 1847, and mean obliquity. The planets place is corrected for stellar, but not for planetary aberration. It is also corrected for planetary parallax. The residual errors, though small, show in the course of six months a sensible deviation of the orbit from a circular form. They show at the same time that if the eccentricity is greater than 0,06,

the true anomaly must be nearly $\pm 90^\circ$, a possible, but, as was said before, an improbable case.

The next step in the investigations was to make equations of condition of the form, $0 = ax + by + cz + n$, in which x is $50 \times \Delta r$, $y = 10 \times \Delta v$, $z = \Delta \lambda_{300}$, a , b , and c are computed coefficients, v the daily increase of λ , and λ_{300} the heliocentric longitude of Neptune on the 300th day of the

year. Finally, n is the equivalent heliocentric value of $\Delta\alpha$ above with sign changed. The number of equations was reduced to 9, by taking for the first the one third of Nos. 1 and 2, above. Then follow the next five unchanged, then the mean of Nos. 8, 9, and 10. No. 11 is rejected, then the mean of Nos. 12 and 13. Lastly, the mean of Nos. 14, 15, and 16. The 9 conditional equations have then equal weight, and stand thus:—

				Residual Error. ϵ
$0 = -0,303 \times x - 2,700 \times y + \frac{1}{3} \times z + 3''88,$				$-0''08$
$= +3,016$	$-3,000$	$+1$	$+1,00,$	$+0,49$
$= +3,363$	$-2,400$	$+1$	$-0,27,$	$+0,19$
$= +3,685$	$-1,800$	$+1$	$-1,10,$	$+0,22$
$= +4,038$	$-1,000$	$+1$	$-3,07,$	$-1,03$
$= +4,268$	$-0,200$	$+1$	$-4,12,$	$-1,31$
$= +4,594$	$+1,267$	$+1$	$-2,44,$	$+1,03$
$= +4,248$	$+3,950$	$+1$	$-1,73,$	$-0,13$
$= +3,332$	$+6,133$	$+1$	$+1,81,$	$-0,16$

The solution by least squares gives,

$$\begin{aligned} 0 &= 118,879 \times x + 7,477 \times y + 30,443 \times z - 45''629. \\ 0 &= 7,477 \times x + 85,149 \times y + 0,250 \times z + 1,687. \\ 0 &= 30,443 \times x + 0,250 \times y + 8,111 \times z - 8,627. \end{aligned}$$

Whence, $\sum \epsilon \epsilon = 4''21$

$$x = +3,255712$$

$$y = -0,272963$$

$$z = -11,1475$$

$$r = 29,939950 + \frac{x}{50} = 30,00506$$

$$n = \frac{v_{359} - v_{270}}{t_{359} - t_{270}}, \text{ corrected for aber.} = 21''65789$$

$$a = \frac{1}{\frac{2}{r} - \left(\frac{rn}{k^2}\right)^2} = 30,20058$$

$$k = \text{Gauss's constant of the earth's velocity}$$

$$\mu = ka^{-\frac{3}{2}} = 21''37881$$

$$\text{Period} = T = 165,97030 \text{ tropical years.}$$

Thus it appeared that Elements II., assuming the eccentricity and perihelion point unknown, and neglecting the daily variation of the radius vector, would give an ephemeris following the planet's path for five months so closely, that the sum of the squares of 9 comparisons of theory with observation was only $4''21$. This residual quantity might have been still further reduced by inserting a fourth term of the form du , in which u is the daily increase of r_{300} , and d , a coefficient of the form

$$d = a \Delta r + \left(\frac{d\lambda}{dt}\right) \Delta t,$$

where a is the former coefficient of x , and $\left(\frac{d\lambda}{dt}\right)$ is the daily variation of λ for conservation of areas. Since these terms become sensible in the course of a few additional months, it

was thought preferable to postpone the research after the final values of e and π , and by assigning them suitable limits, that of $e < 0,06$, and to π its corresponding value from the equation, $\cos. \nu = \frac{a(1-e^2)-r}{er}$, then to compute the locus of Neptune's orbit for these limits for any given date, and search for an observation of a missing star in Neptune's path on the same night in some of the ancient catalogues. The fact that $(n-\mu)$ is at this time only $0''28$, shows the limit of $\nu < \pm 90^\circ$. The following table of limits of ν was computed.

Assumed values of e .	Concluded values of ν .
1,00	$\pm 90^\circ 0$
0,06	$\pm 87,2$
0,05	$\pm 85,4$
0,04	$\pm 83,0$
0,03	$\pm 79,2$
0,02	$\pm 72,2$
0,01	$\pm 50,1$
Minimum limit, 0,006474	0,0

Now, since all values of $\pi = (\omega - \nu)$, ω being the longitude on the orbit, are a priori equally probable, and since the maximum value of ν is $\pm 90^\circ$, for $e = 1$, we have the a priori probabilities for e as follows:—

Limit of e .	A priori probability.
Limit $e > 0,06$ and $< 1,00$	$\frac{2,7}{9,0}$
$e > 0,05$ „ $< 0,06$	$\frac{1,8}{9,0}$
$e > 0,04$ „ $< 0,05$	$\frac{2,4}{9,0}$
$e > 0,03$ „ $< 0,04$	$\frac{3,8}{9,0}$
$e > 0,02$ „ $< 0,03$	$\frac{7,0}{9,0}$
$e > 0,01$ „ $< 0,02$	$\frac{22,1}{9,0}$
$e > 0,006474 < 0,01$	$\frac{50,1}{9,0}$

This a priori probability that e falls between 0,01 and 0,006474 of $\frac{5}{9}$, is derived from a theory, which in a half-year's path of Neptune presents throughout a probable discrepancy of $0''49$ between theory and observation.

The next inquiry is, how far this value of e is consistent with the equations of condition between e and a , derived from the residual perturbations of Uranus.

From the two full computations of Mr. Adams's Supplement to the Nautical Almanac for 1852, for values of $\frac{a}{a'} = 0,50$ and $0,51$, e varies from 0,16103 to 0,12062. Hence Mr. Walker found the conditional equation,

$$e = 0,16103 \left[\frac{0,12062}{0,16103} \right] \left(\frac{\log. \frac{a}{a'}}{\log. \frac{1,03}{1,03}} \right).$$

Whence for $a = 30,20058$, $e = 0,0153883$, which is the eccentricity from Adams's computations, with this value of the mean distance. The mean longitude of Neptune, according to Mr. Adams's remark, also comes out right for this hypothesis.

It remains to consider *Le Verrier's* limits in his additions to the *Connaissance des Temps* for 1849. In his first solution, he gives for the minimum limit of the mean longitude of Neptune for 1800, 234° , whereas *Elements II.*, with $e < 0,0153883$, would require at that date a mean longitude of 226° . In his final solution, *Le Verrier* finds the most probable value 240° nearly. The limit $\pm 5^\circ$ gives for the minimum 235° . If it be asked why *Le Verrier* and *Adams* differ in their conclusions, it may be answered, that they differ in their residual perturbations required, from the more complete computations of *Le Verrier*.

It was during a discussion of this subject by Professor *Peirce* and Mr. *Walker*, that the possible omission of some inequality of long period by *Le Verrier* was suggested by Professor *Peirce*. On comparing the mean motions of Uranus and Neptune by *Elements II.*, it was found that if the mean distance of Neptune is 30 nearly, such an inequality of great power has been omitted. Thus we have

$$\begin{aligned} \text{For Uranus, } \mu' &= 42''2331 & \mu' &= 42''2331. \\ \text{Elements II., Neptune, } \mu &= 21,3788 & 2\mu &= 42,7576. \\ & & 2\mu - \mu' &= 0,5245. \end{aligned}$$

Instead of this, *Le Verrier* retained only the inequality from $(3\mu - \mu')$, which was suited to the mean distance, 38.

It is impossible to decide, without a revision of the calculations of *Le Verrier*, substituting the new inequality depending on $2\mu - \mu'$, whether the limits would not be so far modified as to include Mr. *Walker's* *Elements II.* The inequality of $(2\mu - \mu')$, if the mean distance is nearly 30, is the most remarkable yet discovered in the primary solar system, and merits a thorough analytical investigation.

In conclusion, then, it may be remarked, that the hypothesis of a very small eccentricity is strongly probable by the *Elements II.*, probable by *Adams's* computations from the residual perturbations of Uranus, and not necessarily contradicted by *Le Verrier's* computed limits, unless we admit also that they exclude at the same time the possibility of the semi-axis major which results so directly from Mr. *Walker's* normal places.

It remains to consider the question, whether any light can be thrown on the subject of the orbit of Neptune by the ancient catalogues. On this head it may be remarked that *Bradley*, *Lacaille*, and *Mayer* seldom observed stars of the magnitude 7,8. In the first three volumes of *Piazzi's* original observations, now in course of publication by the Vienna Observatory, there is no one of those „not found in the catalogues,” that was near enough to the path of Neptune, on the night of observation, to authorize the supposition of its having been that planet. *Bessel*, in preparing his zones, never

swept so far south as the actual position of Neptune. The *Paramatta Catalogue* seldom extends north of the 33° parallel of south declination. The *Madras* observations were generally confined to the stars of *Piazzi's* or *Baily's* Catalogues. The only remaining chance at present for finding an ancient observation of Neptune (though doubtless others will be found hereafter) was in the *Histoire Céleste Française*.

Mr. *Walker* found that *Lalande* had twice included the Neptunian region in his sweeps, viz. May 8th and 10th 1795. Accordingly, he computed the locus of Neptune on the latter night for all values of e , from 0,006474 to 0,06, and for the two cases of $\pm \nu$ at the present time. This locus referred to the mean longitude and obliquity for 1800, so as to compare with *Hussey's* Hour XIV. in the *Catalogue* of the Berlin Academy, is as follows:—

Locus of Neptune, May 10th, 1795, for various Eccentricities.

		Neptune's R.A. 1800. Dec. 1800.	
For $+\nu$, and for $e = 0,06$		$13^h 45^m 50^s$	$-9^\circ 3' 1''$
„ „ „ $e = 0,05$		$13 49 48$	$-9 24,6$
„ „ „ $e = 0,04$		$13 53 51$	$-9 47,0$
„ „ „ $e = 0,03$		$13 57 52$	$-10 8,6$
„ „ „ $e = 0,02$		$14 1 56$	$-10 29,6$
„ „ „ $e = 0,01$		$14 3 52,2$	$-10 40,40$
For $\nu = 0$	$e = 0,006474$	$14 9 18,0$	$-11 8,75$
For $-\nu$	$e = 0,01$	$14 12 9,1$	$-11 23,46$
„ „ „ $e = 0,02$		$14 16 36$	$-11 44,5$
„ „ „ $e = 0,03$		$14 20 35$	$-12 6,1$
„ „ „ $e = 0,04$		$14 24 29$	$-12 25,3$
„ „ „ $e = 0,05$		$14 28 19$	$-12 44,2$
„ „ „ $e = 0,06$		$14 32 8$	$-13 2,6$

Mr. *W.* then selected from the *Histoire Céleste* all the stars within $15'$ of the locus of Neptune in the above table.

No.	Mag.	R. A. 1800.	Dec. 1800.	Authority.
1	9,10	$13^h 50^m 36^s$	$-9^\circ 24' 0''$	L ¹
2	7,8	$13 52 48$	$-9 58,8$	L ¹
3	7,8	$13 52 53$	$-9 45,7$	L ¹ B ¹
4	8,9	$13 57 13$	$-10 11,7$	L ¹ B ¹
5	9	$13 59 54$	$-10 26,4$	L ¹ B ¹
6	8	$14 10 0$	$-11 26,5$	L ¹ B ¹
7	8	$14 12 0$	$-11 8,3$	L ¹ B ¹
8	7,8	$14 12 0,9$	$-11 20,96$	L ¹ missing star.
9	8	$14 29 37$	$-13 10,7$	L ¹ B ¹

The only stars in this list, not found in *Bessel's* Zones, are Nos. 1, 2, and 8. Of these No. 1 is too small. No. 2 is too far south ($17'$) of Neptune's computed path. No. 8 is within $2'$ of the computed path, and if missing now in the heavens may have been the planet. This comparison was made by Mr. *Walker* on the 2d of February, a cloudy night. He immediately, by letter of that date, notified the Superintendent of the Observatory, Lieutenant *Mauray*, of his expectation that on examination of that region on the next clear night No. 8 would be missing.

On the 4th of February, Prof. *Hubbard* examined the region and verified the expectation of Mr. *Walker*. The star was in fact missing.

Here, then, was a presumption in favor of their identity. Mr. *Walker* believed his limits sufficiently extensive to comprise the Neptunian region of May 8th and 10th, 1795. The planet was of the size 7, 8 magnitude, seldom omitted by *Lalande*. No star except No. 8 fulfilled all the conditions.

There is, however, a : placed by *Lalande* after the observation of the star No. 8, which indicates that the declination is doubtful $\pm 5'$. Mr. *Walker's* attention was first called to this circumstance by Prof. *Peirce*. The entries in the *Histoire Céleste* are as follows: —

Mag.	Wire I.	Mid. Wire.	Wire II.	Zen. Dis.
7,8		14 ^h 11 ^m 23 ^s .5		60° 7' 19" : (now missing.)
2 $\frac{7}{8}$ 6		14 12 3,4		59 33 59
8			14 ^h 11 ^m 50 ^s .5:	59 54 40

If then the two: only indicate a doubt of 5' there is no contradiction of the possible identity of the star 7,8 mag. and the planet. If, however, the two: may be so construed as to make the 1st and 3d star the same in declination as they are in right ascension, then no star is missing, and the heavens are now as in *Lalande's* time. The difference of magnitudes 7,8 and 8 militates against this supposition, that stars 1 and 3 are the same. *)

Mr. *Walker* concludes by remarking, that he has stated all the circumstances known to him favorable or unfavorable to the supposition of identity of the star and planet. The decision of the question must be the work of time. In order to establish the priority in determining these elements if the identity should be confirmed, he had computed his Elements III. upon this hypothesis of identity. The three sets of elements are here given, referred to the mean equinox of January 1st, 1847, and to mean time Greenwich.

Elements of Neptune.		Circular Hypothesis		Elements if the missing star was Neptune.	
		I.		Elliptic Hypothesis	
				II.	
				III.	
Longitude of perihelion	π	unknown		0°12'25".51	
" " ascend. node,	Ω	129°48'23".16		131 17 35,80	
Inclination,	i	1 45 19,88		1 54 53,83	
Long. of epoch, Jan. 1, 1847,	ϵ	unknown		328° 7' 56".64	
True heliocentric long. on orbit, Sept. 28, 1847	w	326°59'41".50		326°59'34".74	
Eccentricity,	e	0.		0,0088407	
Radius vector, Sept. 28, 1847,	r	29,93995		30,00506	
True sid. orb. mot., "	n	21"65857		21"65789	
Mean distance,	a	29,93995		30,200385	
Mean daily sidereal motion,	μ	21"65857		21"37881	
Period in tropical years,	T	163,78259		165,797030	
				166,738134	

Professor *Peirce* remarked, that the orbits given by Mr. *Walker* differ so widely from the predictions, that he has been induced to make a careful reëxamination of the observations. He has not only himself verified Mr. *Walker's* distance of 30, and the consequent angular motion; but Mr. *George P. Bond*, of the Cambridge Observatory, has also, at his request, verified this distance and motion from the Cambridge observations alone. From these data, without any hypothesis in regard to the character of the orbit, he has arrived at the conclusion, that the planet Neptune is not the planet to which geometrical analysis had directed the telescope; its orbit is not contained within the limits of space which have been explored by geometers searching for the source of the disturbances of Uranus; and its discovery by *Galle* must be regarded as a happy accident. Mr. *Adams*, in his „Explanation of the Observed Irregularities of Uranus,“ considered two hypothetical orbits, in one of which the mean distance is 38,4,

or just double that of Uranus, and in the other it is 37,6; while M. *Le Verrier*, in his „Researches into the Motions of the Planet *Herschel*, called Uranus,“ after deriving some rough approximations from the consideration of the mean distance 38,4, proceeds to the accurate examination of the three distances 39,1, 37,6, and 36,2. The extension of the investigations to any other mean distances can be made only by assuming a continuous law to pervade the subject of inquiry, and that there is no important change in the character of the resulting perturbations. Guided by this principle, well established, and legitimate, if confined within proper limits,

*) Herr *Walker* konnte nicht wissen dafs Herrn *Mauvais* Nachforschungen in *Lalande's* Manuscripten es aufser Zweifel gesetzt haben, dafs der am 10^{ten} Mai 1795 beobachtete Stern der Planet war.

M. *Le Verrier* narrowed with consummate skill the field of research, and arrived at two fundamental propositions, namely,—

1st. That the mean distance of the planet cannot be less than 35, or more than 37,9. The corresponding limits of the time of sidereal revolution are about 207 and 233 years.

2d. „That there is only one region in which the disturbing planet can be placed, in order to account for the motions of Uranus; that the mean longitude of this planet must have been, on January 1st, 1800, between 243° and 252° .“

Neither of these propositions is of itself necessarily opposed to the observations which have been made upon Neptune, but the two combined are decidedly inconsistent with observation. It is impossible to find an orbit, which, satisfying the observed distance and motion, is subject at the same time to both of these propositions, or even approximately subject to them. If, for instance, a mean longitude and time of revolution are adopted according with the first, the corresponding mean longitude in 1800 must have been at least 40° distant from the limits of the second proposition. And again, if the planet is assumed to have had in 1800 a mean longitude near the limits of the second proposition, the corresponding time of revolution with which its motions satisfy the present observations cannot exceed 170 years, and must therefore be about 40 years less than the limits of the first proposition. Neptune cannot, then, be the planet of M. *Le Verrier's* theory, and cannot account for the observed perturbations of Uranus under the form of the inequalities involved in his analysis.

It is not, however, a necessary conclusion that Neptune will not account for the perturbations of Uranus, for its probable mean distance of about 30 is so much less than the limits of the previous researches, that no inference from them can be safely extended to it. An important change, indeed, in the character of the perturbations takes place near the distance 35,3; so that the continuous law by which such inferences are justified is abruptly broken at this point, and it was hence an oversight in M. *Le Verrier* to extend his inner limit to the distance 35. A planet at the distance 35,3 would revolve about the Sun in 210 years, which is exactly two and a half times the period of the revolution of Uranus. Now, if the times of revolution of two planets were exactly as 2 to 5, the effects of their mutual influence would be peculiar and complicated, and even a near approach to this ratio gives rise to those remarkable irregularities of motion which are exhibited in Jupiter and Saturn, and which greatly perplexed geometers until they were traced to their origin by *Laplace*. This distance of 35,3, then, is a complete barrier to any logical deduction, and the investigations with regard to the outer space cannot be extended to the interior.

The observed distance 30, which is probably not very far from the mean distance, belongs to a region which is even more interesting in reference to Uranus than that of 35,3. The time of revolution which corresponds to the mean distance 30,4 is 168 years, being exactly double the year of Uranus, and the influence of a mass revolving in this time would give rise to very singular and marked irregularities in the motions of this planet. The effect of a near approach to this ratio in the mean motions is partially developed by *Laplace*, in his theory of the motions of the three inner satellites of Jupiter. The whole perturbation arising from this source may be divided into two portions or inequalities, one of which, having the same period with the time of revolution of the inner planet, is masked to a great extent behind the ordinary elliptic motions, while the other has a very long period, and is exhibited for a great length of time under the form of a uniform increase or diminution of the mean motion of the disturbed planet. But it is highly probable that the case of Neptune and Uranus is not merely that of a near approach to the ratio of 2 to 1 in their times of revolution, but that this ratio is exactly preserved by those planets; for it may be shown, as was shown by *Laplace* for the ratio two fifths, that a sufficiently near approach to it must, on account of the mutual action of the planets, result in the permanent establishment of this remarkable ratio. Thus, if

ν = the mean longitude of Uranus,

ν' = that of Neptune,

$V = 2\nu' - \nu$; and if D expresses the differential coefficient relatively to the time, a near approach to the ratio of 2 to 1 gives the equations,

$$D^2\nu = p \sin. (2\nu' - \nu + A) = p \sin. (V + A),$$

$$D^2\nu' = q \sin. (2\nu' - \nu + A) = q \sin. (V + A);$$

in which p , q , and A are known functions of the masses and different elements of the orbits. These equations give at once

$$D^2V = 2D^2\nu' - D^2\nu = (2q - p) \sin. (V + A),$$

which, multiplied by $2D\nu$ and integrated, becomes

$$DV = \sqrt{[H^2 - (4q - 2p) \cos. (V + A)]},$$

in which $H = 2n' - n$;

if n = the mean motion of Uranus,

and n' = that of Neptune.

It follows from the value of DV , that if

$$H^2 < 4q - 2p,$$

$V + A$ cannot increase indefinitely, and that, therefore, the term $(2n' - n)t$,

upon which its indefinite increase depends, must vanish, or in other words $2n' - n = 0$,

and $V + A$ must oscillate in value either

about zero when $2p - 4q$ is positive,

or about 180° when $2p - 4q$ is negative.

The probability of the occurrence of this ratio depends, it will be seen, upon the magnitudes of p and q , which are always of opposite signs. It is evident, from inspecting the computations of Mr. *Walker*, that Neptune's period of revolution is not less than in his second hypothesis of 166 years; and Professor *Peirce* infers from the investigations which he has already made, that a period of $166\frac{1}{2}$ years, which

involves only a slight additional eccentricity, is already a sufficiently near approximation to establish the exact permanency of the period of 168 years. As soon, then, as there may be observations sufficient to prove that Neptune revolves in more than $166\frac{1}{2}$, and in less than $169\frac{1}{2}$ years, the conclusion is inevitable, that its year is precisely twice as long as that of Uranus.

Berichtigungen.

Astron. Nachr. Nr. 595	pag. 307	Zeile 2 v. oben	statt	Z. 192 Nr. 104	lies	Z. 90 Nr. 104
_____	_____	596 „ 317 „ 26 „	„	März 20	„	März 21.
_____	_____	596 „ 317 „ 27 „	„	März *21	„	März *22.
_____	_____	597 „ 339 „ 10 „	„	observations	„	observations.
_____	_____	597 „ 339 „ 16 „	„	3 ^{em} fil	„	3 ^{eme} fil.

A n z e i g e.

Es ist schon in den früheren Bänden dieser Nachrichten bemerkt, daß ohne ausdrückliche Bestellung und Vorausbezahlung keine Nummer eines neuen Bandes versandt wird. Die Herren Abonnenten, welche diese Blätter fortzusetzen wünschen, werden also ersucht, um Unterbrechungen zu vermeiden, baldmöglichst ihre Bestellungen einzusenden.

Man pränumerirt mit 8 $\frac{1}{2}$ Hamburger GrobCourant, und von diesem Preise wird auch den Buchhandlungen und Postämtern kein Rabatt gegeben, die also nothwendig ihren Abnehmern höhere Preise berechnen müssen. Ueberhaupt sind alle in dieser Anzeige bemerkten Preise, Nettopreise.

Einzelne Nummern werden nur zur Completirung, wenn sie vorrätbig sind, à 4 ggr. abgelassen.

Da sehr wenig Exemplare mehr gedruckt werden als bestellt sind, so kann ein Band, der schon geschlossen ist, nicht unter 12 $\frac{1}{2}$ Hamburger GrobCourant verkauft werden. Die einzige Ausnahme ist, wenn alle schon geschlossenen Bände, vom 5^{ten} (inclusive) an, auf einmal genommen werden, wodurch keines von den wenigen noch übrigen Exemplaren des ganzen Werkes incomplet gemacht wird. In diesem Falle wird der Band nur zu 8 $\frac{1}{2}$ gerechnet. Die 4 ersten Bände sind ganz vergriffen.

Man ersucht, um Unterbrechungen in der Zusendung zu vermeiden, die Bestellungen auf den nächsten Band sobald als möglich zu machen.

I n h a l t.

(Zu Nr. 597.) Auszug aus einem Briefe des Herrn Professors *Hansen* an den Herausgeber p. 325. —

Schreiben des Herrn *Hind* an den Herausgeber p. 331. —

Schreiben des Herrn *Hackel*, Astronomen auf der Freiherrlich Senftenberg'schen Sternwarte, an den Herausgeber p. 333. —

Sternbedeckungen, beobachtet auf der Freiherr v. Senftenberg'schen Sternwarte p. 335. —

Schreiben des Herrn Dr. *Jahn* an den Herausgeber p. 337. —

Schreiben des Herrn *Le Verrier* an den Herausgeber p. 339. —

Anzeige p. 339. —

(Zu Nr. 598.) Beobachtungen der Astraea im Frühjahr 1846, von *Otto Struve* p. 341. —

Observations on the Comet of March 4 1847. von *W. C. Bond* p. 355. —

Beobachtungen der Astraea auf der Hamburger Sternwarte p. 355. —

(Zu Nr. 599.) Beobachtungen der Astraea im Frühjahr 1846, von *Otto Struve*. (Beschlufs.) p. 357. —

Ueber die Benutzung der Venus-Stillstände zur Bestimmung der Sonnenparallaxe p. 363. —

Schreiben des Herrn *J. C. J. Schmidt* an den Herausgeber p. 367.

(Beilage zu Nr. 599.) Schreiben des Herrn *v. Littrow* an den Herausgeber p. 373.

On the new Planet Neptune p. 375. —

Berichtigungen und Anzeige p. 387. —