With loss of the higher center, say in the left hemisphere, we have loss of all vision in the right eye, except in a small central area supplied by the right hemisphere, along with loss of vision in the left eye, except in a central area larger than that in the right, and also supplied by the right hemisphere. Vision is lost in the periphery because both higher centers are required for its fulfilment, one being unable to carry on its functions in either eye. Here we may recall the final stage of the spectra in the case which I have fully reported, in which the periphery of both eyes was blind, the central region being sensitive.

The above facts being beyond dispute, Gowers has explained the relationship of the peripheral fields in the two eyes by the theory that when the peripheral visual fibers from the lower center pass to the higher center of the opposite side, they do so by traversing the higher center of the same side, and are therefore involved in disease of either higher center. Each higher center thus containing fibers from the complete periphery of both eyes, an interference with one higher center is sufficient to annul the functions of the complete periphery of both eyes. This theory may or may not be correct, but the bilateral spectra, as illustrated by the cases presented, seem to show a close connection between the visual centers in the two hemispheres, or even that they act almost as one center.

It is interesting to observe here the curious fact noted by Gowers, that the field for color in every area, even in the temporal where the color is seen by one eye alone, is larger when both eyes are open than when one is closed, and farther that "there is a remarkable extension of each areal color field if a small white area is in the central region, and that this extension is almost the same if it is opposite" the one eye only, the other being under examination, though it is greatest where the white area is opposite both eyes.

These demonstrations of the fact that increased stimulation of the retina of either eye produces increase in the area of the periphery sensitive to color in the other eye, supports the theory of the close connection between the two higher visual centers, brought out also by the blindness of the periphery of both eyes on interference with the functions of only one higher center.

DISCUSSION.

Dr. C. W. KOLLOCK, Charleston—I have had some experience with these visual sensations. They have always been traced to indigestion. My first recollection of it was some years ago, when it affected one eye only, beginning with haziness of vision and then the waves appear. At times they appear in the outer fields of both eyes at the same time. The wave passes across the field of vision and goes away at the upper angle of the orbit. Once only did I notice them begin in the center of the field. The attacks last about fifteen minutes.

TREATMENT OF OPTIC NERVE ATROPHY BY MERCURIAL INUNCTIONS IN CON- JUNCTION WITH THE HOT BATHS OF THE HOT SPRINGS OF ARKANSAS.

Read in the Section on Ophthalmology at the Forty-seventh Annual Meeting of the American Medical Association held at Atlanta, Ga., May 5-8, 1896.

BY O. J. SHORT, M.D.

HOT SPRINGS, ARK.

It is not the object of this paper to consider the different pathologic conditions found in optic nerve atrophy, but to give my experience in the treatment

of this condition, no matter what the etiology, by the use of mercurial inunctions of ungt. hydrarg. U. S. P. in conjunction with the baths of the Hot Springs of Arkansas.

I have selected from my records five cases, each arising from a different cause. The mode of treatment pursued is practically the same in each, only varying the conditions and circumstances as indicated. For fear there may be some question as to the correct diagnoses I will state that in each case a like diagnosis was made by at least two oculists, men of unquestionable ability and in most instances members of this society.

Case 1.—Mr. C. C. O., aged 39, referred to me in May, 1893. On examination I found a gray atrophied condition of both optic nerves. I could find neither specific, spinal, nor any other cause to which we could attach the blame. In his history he stated that his mother became partially blind soon after giving him birth and remained so until her death. The opinion of the attending physicians was that the blindness was due to childbirth.

His vision, in right eye, could count fingers at two feet if in good light; in left eye, 20-200. He was ordered mercurial inunctions of 1 dram each night applied to the back and spine, and in addition to take a tub bath of ten minutes at a temperature of 98 degrees F.; at same time to observe all necessary hygienic precautions. On the tenth day symptoms of mercury were manifest in the mouth, on the gums and fauces. The vision was the same as at previous examination. At this visit he was ordered to use the inunctions every second night; increase the baths to 99 degrees F., and fifteen minutes duration, and in addition to apply a douche, 110 degrees F. to the spine and a vapor bath of three minutes. For the benefit of those not familiar with these baths, I will explain. The vapor bath is in a room three by four feet which is full of steam, 130 to 140 degrees F., continually rising through the open slat floor from a stream of hot water which flows underneath. On the twentieth day examination showed no sign of improvement in the vision of the right eye; while in the left eye it was 20-100. The inunctions were discontinued and he was placed upon kali iod. sat. sol. three times a day, commencing with 10 drops and increasing 3 drops each dose, until 60 drops were reached, which was continued until the thirtieth day, when examination revealed no improvement in vision of right eye; in left 20-100 minus. He was ordered to discontinue the kali iod. and to resume the inunctions alternate nights and to continue the baths as before, at the same time I began the use of hypodermic injections of strychn. nit. in the region of the seventh cervical vertebra, beginning with 1/40 grain and gradually increasing to 3/4 grain; toxic symptoms of the strychnia began to manifest themselves and its use was discontinued by gradually decreasing the dose. On the fiftieth day, still no change in vision of right eye; left eye 20-60. He was instructed to use the inunctions as before. On the sixtieth day no change in right eye; left eye 20-50. The patient decided to locate in Hot Springs. Although he has been under my care since and I have experimented with almost everything, there has been no change in his condition since the end of the third month.

Case 2.—A young lady aged 14. Referred to me June, 1894. Her father gave history of syphilis contracted five years before marriage and seven years before the birth of this daughter. The patient had always been healthy except the trouble with her eyes. Her sight began to fail about nine months previous to this consultation. My diagnosis was optic nerve atrophy due to inherited syphilis. I found the vision in right eye, 20-100; in left eye 20-150. She was promptly placed upon the mercurial inunctions, in conjunction with the hot baths, and at the end of the first week upon kali iod. in addition, and instructed to report to me once a week for examination. She improved rapidly from the first and at the end of the second month her vision in both eyes had returned to normal for the central, while the extreme outer fields seemed smoky or blurred. She was discharged and I received a letter some months later from her father saying that she was attending college and that her eyes gave her no trouble.

Case 3.—Mr. G. A. H., age 47, consulted me in November, 1894. He told me that he had gray atrophy of the optic nerve and said he was getting worse all the time, although he had been under the treatment of a prominent oculist since the previous May, and that he had taken mercury pills until he was salivated and potash by the pound. Examination showed atrophy of both optic nerves, with

vision in right eye 20-500; left eye 20-200. I could find no cause except that in March previous while on a railroad train, a steam heater burst, causing him to be unconscious for twelve hours and laying open his abdomen, in right iliac region, for four or five inches, allowing the intestines to protrude. This was properly attended to at the time and he made a good recovery and has felt no bad effects from it since. He noticed that his vision began to fail in May following, when he immediately consulted an oculist and had been under constant treatment since with the above results. He was ordered to begin the inunctions of mercury, 1 dram each night, applied to his back and spine, and to take a bath of 98 degrees F.; this was continued, with interruptions as the necessities of his physical condition demanded. The progress of the disease was almost immediately arrested and he claims he could see better at the end of the second week, but this I failed to verify by actual tests. At the end of the third month I could find no improvement in his vision, although he persisted in saying he could see better. He was then advised that he could gain nothing by remaining longer at the Springs at that time. He resumed his duties on the road until the following June, when he again consulted me; I found his vision the same as when he left. He was again put upon the inunctions and baths. In twenty-one days his vision improved to, for right eye 20-400; left eye 20-100. Although he has taken two courses of treatment since, there has been no change in his vision.

Case 4.—Mr. T. P. M., consulted me in June, 1895. He had been poisoned by lead fourteen months previously. His vision began to fail a month later, when he consulted an oculist who treated him for several months, and as he continued to grow worse he went to Denver where he was told he had optic nerve atrophy, due to lead poisoning and advised to visit the Hot Springs for the baths. On examination I found an atrophied condition of both optic nerves with vision in right eye 20-80; in left eye 20-300. He was instructed to begin the use of the mercurial inunctions, and in addition to take a bath of ten minutes at a temperature of 98 F. On the eleventh day toxic effects of mercury were noticed; the inunctions were ordered every third day. In addition he was put upon kali iod. beginning with 10 drops and increasing 3 drops each dose, and ordered vapor baths. This treatment was continued until he was taking 300 grains of kali iod. three times a day, when symptoms of iodism presented and its use was discontinued by decreasing 10 drops per dose. He was then ordered ten baths without the use of any medicine and dismissed at the end of the twelfth week. His vision began to improve during the second week; final examination showed right eye 20-40; left eye 20-200. He is still using the inunctions of mercury two or three times a week and writes me that his vision is improving.

Case 5.—Mr. W. T. R., referred to me in December, 1895. He gave history of syphilis contracted twenty years ago. His sight began to fail two years ago and he had been under the treatment of an oculist. On examination I found a dirty gray atrophied condition of both optic nerves, accompanied by Argyll Robertson pupil, ataxic gait, and knee reflexes in right knee entirely gone; in left greatly diminished. Vision of right eye 20-800; left eye only light perception. He was placed upon mercurial inunctions together with the hot baths. He insisted that he was getting worse both in vision and gait until the eighth day when he reported that he could walk much better and could tell the time of day by his watch, which I verified. The reflexes had slightly improved. His ataxia has continued to improve up to the present time, March 1, 1896, with a vision in right eye of 20-500; left eye no change.

My conclusions, based upon my observation of forty-one cases treated since residing in Hot Springs are:

1. That in those cases having only light perception, treatment is of no avail; but where there is any vision left, that the progress can be checked, and if the vision has not gone below 20-200, unless from traumatic origin, we may expect benefit.
2. That the mercurial inunctions in conjunction with the hot baths are always beneficial.
3. That unless the origin is from specific, blood or mineral poisons that the use of kali iod. is more often harmful than beneficial.
4. That strychnia, as a rule, is good only in cases of spinal origin.

Let us have a Department of Public Health!

THE INCREASE OF INSANITY AND TUBERCULOSIS IN THE SOUTHERN NEGRO SINCE 1860, AND ITS ALLIANCE, AND SOME OF THE SUGGESTED CAUSES.

Read in the Section on Neurology and Medical Jurisprudence, at the Forty-seventh Annual Meeting of the American Medical Association, held at Atlanta, Ga., May 5-8, 1896.

BY THEOPHILUS O. POWELL, M.D.

MILLEDGEVILLE, GA.

The negroes of Georgia, and I might say of the Southern States, up to 1860 enjoyed remarkable mental and physical health, and they were almost entirely exempt from certain diseases to which they are now not only very susceptible, but are dying much more rapidly from these maladies than the whites; namely, insanity and consumption.

While they may indicate much less susceptibility to miasmatic fevers, they are becoming more and more susceptible to them every year, and when we consider that heredity is one of the leading factors in the causation of insanity with the whites, and know that twenty-five or thirty years ago insanity and consumption in the negro could not have been attributed to hereditary predisposition save in a very few cases; couple this phenomenal increase in this race since 1860 with the inherited tendencies to these maladies, which we might say is just beginning to be manifested in full force, and when we remember that it is the insane and consumptive diathesis that widens and deepens by heredity from generation to generation, unless the laws of health are properly appreciated and conformed to, are we not justified in apprehending that the number of colored insane in the Southern States will soon be as large, if not larger, than the whites in proportion to the population?

The census of 1860 will show that there were forty-four insane negroes in the State of Georgia, and the majority had white blood in them; consumption in the full-blooded negro appeared to be just as rare at that time. Up to and during the war, the negroes were the principal nurses for the consumptives of the South. They washed the cuspidores, bedding and clothing of the consumptives, swept and dusted and in many cases slept in the rooms with them, and were more exposed to the tubercle bacillus than now as a general thing; still they resisted the disease. Why then this sudden and radical change which makes them so susceptible to both consumption and insanity? These 44 were of a population of 465,698, or one insane negro to every 10,584 of population; and there is no good reason from family pride, or fear that their future hopes would be blasted, to suppose that the census of 1860 was incorrect as so far as it relates to the negro insane.

The census of 1870 shows that there were 129 insane negroes in the State in a population of 545,142, or one colored insane to every 4,225. The census of 1880, colored population 725,133; insane, 411, or one to every 1,764. The census of 1890, population 858,815, insane 910, or one colored insane to every 943 of population.

The rapid increase of insanity and consumption in this race, is due to a combination of causes and conditions. This race has developed a highly insane, consumptive, syphilitic and alcoholic constitution which predisposes them to diseases which formerly they were free from. In this disturbed and unstable condition they seem to be totally unable to resist the