not perfectly identified. Extensive pleuritic effusion was discovered on the right side of the chest, with buttery lymph, and the same on the left side, but smaller in quantity. Thus, acute pleurisy may be considered as having had a large share in the fatal issue of the case. Several suppurative sources, or lobular phlebotic spots of pneumonia, were found on the surface of either lung, but the liver, spleen; heart, kidneys, &c., were healthy. Cases like this are well calculated to test M. Sédillot's opinions on pyæmia; we will, however, take another opportunity of touching on this subject, as this valuable case necessarily requires a somewhat extended report.

KING'S COLLEGE HOSPITAL.

Chorea.

(Under the care of Dr. Todd.)

Dr. Todd, in the Lumleian Lectures for 1849, says, in speaking of chorea—"No one now-a-days thinks of adopting for this malady any other treatment but one, eliminatory and corrective of the various secretions by the use of purgatives, and, at the same time, tonic and bracing to the nervous system by the free application of cold water to the surface, and also tending to improve the quality of the blood by a nourishing diet, by chalybeates, quinine, and various metallic tonics." We have seen this treatment so effective in the hands of Dr. Todd, that we deem it practically useful to bring forward some of the cases we noticed in our nosocomial peregrinations.

It will be interesting, before entering upon details, to inquire what are Dr. Todd's opinions on the pathology of chorea, and we cannot do better than to offer a few extracts from the lectures quoted above. "In chorea, the part affected primarily is the nervous system, but in all cases the muscular system participates in a depraved nutrition, which renders it more prone to sympathize with the irregular motions of the nervous system. . . . All evidence tends to show, that the part of the nervous system more especially affected in chorea is the *brain*." From various reasons, Dr. Todd believes that the corpora quadrigemina and the mesencephale are the parts principally engaged, one reason being that the nerves supplying the organs of sense take their rise in those localities. Among the circumstances which tend to show that the brain is more concerned than the spinal cord, the cessation of the convulsive movements in sleep may be mentioned.

Finally, Dr. Todd states—"1st, that chorea is a disease occurring at a time when the nutrition of the brain is passing, as it were, through a state of transition from that of infancy or very early childhood to that of the adult period, when that organ is particularly prone to suffer from a mental shock, or other causes of disturbance of the system; and more especially when the blood is in an unhealthy state, deficient in some of its staminal principles, or containing some morbid element. 2nd. The part of the brain most affected is the centre of emotion. 3rd. The nature of the cerebral affection is one of weakened nutrition. 4th. The disease is essentially one of depraved nutrition."

A glance at these pathological inquiries will show how logically the treatment has been deduced from the pathology of the disease, and we proceed to give two of the cases which we have seen treated at this hospital, availing ourselves of Mr. J. Dickinson's notes.

R. K—, aged four years and a half, a native of London, but for the last three years at nurse in Brentford. The patient was a very healthy child when he left home, nor suffered from worms, or had any illness whatsoever. The nurse wrote a short time ago to say that the child was ill, and when brought home he had convulsive movements of the face and limbs, continually throwing about his arms and legs, and at length falling asleep, apparently from exhaustion. On admission, the eyes and countenance were strangely distorted, and the convulsive movements followed by a vacant expression of countenance. While lying on his back, although at rest, both legs, and especially the left, are constantly tossing to and fro. When the patient stands up in bed, both legs appear much weakened, and he cannot walk at all without assistance, and even then his feet, instead of being placed flat on the ground, are turned outwards, his legs being drawn after him, as if paralyzed. His arms, too, are abruptly and convulsively drawn over his face and chest: this has caused chafing of the chin. When he attempts to speak he hesitates very much, and seems to express himself with much difficulty; when desired to put out his tongue, it is protruded with a peculiarly comical jerk.* The patient is always in a state of agitation, except during sleep, when he is perfectly tranquil. The sounds of the heart and lungs are natural.

Dr. Todd ordered a warm bath, low diet, with a slice of mutton, and a cold douche every morning. The latter ordeal was carefully applied, and the bowels kept regular with calomel and scammony. The improvement began, in a short time, to be well marked; most of the symptoms lost in intensity, with the exception of the jerking of the tongue. The legs remained uncontrollable, and the act of progression was more of the nature of stumbling than walking, the body being at the same time thrown forwards. Dr. Todd now ordered one grain of disulphate of quinine to be taken twice a day. This tonic medicine, being combined with the cold douche, had very satisfactory results. In about a dozen days the little patient began to walk more steadily, though he could not keep his equilibrium without assistance; the tongue was put out with much less jerking; and a few days afterwards he could walk by the beds of the ward without assistance. He improved so rapidly from that time, that he soon walked very fairly by himself, and put out his tongue much more steadily. About one month after admission he began to run about all day, from one ward to the other, and was very talkative and cheerful. A few days afterwards, the movements of the arms having quite ceased, and progression being perfectly steady, he was discharged cured.

The second case, from Mr. J. Dickinson's notes, is as follows:—

Edward J—, sixteen years of age, admitted under the care of Dr. Todd, has always enjoyed good health, with the exception of one or two of the diseases incidental to childhood.

[Dr. William Addison, in his work on Scrofula, lays much stress on the effect of these diseases of childhood upon the constitution; and it would appear but reasonable that attention should be paid to these first steps in the path of morbid phenomena, both in the history and treatment of disease.]

On one occasion he had a slight attack of rheumatism, and his legs were rather swelled. The family is healthy. The present illness began about a fortnight before admission, when he felt a sort of fidgeting in his right hand, could not keep it still, and then in a short time his whole body was affected. As a cause he assigns overwork; he looks, however, a strong and well-made lad. The disease is of so decided a nature, that the patient cannot remain still for ten seconds, and he throws himself, when lying in bed, into every imaginable attitude. The legs are the most affected, and by observing him for a short time, almost all the muscles of the body may be seen alternately in convulsive action. In walking he has an extremely awkward gait, rolls about somewhat like a sailor, and he protrudes his tongue, which is slightly furred, with a jerk. He speaks somewhat thick; hesitates very much in answering questions; and states that about six months ago he was very much *frightened*, when jumping about, by finding himself on the point of falling into a pit. This circumstance was, by his father, looked upon as a cause of his present complaint. General health good.

Dr. Todd adopted in this case the same line of treatment as in the preceding. The patient rapidly improved; so short a time as two days after admission he was considerably better, the left leg, which was the most affected, became tranquil, and the upper extremities were tolerably quiet. The occipito frontalis, and the muscles of the face generally were, however, much excited. On the fifth day he walked much more steadily, and did not roll about so much. The urine was loaded with lithates. On the seventh, the muscles, generally, were pretty quiet; the tongue was not thrust out with such a jerk as a day or two ago; his speech became clear and distinct, and he no longer hesitated in answering questions, as he formerly did.

On the tenth day after admission the improvement continued, the walking, though still awkward, was more steady, and the tongue was not protruded with such force. In short, all the symptoms gradually disappeared, and he was discharged, twenty-two days after admission, with a slight rotation of the head, and a little occasional fidgeting of the right arm. It will be perceived that the patients, the outline of whose cases has just been given, were of the male sex. Though the statistics of the disease gave a majority of females, it is useful to note how severely very young boys (four years and a half) may be affected.

THE COLLEGE ELECTIONS.—As the time approaches for the fellows to meet in the theatre of the Royal College of Surgeons, to elect and re-elect from this body into the council, we have obtained the list of the next dozen candidates, commencing from Mr. Bishop, the gentleman last elected, Messrs. J. G. Perry, G. Simpson, G. W. Mackmurdo, J. Kiernan, H. Heath, H. Russell, J. Harrison, T. Green, R. Hey, G. Gulliver, E. W. Tuson, and H. Clark. This list comprises six town and six provincial surgeons, the latter are of course ineligible, unless, like Mr. Hodgson, they remove to London.

* Dr. Todd looks upon this symptom as very important. He says, in the Lumleian Lectures for 1849:—"Of all the movements which accompany chorea, that of the tongue is the most characteristic; indeed, I would call it pathognomonic."