by the most famous therapists. As a matter of fact, these books retard scientific prescribing by keeping alive a lot of shotgun prescriptions which the therapeutic leaders of to-day would never think of employing. The science of therapeutics would take a long step forward if all such props should be cast into the eternal fires of oblivion.

But the outlook is not entirely dark. The very fact that we notice such methods shows that we know better. Manufacturing houses are finding out that the medical profession is beginning to do its own thinking on therapeutic problems, that absurd claims which manufacturers were once able to make for their preparations are no longer accepted, and that the old time-worn proverb, "Honesty is the best policy," is a maxim not altogether out of date.

On the part of the physicians it may be said that there is a large and ever-increasing number who, knowing the action of drugs, prescribe them scientifically, that is, simply and with a definite object in view and in sufficient doses to produce the physiologic effects characteristic of each.

It would be of the greatest aid to therapeutics if more physicians would not only keep careful clinical records of the effects of the drugs but would also publish from time to time critical comments on them. By work of this sort the practitioner would contribute greatly to our knowledge of drug action, and it is only by such mutual cooperation of the man at the bedside with the man in the laboratory that therapeutics will make the progress that it should.

THE ESSENTIAL CHARACTERISTICS OF THE TRUE PHYSICIAN

And now, to you who to-day come for the first time seeking to be followers of Æsculapius and true disciples of the healing art, in the name and on behalf of the faculty it is my privilege to bid you welcome. You have chosen a noble profession and one which before all others will give you an opportunity to serve your fellow men.

Medicine never offered a more inviting field for work than it does to-day; and, on the other hand, the demand for the highest type of manhood was never greater. For other types medicine has no place.

If you would know the kind of men that medicine demands, then listen to this quotation from the Vedas, which describes what the ancient Hindu required of the prospective student of medicine: "He should be of a mild disposition—noble by nature—not mean in acts—possessed of intelligence—free from pride—endowed with a large understanding—with a power of memory and judgment—having a liberal mind—devoted to truth—disposed to solitude, free from haughtiness—of a thoughtful disposition—not prone to wrath—endowed with purity of behavior and compassion for all—devotedly attached to the study of medicine—free from cupidity—without sloth—seeking the good of all creatures."

Such is, indeed, a high standard, but surely medicine of 1908 can not accept lower ideals than did the Hindu thousands of years ago. No standard can be too high, for, as Cicero said: "There is nothing in which men so approach the gods as in giving health to other men."

Health and Education.—C. G. Kerley states that as a general proposition there should be no choice between health and education; only that education is effective which is founded on and which secures good health.—*Archives of Pediatrics*.

Original Articles

COMBATING TROPICAL DISEASES IN THE PHILIPPINES BY SCIENTIFIC METHODS

R. P. STRONG, M.D.

Chief of Biological Laboratory, Bureau of Science
MANILA, P. I.

TROPICAL DYSENTERY

At the time of the American occupation of the Philippine Islands tropical dysentery was the most serious and common disease which attacked both Americans and Europeans in Manila and the vicinity. Hence it was this affection which first attracted the attention of the medical scientists who came here at this period to study the prevailing diseases of the country. Some idea of the importance of this malady may be derived from a consideration of the fact that during the year 1900 in the United States Army here there occurred four times as many deaths from dysentery as from any other single disease. Little knowledge of the cause and nature, and even of the intelligent treatment, of this affection could be obtained from the physicians resident at that time in the Philippines, since no accurate study had previously been made of it in these islands. It was, therefore, important that such an investigation as would result in the satisfactory control and prevention of the disease should be undertaken as speedily as possible.

An extensive research on this subject, carried on in the laboratories which had been established in Manila by the American government, finally revealed the fact that the so-called tropical dysentery of the Philippine Islands really included three distinct diseases and that the most satisfactory forms of treatment were widely different for these various affections; thus one form (amebic dysentery) in Manila was found to be due to a species of protozoa, an ameba; a second, with an acute onset (bacillary dysentery), to a species of bacteria known as *Bacillus dysenteriae*; and a third (catarrhal dysentery), a less common and a less serious form of dysentery, to the action of one of several bacteria which produced a catarrhal condition of the large intestine. The first type was found to be identical with the form usually known in certain portions of Africa under the term of "tropical dysentery," while the second was apparently the same as the prevailing form of intestinal flux occurring in Japan, and the third corresponded to the forms of catarrhal dysentery encountered more or less throughout the world.

It was, moreover, demonstrated that these three forms of dysentery could easily be distinguished from one another by laboratory methods. Thus the amebic form could always be recognized by a microscopic examination of the dejecta and by the presence of the specific parasite, the ameba, therein; while in the bacillary form the amebas were absent from the excreta; and the blood serum of the patient, when added to the *Bacillus dysenteriae* (the germ giving rise to this form of the disease), caused a clumping of the bacteria into groups and their ultimate destruction, a reaction not given by the blood serum of patients suffering with other forms of dysentery or with other diseases.

After these discoveries in the differentiation of the varieties of the disease its treatment was soon placed on an intelligent basis, and it was found that while the most effective treatment for the amebic form consisted in attacking the parasite by local applications (enemas)

the same therapy in the bacillary dysentery was not only not efficacious but very harmful, and that the best results in this variety of the disease were obtained by a general method of treatment and by the inoculation of antidyenteric serum. In the catarrhal type it was found that neither the local treatment nor the serum inoculations were to be recommended and that the disease generally yielded to much simpler therapy. It was further demonstrated that the amebic and bacillary dysenteries were usually acquired by drinking infected water and that the catarrhal form was, at least frequently, contracted in this way. Moreover, it was shown that the city water supply itself was practically always infected with amebas, and although it has not been definitely established that these same organisms are always capable of producing dysentery in man, nevertheless such water must be regarded as at least dangerous. So far it has been found quite impossible to destroy the amebas in the water supply by the introduction into the city reservoirs of chemical substances, such as copper sulphate, in safe amounts for drinking purposes, or to remove them by large filters, and for this reason, as well as for many others, the city water can not now, and probably never can be, recommended for drinking purposes unless it has been previously sterilized.

As a result of these discoveries both the mortality and the amount of dysentery among Americans and Europeans has become rapidly reduced during the past few years. The people have become educated to the necessity of avoiding the use of infected drinking water and of employing only boiled and distilled or imported aerated waters for drinking purposes. To-day bacillary dysentery, which formerly was epidemic here, and which frequently carried off its victims in from three to four days, is very rare in this city, and amebic dysentery is becoming less and less frequent each year among the more intelligent population. Physicians here have been taught to recognize these diseases in their very early stages, at a time when the symptoms may sometimes appear trivial, and then to institute the proper treatment. Therefore, the disease does not so often become chronic, and for this reason the amount of invaliding to the United States for dysentery has become much reduced. A decided advance in our knowledge of the amebas has been brought about by the demonstration in Manila of the ease with which they can be cultivated on solid media, and of the fact that they can be grown in the test-tube under certain conditions with almost the same facility as bacteria.

PLAGUE

In 1900 bubonic plague appeared in the city of Manila, the first case having been introduced from Hong-kong. For a time it was feared that the disease might gain a firm foothold as it had, for example, in India, and, therefore, strenuous methods were adopted for its control. Practically all vessels arriving from infected ports were fumigated at once and other means taken to prevent the escape of rats from them to the land. The most rigid isolation of the patients sick with the disease was carried out. Since rats were known to suffer from infection with the plague bacillus and to scatter this organism about by means of their excreta, and, in fact, to be the most important disseminators of the disease both in this manner and by means of fleas which they carry about on their bodies, the wholesale extermination of these rodents was undertaken. Squads of rat-

catchers, armed with rat traps, rat poison and every necessary implement, were assigned to the various districts of the city under the supervision of district sanitary inspectors. A bounty of 2½ cents gold was paid for each and every rat turned over to the health authorities and stations were established at convenient points throughout the city where they could be received. Each rat was tagged with the street and number of the building and lot from which it came and then dropped into a strong antiseptic solution and sent to the government laboratory, where it was subjected to a careful bacteriologic examination for plague. The premises where plague occurred, or where the rats infected with plague had been found, were subjected to a thorough disinfection and their owners compelled, under the direction of a sanitary engineer, to make necessary alterations. Cement ground floors were laid, double walls and ceilings affording refuge for rats were removed, defective plumbing was remedied, whitewash was liberally used for the destruction of fleas, and in general everything was done that could render the buildings where plague had occurred safe for human occupancy. Wherever these changes were not practicable, houses, or even districts, were frequently destroyed by fire.

The Board of Health in Manila considered, however, that by far the most effective measure in combating the epidemic of plague consisted in the immunization of the people living in the badly infected districts by means of inoculation with the prophylactic against the plague. All Chinese arriving from infected ports in China, and all those individuals living in a house where a case of plague occurred, were also inoculated with the prophylactic or with antiplague serum. To quote from the report of the director of health at this time in regard to the preventive inoculation against plague carried on in Manila: "The method has given results almost beyond expectation."

Partly for this reason and partly from the character of the disease, it seemed hopeful that a considerably greater success might be obtained against this malady by improved methods of protective inoculation; therefore particular attention was paid to the investigation of this subject in the government laboratories in Manila and a method of vaccination has been finally obtained which will probably reduce the mortality of plague among the vaccinated to under 20 per cent. usually, whereas it is well known that the usual mortality of plague is in the neighborhood of 80 per cent.

At present there is no plague in the Philippine Islands nor has a case occurred in them for over a year. It is hoped, however, that the method of vaccination against this disease will be employed in India, where the plague has raged for the past ten years in spite of all hygienic measures employed there and during which time the mortality has amounted to over 5,000,000.

CHOLERA

The first appearance of cholera in the Philippine Islands under American occupation occurred in March, 1902, and measures were at once undertaken to prevent the occurrence of an epidemic. Unfortunately, it was impossible to convince the majority of native physicians and the population in general that cholera had really invaded the islands and that a severe epidemic was imminent, notwithstanding the fact that the first cases found in the city had in the laboratory been carefully identified bacteriologically as those of cholera. For

weeks, however, the presence of cholera was denied, the Filipinos refusing to believe in its existence because the daily deaths did not reach up into the thousands as they had done in the epidemic of this disease which occurred here prior to American occupation. Therefore, as was to be expected, the rigorous measures enforced by the board of health produced bitter opposition. The minds of the common people were poisoned by tales of abuses and even of murder in the cholera hospitals. The rumor was also circulated that the cholera was due to the Americans having poisoned the wells and that the quarantine measures instituted were evidences of revenge on their part. These absurd tales gained credence and proved a very serious factor in the control of the situation, since they led to the concealment of the sick, the escape of those who had been in close contact with cholera patients, and the throwing of the dead cholera bodies into the esteros (canals) throughout the city and into the Pasig River (the waters of the latter were frequently used for drinking purposes by the natives). In fact, for the first few weeks of the epidemic there was a general disposition on the part of the population to conceal every case of cholera which occurred. As a result, in spite of all measures of control, the disease spread throughout the city. The following measures for its suppression were, however, carried on and served to keep the epidemic under control:

A cholera hospital was established and, with a few exceptions, all definite and suspected cases were speedily removed to it for treatment. All cases of the disease were isolated. The city was divided into districts and each district was inspected during night and day for the purpose of detecting and diagnosing cases of the disease. All wells throughout the city were closed and distilled water was distributed for drinking purposes. A rigorous quarantine was placed around the city, and the Mariquina River, from which the city water supply was obtained, was carefully guarded by a patrol of regular troops in order to prevent it becoming infected. Certain districts, which were thoroughly infected with the germ of cholera and which could not be disinfected properly owing to the character of the dwellings, were destroyed by fire; notable among these was the so-called Farola District. The natives were reimbursed by the government for the loss of their houses and their property.

In order to aid the board of health in most of this work, measures for the early bacteriologic examination and diagnosis of the disease by means of cultures and immune sera were undertaken. An assistant was on duty day and night at the laboratory for the purpose of performing such diagnoses. The city water supply was examined from time to time throughout the epidemic for the cholera germ. This water was never found infected with this organism. Had it been so, the cases of cholera in Manila would have reached into the thousands daily. During the epidemic the highest death rate reached in Manila was ninety-one per diem.

Since it became evident early in the epidemic from bacteriologic examination that the city water supply was not infected with cholera, investigations were undertaken in the laboratory to find out by what means the disease was being spread. Visits were made from house to house in the badly infected cholera districts and cultures were taken from various foods, fruits, linen and articles of clothing and from various samples of water found in these houses. These studies were continued

throughout the year and as a result of them it was determined that one of the chief means of the dissemination of the disease in Manila was through infected uncooked foods. As a result of these studies a number of uncooked foods and fruits were forbidden for sale in the markets. Flies were also shown to have played a part in the infection of certain foods where unsterilized excreta of cholera patients were exposed in the neighborhood. Certain wells and canals in the city were found to be infected with the cholera germ, and these also played a part in the spread of the disease; but, as mentioned, the general city water supply has never yet been found to be infected with the cholera organism.

Since it was recognized that individuals who had recovered from cholera might carry about with them the germ of the disease in their intestines for several weeks after convalescence and thus serve as disseminators of the infection, an attempt was made to keep each patient at the cholera hospital until his excreta were shown to be bacteriologically free from the cholera germ. It was later found, however, that some individuals who had only very mild symptoms of cholera, and who had not been taken to the hospital, also had cholera spirilla in their dejecta and thus they were able to act in spreading the infection.

Partially on this account, since such individuals must frequently escape quarantine, and partially as a general measure in the prevention of the disease, the subject of immunization in human beings against cholera was studied in the Manila government laboratories, and after extensive investigations a method of protective inoculation against cholera was devised which is perfectly harmless and which is not objected to by the natives. This method has been given an extensive trial in some of the smaller towns of the Philippines where cholera has been epidemic, and in Manila, and in all of these places large numbers of men, women and children volunteered for inoculation.

Although cholera has on several occasions since 1902 been banished from Manila and, indeed, supposedly from the entire Philippine Islands, it has always reappeared therein and has probably become, for the present at least, endemic in the islands. While the present hygienic measures are enforced, however, there is little danger of a large epidemic. It seems not unlikely that by means of protective inoculation we have a valuable adjunct in ridding a community of this disease, and it may be that we shall here finally come to resort to this method more generally for the extermination of the malady. In certain provinces of the islands statistics already show that the number of cases of cholera which have developed in the inoculated is only one-sixth of that which has occurred among the uninoculated.

MALARIA

In regard to malaria, it has been shown that nearly 50 per cent. of the native children in certain portions of the islands suffer from a latent form of this disease; that is, while many of them may not appear very ill to the casual observer, they are carrying about with them malarial parasites in their blood and furnish a constant source for the transmission of the disease to other individuals through the medium of certain species of the anopheles mosquito. Presenting few or only very indefinite symptoms, these children frequently remain untreated for the malarial infection. It is a well-recognized fact that the greatest source of danger to the white man

in a malarial locality lies in these latent infections in the native, and particularly in the native children. Studies in the Philippines have demonstrated the importance of discovering these cases by means of a routine microscopic examination of the blood, and of the treatment of the patient and destruction of the parasites by means of quinin. Such measures play an important rôle in the suppression of malaria in certain portions of the islands.

The amount of anemia and general debility produced by malaria in the native population of the Philippines has been greatly underestimated. In some districts over 50 per cent. of the adults show either active or latent infection. Malaria is uncommon among white people in the islands, however, particularly in the large cities.

DENGUE FEVER

Studies in dengue fever have been recently made here which seem to show that this disease is perhaps conveyed by means of the bite of the mosquito *Culex fatigans*, which also transmits the parasite (a blood filaria) which gives rise to the affection known as elephantiasis. These investigations have also shown that the organism causing dengue fever is probably too small to be observed by the microscope (that is, it is ultramicroscopic), and that it is present in the circulating blood of the patient suffering with the disease.

INFECTION WITH INTESTINAL WORMS

An investigation has recently been undertaken in regard to the prevalence of animal parasites in the native population and their possible influence on the public health. In this study the dejecta of 4,106 prisoners were investigated, in all 15,000 microscopic examinations being made. It was found that 84 per cent. of these prisoners were infected with one or more animal parasites.

The most usual infection was with the common whipworm (*trichuris*), which was present in 59 per cent. Fifty-two per cent. were infected with hookworms (*uncinaria*), though very rarely were these infections severe ones. Indeed, in the Philippines this parasite does not play the important rôle that it does in Porto Rico, Egypt and several other countries in the production of a grave anemia. The great majority of those infected here show only mild infections, practically having no symptoms of disease and no visible anemia. The dejecta of 23 per cent. of the prisoners contained amebas and those of 26 per cent. round worms (*ascarides*). The numerous other parasites encountered were found only in a comparatively small percentage of those examined.

The question has arisen as to what rôle these infections play in regard to public health. Attention has been called to the fact that if these percentages of infection obtained from the examination of the 4,106 prisoners were applied to the total native population of the islands it would appear that no less than five million of the inhabitants of the Philippine Islands were infected with intestinal worms.

It has been argued that since the systematic treatment of the prisoners for intestinal worms was begun and vigorously carried out the death rate of the prison where these examinations were performed fell from 75 per 1,000 to 13 per 1,000 per annum. It appears, however, that this decrease in the death rate was very largely due to the gradual improvement in the general

sanitary conditions in the prison, which lead to a very great reduction in the number of cases of beriberi present, many of which had been formerly unrecognized, and to the fact that the new prisoners arriving at the prison were at once surrounded by better hygienic conditions than the older prisoners had formerly been. While there is practically no evidence to show that the moderate infections with whipworms and round worms exercise a deleterious effect on the health of the individual, the infections with hookworms and with amebas must be regarded in a different light. Both of these parasites are pathogenic or may come to have decided pathologic importance in the individual infected. There can be but little doubt that the patients in a number of the cases in which there were amebas in the dejecta would sooner or later develop amebic dysentery, and it is very probable that a certain percentage of these would succumb to the infection if not treated. In this way, by the detection of these cases of amebic dysentery, and by instituting proper treatment, the mortality among the number infected would be reduced. As has been pointed out, the cases of hookworm infection were so mild that the hookworms themselves probably had but little influence on the death rate. However, in the more severe infections, both with hookworms and amebas, these parasites undoubtedly weakened the individual and rendered him more susceptible to the attack of other infectious diseases.

Since the source of all these intestinal worms is the dejecta of some individual infected, the most important factor in the extermination of this infection lies in the proper disposition of human excreta. The most satisfactory means for such disposal are now being investigated.

To obtain a working system, practical, economical and adapted to the conditions throughout the islands, is a special problem which will require considerable further study. When such a system is perfected, however, it will remove one of the most dangerous channels, not only for the dissemination of animal parasites, but for the spread of other intestinal infectious diseases of bacterial origin.

SMALLPOX

In relation to smallpox, attention must be called to the fact that since the beginning of systematic vaccinations in the islands over 3,515,000 people have been vaccinated against this disease with virus prepared in the government laboratory at Manila. A cessation of deaths from smallpox has followed in the provinces in which the people have been thoroughly vaccinated. In provinces where there were formerly 6,000 cases of the disease annually not one death has been reported from this disease for the past year. No case of loss of life on account of vaccination has occurred, nor have any limbs been sacrificed, nor has there been a case of very severe infection resulting from vaccination reported. The disease has been thoroughly kept under control by vaccination alone.

GENERAL HEALTH OF MANILA

After nearly ten years' residence in the Philippines I think that I am justified in saying that Manila to-day can certainly be regarded as a very healthy city for those who exercise proper hygienic precautions in regard to their mode of living. The death rate here for the past year among American residents has been only 5.59 and among Spaniards but 15.84 per 1,000.