This article was downloaded by: [UZH Hauptbibliothek / Zentralbibliothek Zürich]
On: 26 December 2014, At: 08:26
Publisher: Taylor & Francis
Informa Ltd Registered in England and Wales Registered Number: 1072954 Registered office: Mortimer House, 37-41 Mortimer Street, London W1T 3JH, UK

Annals and Magazine of Natural History: Series 6

Publication details, including instructions for authors and subscription information:
<http://www.tandfonline.com/loi/tnah12>

XXII.—On Strauch's Triton longipes

G. A. Boulenger

Published online: 02 Oct 2009.

To cite this article: G. A. Boulenger (1892) XXII.—On Strauch's Triton longipes, *Annals and Magazine of Natural History: Series 6*, 9:50, 144-145, DOI: [10.1080/00222939208677290](https://doi.org/10.1080/00222939208677290)

To link to this article: <http://dx.doi.org/10.1080/00222939208677290>

PLEASE SCROLL DOWN FOR ARTICLE

Taylor & Francis makes every effort to ensure the accuracy of all the information (the "Content") contained in the publications on our platform. However, Taylor & Francis, our agents, and our licensors make no representations or warranties whatsoever as to the accuracy, completeness, or suitability for any purpose of the Content. Any opinions and views expressed in this publication are the opinions and views of the authors, and are not the views of or endorsed by Taylor & Francis. The accuracy of the Content should not be relied upon and should be independently verified with primary sources of information. Taylor and Francis shall not be

liable for any losses, actions, claims, proceedings, demands, costs, expenses, damages, and other liabilities whatsoever or howsoever caused arising directly or indirectly in connection with, in relation to or arising out of the use of the Content.

This article may be used for research, teaching, and private study purposes. Any substantial or systematic reproduction, redistribution, reselling, loan, sub-licensing, systematic supply, or distribution in any form to anyone is expressly forbidden. Terms & Conditions of access and use can be found at <http://www.tandfonline.com/page/terms-and-conditions>

Dimensions.

	<i>T. microlepis.</i>		<i>T. Blythii.</i>	
	Ad. millim.	Yg. millim.	Ad. millim.	Yg. millim.
Total length.....	150	80	163	78
" " (without caudal)	127	66	140	64
Depth of body	64	32	59	29
Length of head	44	26	48	26
Diameter of eye	12	8	13	8
Length of 1st dorsal spine	9	5	9	4½
" 2nd " "	17	10	17	9
" 3rd " "	21	14	25	?
" 4th " "	26	16	26	13
" 5th " "	25	13	25	13
" 1st anal " "	13	6	13	6
" 2nd " "	17	9	19	10
" 3rd " "	18	11	25	14

XXII.—On *Strauch's* Triton longipes.

By G. A. BOULENGER.

IN my revision of the newts, published in 1882*, I expressed the opinion that *Strauch's* *Triton longipes* † from Astrabad, N. Persia, should probably be considered a variety of *Molge cristata*, agreeing with var. *Karelinii*, *Strauch*, except in the more elongate digits. This view was endorsed shortly afterwards by *Camerano* ‡, who referred some Italian specimens (from the Gran Sasso) to subsp. *longipes*, not, however, without some hesitation. I am now in a position, thanks to the kindness of Dr. F. S. Monticelli, who sent me numerous specimens of *M. cristata* from near Naples, to state that *T. longipes* represents merely an individual variation of *Molge cristata*, var. *Karelinii*. Among the Naples specimens there is one, a male *post nuptias*, which is in every respect referable to *T. longipes*; its digits are extremely slender and elongate, all the fingers except the first extending beyond the snout when the fore limb is stretched forwards, and the longest toe reaching the elbow when the limbs are pressed against the body. As there exists in the same locality every passage between such a specimen and a normal *M. cristata*,

* Cat. Batr. Grad. s. Caud. 1882.

† Rev. Salam. Gatt., Mém. Ac. St. Pétersb. xvi. no. 4, 1870, p. 44.

‡ P. Z. S. 1884, p. 423, and Mon. Anf. Urod. Ital., Mem. Acc. Tor. (2) xxxvi. 1884.

var. *Karelinii*, it is clear that the form *longipes* cannot be upheld even as a subspecies. The measurements are here given of the specimen referable to *T. longipes* (a) and of a normal *M. cristata*, var. *Karelinii* (b), both males from Naples.

	a. millim.	b. millim.
Total length	108	108
From snout to cloaca	60	56
Head (skull)	14	14
Width of head	11	11
Fore limb	25	22
Third finger	8	6
Hind limb	28	22
Third toe	10	7
Tail	48	52

It must be borne in mind that the typical form also varies considerably as regards the elongation of the digits, and that Strauch's statement as to the proportions of the limbs in *M. cristata* applies only to females, which have the digits much shorter than in the males. It is quite clear, from the measurements given on p. 46 of his memoir, that the specimen from the Gov. Kursk, with which he compares the (male) type specimen of *T. longipes*, is a female; it need hardly be added that such a course, in a group of animals in which the secondary sexual characters are so highly developed, can only convey a fallacious impression, especially if no mention is made of the sexes of the specimens the measurements of which are tabulated.

XXIII.—*Note on the Gibbon of the Island of Hainan*
(*Hylobates hainanus*, *sp. n.*). By OLDFIELD THOMAS.

IN his paper on the mammals of Hainan * Mr. Robert Swinhoe has given an account, mainly compiled from native authorities, of the Gibbon which inhabits that island; but no specimen appears hitherto to have been brought to Europe. The animal was referred by Mr. Swinhoe, with some doubt, to *Hylobates pileatus*, Gray, the Cambodian species; but as he was unable to obtain a specimen, this reference has never been confirmed.

At length a specimen of this Gibbon has been presented to the British Museum by Mr. W. T. Lay, to whom it was

* Proc. Zool. Soc. 1870, p. 224.