sented only in the wall of the bladder, and does not enter the cavity, so the danger of its forming nuclei for other calculi is entirely avoided.

Kocher has performed this operation six times during the winter semester. All the patients recovered, without rise of temperature, in eight days, and were sent home in two weeks' time. This illustrates the simplicity of operation at this early period as soon as the diagnosis is established and before complications have rendered the intervention more hazardous.

He does not believe that any form of medical treatment can remove these calculi or prevent the patient from having recurrent attacks until the calculi are removed. Recurrence is not probable after all the calculi are removed, but the patient should be put upon a regulated diet, with the end in view of preventing the formation of other calculi.

PEDIATRICS.

UNDER THE CHARGE OF

LOUIS STARR, M.D.,
OF PHILADELPHIA,

AND

THOMPSON S. WESTCOTT, M.D.,
OF PHILADELPHIA.

Pulmonary Affections Consecutive to Infectious Diseases and Gastro-enteritis.—H. SPIEGELBERG (*Archiv für Kinderheilkunde*, 1899, Band xxvii., S. 367) discusses the question of the bronchopneumonias of intestinal origin, which has been sustained by Czerny in Germany, and reaches the conclusion that these bronchopneumonias are to be attributed to secondary infection of bronchial origin.

The researches were made upon 111 children, 90 of whom were one year old, 18 two years old, and 2 three years old. All had succumbed in the course of various infectious diseases, and of the number there were 29 cases of acute, subacute, or chronic gastro-enteritis, with terminal bronchopneumonia.

In all the cases associated with gastro-enteritis the lungs were found to be more or less congested, succulent, and presenting foci of lobular infiltration of variable size. Microscopically these foci were composed ordinarily of bronchioles and alveoli filled with an exudate containing round and epithelial cells. In the severer cases cellular infiltration extended into the alveolar septa and the interstitial tissue, giving a homogeneous appearance to the affected areas. In some cases in which the infectious agents were particularly virulent a superficial necrosis of the mucosa was found.

The mucous lining of the larger bronchi was congested, while that of the smaller branches was covered with purulent exudate, almost completely

tion of the cord showed the presence in pure culture of a streptococcus analogous to that observed in the first case.

The same organism, therefore, determined in one case a serous inflammation, in the other a seropurulent one. In discussing the reason for this difference of effect the authors think that a difference in the virulence of the germ in the two cases can scarcely be invoked, since in the seropurulent case the organism was innocuous for animals; neither could the duration of the disease be considered important, since the serous inflammation evolved more slowly than the other. The hypothesis of a special resistance or a peculiar predisposition of the meninges alone remains, but these are factors which are not clearly understood.

The organism presented some interesting peculiarities. It grew poorly at first upon gelose, giving very minute punctiform colonies, resembling those of the pneumococcus. After several removes it acquired the typical characteristics of a streptococcus, giving in liquid media a deposit formed of large clumps. Morphologically it had at first the form of a diplococcus; later there were long chains when cultivated in the blood-serum of the rabbit. The organism possessed no capsule taking stain. It was virulent for neither rabbits nor mice.

THERAPEUTICS.

UNDER THE CHARGE OF

REYNOLD W. WILCOX, M.D., LL.D.,

PROFESSOR OF MEDICINE AND THERAPEUTICS AT THE NEW YORK POST-GRADUATE MEDICAL SCHOOL AND HOSPITAL; VISITING PHYSICIAN TO ST. MARK'S HOSPITAL.

Atmokaussis.—DR. A. VON GUÉRARD reports his results from the use of this method in twenty-one instances. For uterine hemorrhage, especially at the climacteric, this method is serviceable. The results in putrid abortion, endometritis, carcinoma, and dysmenorrhœa are uncertain. It was successfully used in a bleeder whose mother and sister had died from hemorrhage. The application was made at 239° F. for thirty seconds. Of these twenty-one patients seventeen suffered pain, in five of whom it was excessive, and this symptom was accompanied by fever. This method may be used successfully when contraindications exist to others.—*Therapeutische Monatshefte*, 1900, Heft 3, S. 154.

Soluble Silver.—DR. A. DWARETSKY reports the histories of two patients suffering from phlegmon of the hand, with septicæmia; one of the same condition of the foot, one of furunculosis with septic infection, and one of chronic osteomyelitis of the femur, with chronic septicæmia. In all instances with the ointment was practised, and this resulted in benefit both to the local sepsis and the general septic conditions. These instances of its use fully confirm the claims of Credé and others.—*Therapeutische Monatshefte*, 1900, Heft 3, S. 145.