Scar-contraction can never injuriously diminish the lumen of the tube, because the scar which encircles the ureter after union by this method is equal in length to twice the extent of the incision in the side of the lower ureteral stump.

Transverse suture is difficult and seems likely to be followed by stricture, unless an effort is made, by splitting the ends, to increase the lumen. Hochenegg is reported to have made use of this method. Bovée reports having successfully employed direct suture after having divided the ends obliquely.

Paggi is said to have been the first to propose end-in-end anastomosis, but I have been unable to find a description of his method or the technique of the method as carried out by others. The method of end-in-end anastomosis as carried out in this case seems to have all the advantages of Van Hook's method, with one or two additional: 1. It is simple and more quickly done. 2. There is less waste of substance, and it would be available when the Van Hook method would produce too great tension, or where the distal fragment is so short that union by Van Hook's method could not be made. It is much quicker, simpler and surer than by transverse suture or oblique end-to-end suture as employed by Bovée.

When division of the ureter occurs, the following would seem to be the rational mode of choosing the method of repair: 1. When possible to perform uretero-ureteral anastomosis this is the preferable procedure: end-in-end anastomosis, as employed in this case, seeming to me simpler and better. 2. When the distal portion is too short for uretero-ureteral anastomosis, implantation into the bladder should be performed. 3. When there is too much loss of substance to permit uretero-ureteral anastomosis, and the proximal end can not be brought down to the bladder even with the assistance of a diverticulum of the bladder, as devised by Van Hook, the procedure with the least objection is probably implantation into the colon.

EPISPADIC EXSTROPHY OF THE BLADDER COMPLETE.*

AP MORGAN VANCE, M.D.

LOUISVILLE, KY.

Dr. Rudolph Matas,¹ at the last meeting of the Association, gave such a complete résumé of the subject that it would be simply repetition to now enter into a general discussion of the topic—and in his conclusions favored Maydl's operation, holding that the autoplasmic operations could be little more than palliative, and never satisfactory. His paper decided me to report a case and exhibit the patient, showing the result of autoplasmic work. I do not know how original the procedure to be described may be, though I have been unable to find a report of such a one.

Mr. J. McE., aged 17 years, was referred to me by Dr. J. M. Mathews, on Oct. 26, 1897. He had come to Louisville to obtain a urinal and, if possible, get relief from the miserable condition of complete bladder exstrophy. Fig. 1 shows the condition before operation.

He was fairly well developed for his age, but his life was rendered unendurable as the result of excoriations from the urine and the contact of the clothing with the bulging posterior bladder wall, where ulceration with the deposit of calcareous material around the edges rendered the local condition horrible in the extreme. On Nov. 1,

1897, the first operation was done. This was much too soon, as five days was not sufficient to get the parts into proper condition, but it was done at this time because his family physician, Dr. Cowen, of Girard, Ill., who came with him, wished to return.

The epispadic penis was very short and drawn back into a sulcus at the lower part of the opening. The posterior bladder wall, with discharging ureters, presented a convex surface far in front of the abdominal plane, about the size of the half of a baseball or larger—when recumbent or in the Trendelenburg position a concave sulcus of corresponding dimensions presented.

The first step in the operation, as illustrated by Fig. 2, was to score as deeply as possible in a triangular manner.

FIGURE 1.

wide out into the iliac regions, then with strong forceps to drag upward and inward, still further loosening the tissues thus freed, bringing the mucous membrane along with the skin. When this loosening and stretching had been done sufficiently to allow the separated pudendal structure to be brought together in the middle line, the edges down to the end of the penis were vivified by splitting and brought together by two sets of sutures: one line of buried chromicized catgut, No. 1, which inverted the inner or mucous membrane side, and another of interrupted silkworm-gut which everted the outer or skin side. Over this a dressing of collodion and cotton was applied. The button-hole-like opening left above was used to drain the urine away, the patient being required to maintain the Trendelenburg position. No

*Presented to the Section on Surgery and Anatomy, at the Fifty-first Annual Meeting of the American Medical Association, held at Atlantic City, N. J., June 5-8, 1900.

1. THE JOURNAL, xxxiii, p. 260.

good resulted from this operation, except probably the piling up of tissue to be used to better advantage at the next attempt.

On Jan. 16, 1898, all the parts being healed and the resulting infiltration having disappeared, the above-described flap-splitting and suturing was repeated, the only difference being that I had less tension to combat and was gratified on removing the sutures on the tenth day to find that the penis was united and a fairly good bridge of tissue at the upper part of the pudendal portion also held together. Fig. 3 shows this result.

On July 16, six months later, the third operation was done. It consisted in the closure of the transverse opening at the upper part of the original open space. This

The new bladder has now been performing its function six months and is in a perfectly healthy condition. The penis has developed quite a good deal. The young man has been able to fulfill the duties of a salesman in a store, is able to keep himself dry and, in fact, is altogether a new man in every way. While recumbent about four ounces of urine will accumulate, and this is voided when he arises and can be passed out with some degree of force by voluntary effort. When upright the urine is received into an ordinary urinal. Fig. 4 shows the final result.

I had at the same time a second similar case, but the child was only 7 years of age. The result promised to be quite as good and was almost complete. He was allowed

FIGURE 2.

was done by freshening the edges of the bridge of tissue and the line of skin above by splitting, the same method of suturing being used, except that both lines of sutures were buried and both were chromicized catgut. This healed from end to end, leaving a round hole about the size of a silver quarter still uncovered between the lower edge of the bridge of tissue above referred to and the root of the penis.

By this time the young man's improved condition encouraged great hopes of final success, but four other attempts at closure were made before I succeeded, and another year was required. These last procedures were done by splitting the edges of the opening and suturing, and each trial was followed by a decrease in the uncovered area. The last operation, done just two years after the first, finally finished the case.

FIGURE 3.

to return home on a visit, and has never returned for the finishing touches, much to my disappointment.

In this kind of work much patience is required both on the part of the patient and the surgeon, and many difficulties are to be overcome, principal among which are the septic surroundings, the difficulty of drawing away the urine, and the constant lack of rest of the parts due to the tendency to erections. Notwithstanding all the difficulties, the benefits justify one in the efforts at autoplasmic work.

Maydl's operation, in the light of future experience, may prove better, but I am inclined to believe that in the hands of the rank and file of surgeons it is a very grave procedure, to say nothing of the possible future complications that may arise in the way of disease of the kidneys due to septic invasion from below.

DISCUSSION.

DR. DEFOREST WILLARD, Philadelphia—I desire to congratulate Dr. Vance on the excellent results obtained in this case, which is certainly as satisfactory as I have ever seen. In spite, however, of the good results obtained in some of these cases, I think all of us will agree that operation after operation frequently fails to secure success. This class of patients are so disgusting to themselves and to others that they deserve the most careful attention, and none require more surgical skill. Nearly all of the external plastic operations are faulty from the fact that we turn our flaps with skin surface inward. Concretions on the inner surface of the bladder follow from deposition of urinary salts on the hairs, and subsequent opening of the bladder is sometimes necessary. This difficulty Dr. Vance wisely avoided by leaving the skin face of the flaps external.

FIGURE 4.

On the whole, I am convinced that the formation of an intestinal reservoir is better. To obtain this it is preferable to abandon the rectum and go higher up above the sigmoid flexure, since this flexure is a valve and helps to retain the urine. Such a reservoir permits the patient to void urine and feces at stated intervals, and seems at present the best result obtainable under the circumstances. The insertion of the ureter into the bowel is so liable to be followed by kidney infection that it is wiser to remove, at the same time, a portion of the bladder wall, in order to obtain the normal valve of the ureter, since we can never secure a similar valve after division of the tube.

DR. HOWARD A. KELLY, Baltimore—The condition of these exstrophic patients is indeed a sad one. The only operation we can do with certainty of immediate success, is to put the ureter with the neighboring portion of the bladder into the rectum, according to Maydl. The condition of this patient is, how-

ever, remarkable, and does Dr. Vance great credit. I have a case under my care somewhat similar, and I have done a number of these skin-flap operations and tested their utility thoroughly, and have come to reject all operations that turn the skin in with the hair. I broke up the sacro-iliac joint to approximate the symphysis (Trendelenburg), but still it failed; and I then tried what has given me the best result, that is, skinning out the muscular bladder from the peritoneum (Sonnenburg). I dissected it loose from the underlying surface and brought the lax muscularis together. I was thus able to secure a reservoir of considerable size, which has held fairly well so far. I was also able at the same time to form a good penis. I could not have done Dr. Vance's operation in this case, as there were such extensive areas of scar tissue. Skinning up the bladder should be given a more careful trial.

DR. W. J. MEANS, Columbus, Ohio—At the Columbus meeting of the Association I exhibited a case of exstrophy of the bladder, on which I had operated a few weeks previous, similar to Dr. Vance's. The result obtained by the Doctor in this case is certainly excellent, and he is to be congratulated. The operation in my own case was successful in covering the bladder and making a pouch, but a failure in relieving the epispadias. I operated again two months ago for the latter condition with good success. If the patient has the fortitude to persevere, I am satisfied I can still further improve his condition. It was my desire to perform the Maydl operation, but when I mentioned the dangers attending it, his friends objected. It is my opinion that of all the methods tried for the relief of exstrophy of the bladder, none equal that of Maydl.

I had the pleasure of meeting Dr. Allen a few weeks ago. He informed me that the patient on whom he had performed the Maydl operation and presented to this section, was still in perfect health.

The method adopted by Dr. Vance in covering the exposed bladder is certainly ingenious, but I doubt if success follows in many cases. In my case I used the double skin flap. The patient has not had the difficulties from lime deposits that usually follow when the skin is used to form the internal layer of the covering. These plastic operations are, at best, only palliative. A natural reservoir for the urine, that can be emptied at the will of the person, can not be formed by artificial work. The only natural sphincter muscle contiguous to the ureters is the sphincter ani; but to utilize this the urine must be carried into the bowel. The longevity of patients where the rectum is used as a reservoir is a matter of the future to determine.

DR. MAXMILIAN HERZOG, Chicago—Dr. Jacob Frank, of Chicago, has recently devised an operation for exstrophy of the bladder. His operation consists of an anastomosis by the aid of his absorbable bone coupler. Frank has made his operation on a number of dogs, and I have myself made the post-mortem examination and histologic and bacteriologic examination on the animals operated on. Dr. Halstead of Chicago has had a chance to perform Frank's operation on the human being. When operating for exstrophy of the bladder it is undoubtedly very desirable to procure for the urine a receptacle which can be voided at will at certain intervals. The great trouble in making part of the intestines a receptacle for the urine is that the procedure as a rule leads to infection of the kidneys from the gut. Several of Dr. Frank's dogs, however, remained free from infection of the kidneys so that it appears that anastomoses of the bladder and rectum may be made, even in dogs, where the conditions are not as favorable as in the human being, without a subsequent infection of the kidneys.

DR. J. RILEY EASTMAN, Indianapolis—It may interest those members of the section who were present at the Columbus Meeting to know that the case of bladder exstrophy which I reported at that time is still doing well. One of the kidneys in this case was much disorganized and was removed. I removed the bladder mucosa and implanted the end of the ureter from the remaining right kidney at the tip of the clubbed epispadic penis, the penis having been dorsally split in the median sagittal line for reception of the ureter. All the work of secretion of urine has therefore devolved upon the one kidney,

but the boy has enjoyed good health at all times since the operation (eighteen months.) He wears a specially-constructed silver and rubber urinal and is apparently unhampered in work and play by his deformity.

DR. F. G. CONNELL, Chicago—During the last year I have had occasion to look up the literature on exstrophy of the bladder, in connection with some experimental work along this line upon the lower animals, and have found that the methods of treatment have been, for convenience, divided into palliative and radical. Under the former are included all those methods which, when completed, do not give to the individual, absolute control over the evacuation of urine. And those methods wherein the result does admit of this control by the patient are classed as radical. This is accomplished by the utilization of the sphincter ani as a substitute for the sphincter vesicae—which is beyond repair—and the rectum, or some portion of the alimentary canal as a receptacle.

DR. J. E. SUMMERS, JR., Omaha—I have operated on four patients for exstrophy of the bladder. Three of these operations were done on the old-plan methods and were unsatisfactory in results; however, they were as good as could be obtained by such makeshift plastic work. In the fourth case, a female, little trouble was experienced in fixing the ureters in the vagina. If we pass a probe into each ureter, as a guide, there is no difficulty in exposing the ureters and draining them into the vagina. The ureteral openings into the bladder are easily closed by ligation, the ligature being carried, by a full-curved needle, from the vaginal side. The bladder mucous membrane can be excised in whole or in part, and the wound closed after the manner adopted by Dr. Vance.

DR. A. M. VANCE, closing—I think this case is about as extreme a one as we ever meet, but I think the same procedure can be carried out in all cases. I use every precaution to avoid any hair-bearing tissue being introduced into the bladder. It requires great care and all the ingenuity one can bring to bear to combat the many difficulties of such a case. In bringing together the edges of these flaps I do not destroy any tissue but use the flap-splitting method. I believe my experience in these two cases will enable me to do better work in the future.

HEMORRHAGIC INFECTION IN AN INFANT DUE TO THE TYPHOID BACILLUS.*

GEORGE BLUMER, M.D.

DIRECTOR OF THE BENDER HYGIENIC LABORATORY.
ALBANY, N. Y.

The case herein reported, while differing in some respects from the hitherto recorded ones of congenital typhoid infection, probably belongs in the same category. It seems proper, therefore, before detailing the history and findings to review briefly the literature of this somewhat rare class of cases.

It seems hardly necessary to refer to the cases reported in the pre-bacteriologic era, inasmuch as they are so imperfectly recorded, according to modern standards, that considerable doubt must necessarily hang over their authenticity. Even the earlier cases recorded after the discovery of the typhoid bacillus are open to question, on account of the difficulty at the time of their discovery of distinguishing the typhoid bacillus from the organisms of the colon group.

Eberth¹ indirectly hints, and Dürk² positively states that the cases of Reher,³ Neuhauss⁴ and Chantemesse and Widal⁵ are open to doubt on these grounds, and this being the case, but eight or nine positive cases of congenital typhoid fever are left on record. There is a good deal of similarity in all these cases. The infection in all instances occurred in children born at the time that the mother was actually suffering from an

attack of typhoid fever. Many of the cases were premature births, though one or two had gone on to term. The cases of Eberth, Giglio,⁶ Freund and Levy,⁷ and Etienne,⁸ were abortions at about the fourth or fifth month; the cases of Hildebrandt⁹ and Janiszewski¹⁰ were miscarriages at about the eighth month, while in the cases of P. Ernst¹¹ and Dürk the children were born at or near term. Most of the children were born dead, though Dürk's case lived nine hours, Ernst's four days, and Janiszewski's five days. In most cases where the child had only reached the fourth or fifth month of development the post-mortem findings were entirely or almost entirely negative and the diagnosis rested on the bacteriologic findings. In some of the later cases definite macroscopic lesions were present, though in all instances except one the intestines were spared, and in that one, Hildebrandt's case, the lesion consisted of a single swollen solitary follicle.

Several of the authors point out that the disease at this age takes the form of a general infection rather than a local one, and the lesions found were those suggestive of such a state. The most constant lesion was slight enlargement of the spleen. Other lesions present in occasional cases were slight icterus (2 cases), hemorrhages into the skin, liver or kidneys (2 cases) and slight swelling of the mesenteric glands (1 case). The condition of the placenta was recorded in 4 cases and was normal in each instance. The typhoid bacilli were constant in the spleen and frequently present in the blood; they were occasionally present in the lung (2 cases), the kidney and mesenteric glands (1 case) and the liver (1 case). The placenta when examined bacteriologically showed the presence of typhoid bacilli.

In none of the reported cases is there a satisfactory description of the histological lesions in the organs of the child.

The case which I wish to report occurred in the practice of Dr. W. B. Sabin, of Watervliet, N. Y., to whom I am indebted for the clinical notes, and who kindly procured permission for the autopsy.

The mother of the child was apparently in perfect health at the time of its birth, but it was subsequently determined that about four and a half months previously she had had an acute illness, severe enough to confine her to her bed for a period of twenty-one days, characterized by a remittent form of fever, associated with occasional chills and with constipation. As the neighborhood is a non-malarious one, and the history characteristic, the presumption is that the attack was one of typhoid fever. The history of the child was as follows:

Baby H., a female child, was born at term after a difficult labor, the delivery being effected with forceps. Three days after birth the child had a slight convulsion after nursing. At each nursing after this convulsion the child would stiffen out as if in pain. On the fifth day the nurse noticed some spots of blood on the child's diaper, the amount of blood lost gradually increasing so as to become alarming on the sixth day. At this time a consultant was called in, and it was decided that the blood came from the vagina. The vagina was packed with cotton, but the cotton was soon expelled and with it some large blood clots. A further examination was made, and an eroded cervix was discovered, from which the bleeding seemed to come. This was cauterized with nitrate of silver and the vagina was again packed. The packing was again forced out, and hemorrhage from the vagina continued until the death of the child on the ninth day. Before death there also

*Presented to the Section on Pathology, at the Fifty-first Annual Meeting of the American Medical Association, held at Atlantic City, N. J., June 5-8, 1900.