anesthetic has been given with favorable results to a patient with a low hemoglobin, the exceptions do not prove the rule, and, therefore, the conclusion seems justifiable that a general anesthetic must not be given, except under the stress of absolute necessity, if the hemoglobin is low.

On the contrary, Noble,²¹ Deaver,²² Winter,⁷ Shoemaker²³ and others would put no arbitrary limit on a hemoglobin percentage below which it is unjustifiable to operate. This position, I believe, is the correct one, yet it is not to be denied that a hemoglobin percentage of less than thirty is a grave complication and that, under such circumstances, special attention to numerous details, both before and during the operation, must be given. Otherwise operation is not justifiable.

When there is active bleeding, no radical measure should be attempted until every effort to stop the hemorrhage and build the patient up has been exhausted. Curettement and packing the uterus will sometimes, although not often, be effective. Of medicines, hydrastis is better than either ergot or cotarnine preparations. Rest in bed, good food, fresh air and iron are, of course, to be employed.

In spite of the most careful treatment, some patients will not respond. If the bleeding continues active, the blood count very low and the heart's action bad, I believe the ligation of the uterine arteries, as advocated by F. H. Martin,¹² is advisable as a preliminary to hysterectomy. This, I think, should have been done in Case 4. Myomectomy should never be done. When hysterectomy is undertaken, every measure possible must be taken to shorten the time of the operation, minimize the hemorrhage, maintain body heat, and sustain the blood pressure. Carelessness regarding these measures is common, never excusable, under these circumstances criminal.

The reported cases of hysterectomy with hemoglobin under 30 per cent. are too few to justify conclusions as to results. Every gynecologist has had successful cases. Many patients have died, but for the most part the literature contains accounts of the successful cases only.

When there is grave anemia there is great danger of thrombosis and embolism during the convalescence. Phlebitis and thrombosis occurred 19 times in a series of 727 cases of fibroids which the author²⁴ reported from Kelly's clinic. Anemia was present in nearly all of these 19 cases and is undoubtedly a factor in this serious postoperative complication.

In conclusion, it may be said that anemia is frequently associated with fibroids; it is often grave and constitutes a menace to life. Every one has recognized its dangers, but in the literature it has not received the attention its importance demands. More cases should be published in detail and thus more light shed on this phase of the myoma problem.

ABSTRACT OF DISCUSSION.

DR. DANIEL H. CRAIG, Boston: I wish to report a case which came under my care a few months ago. The patient was referred to me by Dr. Courtney, a neurologist, of Boston. The case was one of paralysis agitans with general anemia, apparently spinal, in a single woman, aged 36. There was hemoglobin percentage of 26 and a fraction. The patient was actively bleeding, and every ordinary means had been employed to check the flow before she was referred to me. Hysterectomy

was advised, and at operation a fibroid was found occupying the entire thickness of the uterine wall, impinging on the mucous membrane on one side, and on the other there was a very dangerous deposit of calcareous matter. A distinct submucous polypi was found, filling practically the entire uterine cavity, and very small fibroid nodules. A supravaginal hysterectomy was done. Aside from giving the patient liquids for forty-eight hours before the operation, no unusual precautions were taken, although at the time of operation everything necessary was at hand for transfusion of salt solution should it be necessary. The patient stood the operation well and there was not a single complication; and now, six or seven months afterward, the woman so far as her pelvic condition goes is in perfect health. This was a very marked case of fibromyoma with anemia in which the anemia was an extremely important factor and yet operation was justifiable.

DR. H. T. BYFORD, Chicago: I have had the opportunity of watching some of these patients for a long time after operation, and from my experience I am satisfied that Dr. Schenck is right when he says that fibroid tumors do not of themselves cause vascular changes. Nor do I think that they give rise to any other unfavorable changes in the system until they are large enough to cause some pressure or other interference, unless it be through the anemia. I have had patients who stood the operation well, but who never recovered from the anemia. I remember two patients whom I watched for about ten years without finding the hemoglobin above 60. They had been chlorotic for a long time. I do not know that they ever have or will recover from it. Such cases offer an argument in favor of not letting the patients go too long without operation.

SOME OF THE MORE UNUSUAL FORMS OF EPITHELIAL GROWTHS OF THE SKIN.*

J. A. FORDYCE, M.D.

Professor of Dermatology and Syphilology, University and Bellevue Hospital Medical College.
NEW YORK CITY.

Experience with a large number of epithelial new growths of the skin, studied both clinically and microscopically, has led to some observations which have seemed worthy of being placed on record. Most of these have been confirmatory of the work of others, while a few of them may claim a limited originality. The two large divisions of cutaneous epithelial tumors into basal cell and prickle cell epitheliomas are justified by their histologic structure and clinical course. In the one the basal cells of the epidermis and the outer root sheath of the hair are chiefly or solely active, this class including among others, the well-known rodent ulcer which is so familiar to dermatologists.

Some modern writers on cancer, as Sutton,¹ attribute to rodent ulcer a sebaceous gland origin. It is true that many rodent ulcers in their early stages produce a lobulated growth simulating in a general way the outlines of the sebaceous glands, and it is not disputed that they may originate from the epithelial layer of the gland as they do from the hair follicle or germinal layer of the epidermis, but it is only in this way that a sebaceous gland origin can be invoked. They are not genetically related to the secretory glands, as are cancers of the breast or of other glandular organs. Rodent ulcers, as well as other basal cell growths, only rarely involve the lymph nodes and produce metastases, and no adequate explanation has as yet been offered as to why they spare the nodes.

21. Am. Jour. Obst., September, 1901.

22. Am. Med., April 15, 1903.

23. THE JOURNAL A. M. A. xlviii, 1282.

24. New York Med. Jour., Sept. 6, 1902.

* Read in the Section on Cutaneous Medicine and Surgery of the American Medical Association, at the Fifty-ninth Annual Session, held in Chicago, June, 1908.

1. Tumors Innocent and Malignant, Fourth Ed., 1907.

The spino-cellular type of tumor, on the other hand, originates in the layers above the basal cells and reproduces its own kind, usually resulting in the cycle by which more or less perfect squamous cells are formed. This type of growth, as a rule, is more malignant in that it soon implicates the lymph nodes and leads to metastases. It is the form met with at the muco-cutaneous juncture, the mucous surfaces as well as in the skin. In the latter, tumor formation preceding the ulceration is often the predominant feature, whereas, in the basal-celled epithelioma there is a greater tendency to local ulceration and serpiginous extension with less tumor formation.

Various subdivisions have been made of these two main types, resulting in a lengthy and confused nomenclature. In tumors of the prickle cell variety there is not so much divergence from the type of growth; in some, the cells in their evolution do not proceed to the formation of pearls, but maintain the spinous or polygonal cell throughout. In others, the exaggeration of the normal evolution of the cell is so marked that the growth is almost entirely made up of these centers of cornification. Between these two extremes, of course, there is every intermediate grade. An example of histologic activity is illustrated in Figure 1. The tumor followed an injury to the leg received nine months previously and developed with such rapidity that it terminated fatally in a very short time. In this case a secondary growth was found just beneath the primary one, and it is probable that the secretion containing epithelial cells had been implanted on an abrasion below and thus given rise to the new growth. This is analogous to the infection of the edges of the wound and lines of suture which are sometimes seen after operation for cancer of the breast or other organs.

In the basal-celled variety of tumors the clinical picture is quite as diverse as the histologic one. For instance, I have seen multiple epitheliomata following psoriasis assume the form of fungating growths, which, microscopically, belonged to the rodent ulcer type. Several years ago I reported the case of a small circumscribed pea-sized growth which had developed in the scar tissue of lupus. At that time it was considered an endothelioma, but further study of the sections has led to the conviction that it was a basal cell tumor with cystic degeneration. A tricho-epithelioma, which will be described in detail later, presented the clinical appearance of a morphea without a suggestion, macroscopically, of a tumor formation. The so-called benign cystic epitheliomata, either solitary or multiple, appear under varying guise as the expression of one and the same affection. Certain of them may be of the pure basal cell type with colloid degeneration of the cell masses, while others may be mixed growths with centers of cornification. Both of these forms have been examined histologically by the writer.

The so-called cylindromas probably belong to the basal cell group, although an endothelial origin has been attributed to them by Billroth, Borst, Klebs and others. In addition to the multiple tumors of the scalp described by E. Spiegler,² single tumors of the face and mucous surfaces have been seen. I have removed a number of such growths from the face which, clinically, were brownish-red in color, soft in consistency, and of a relatively benign course with little tendency to recur-

rence after extirpation. Microscopically, they were situated in the corium, had apparently no connection with the epithelial covering or pilo-sebaceous apparatus, and were made up of cells with small well-staining nuclei and little cytoplasm, corresponding closely with the cylindrical cells of the basal layer. About the cell aggregations and inside of them a hyalin degeneration had taken place in the form of cylinders, which circumstance gave the tumor its name (Fig. 2). Darier³ suggested that such growths originated from embryonic rests, possibly glandular, and that their hereditary character speaks for this view.

Epithelial tumors which develop in connection with another morbid process, as a rule, are of the prickle cell variety, though occasionally the basal-cell type is encountered. They afford excellent opportunities for the study of epithelial growths of the skin. Such studies when extended to the precancerous conditions such as senile keratosis, moles, warts, xeroderma pigmentosum, leucoplakia, x-ray dermatitis, Paget's disease, etc., greatly enhance our knowledge of this large subject and emphasize the futility of incriminating one etiologic factor in tumor formation. Dermatologic experience certainly does not support the infective theory of tumors.

Among a large number of secondary growths which I have seen is the so-called Marjolin's ulcer, the name given by Marjolin over more than half a century ago to chronic ulcers which had become malignant. A few years ago DaCosta⁴ revived this name for the condition described, but it will probably be more popular in surgical than in dermatologic nomenclature, as the term is too ambiguous and is not suggestive of the real nature of the affection.

In these chronic ulcerations, the young epithelial cells which lie in the granulation tissue become active when the local conditions are favorable and lead to tumor formation. I have seen bilateral ulcerations from suppurating buboes of the groin simultaneously become carcinomatous; also ulcers of the leg and an extensive ulcer of the forearm from a burn. One case, remarkable for its very chronic course, was that of an ulcer of the thigh from a burn received forty years before. It had broken down and healed several times during this interval and then developed into a squamous-celled cancer (Fig. 3) of slow course, the benignity being explained by the histologic structure which showed the sharp limitation of the epithelial growth by the sclerotic connective tissue beneath (Fig. 4).

The tumors which I have seen secondary to syphilitic ulcerations have been of the squamous-celled variety, but as they were all met with about the lip and tongue, it was natural that they should be, as basal-celled epitheliomas of the mucous membranes are not so commonly met with. They do occur, however, and mention has already been made of the cylindromas of the buccal cavity which have come under my observation. I have also seen a tumor in the roof of the mouth over the hard palate and gums, which consisted of a reddish-brown infiltration and had been mistaken for lupus. A biopsy showed a growth of basal cells with a tubular arrangement.

An examination of leucokeratosis of the mouth reveals a condition analogous to that met with in many other precancerous keratoses, viz., a proliferated epidermis, which, when assuming malignancy, shows hy-

2. Endotheliome der Haut: *Archiv f. Derm. u. Syph.*, 1899, i, 165.

3. *La Pratique Dermatologique*, iv, 687.
4. *Ann. Surg.*, 1903, xxxvii, 496.

perchromatic nuclei and mitotic figures. A lymphocytic and plasma cell infiltration is present in the upper corium, which reaction is essentially the same wherever growing epithelium is beginning to show malignancy.

It seems reasonable to assume that nevi do not always originate from the same embryonic structure, for it is possible that the mesoblast as well as the epiblast may give rise to these developmental anomalies. In my experience there is more evidence in favor of an epidermal than an endothelial origin. I have found, however, that the type of cell varies somewhat, even in nevi of the same class. In some of the pigmented warty moles I have seen perpendicular columns of cells which were closely related to the basal cells of the epidermis. They were elongated or cylindrical with deep-staining nuclei and relatively little cytoplasm (Fig. 5). Again, in nevi which were similar, clinically, to the preceding, the cells had the arrangement of a mosaic and were of a transitional type, being large and globular or polygonal, with vesicular nuclei, clear cytoplasmic bodies and distinct borders (Fig. 6). In none of them was it possible to demonstrate prickles. Sometimes these varieties occur mixed and next to a column of small deeply-staining cells one will find a nest of the larger pale kind.

I agree with the contention of Unna, Gilchrist and others that the majority of pigmented tumors arising from nevi are of epithelial origin, though it is not possible in every case to exclude a mesodermic origin of certain of them. This is opposed to the views of Johnston and other supporters of the endothelial genesis of moles that melanomas from nevi are sarcomas.

CARCINOMATOUS LYMPHANGITIS SECONDARY TO CANCER OF THE BREASTS.

Malignant disease of the breast seldom comes to the dermatologist unless it is complicated with a cancerous lymphangitis or unless the latter is the predominating feature of the disease, as in the case about to be related. In this case nodular lesions appeared in the skin over the mammary region simultaneous with or shortly after the occurrence of the retraction of the nipple. From the history which the patient gave, it seemed likely that carcinoma of the breast followed on a mastitis subsequent to childbirth.

History.—Mrs. C. C., aged 28, had had six children, two of whom were dead. Her mother was living and well; her father died of "jaundice" at the age of 49. She had one sister living. For nine years she had had trouble with her right breast, following an abscess after the birth of her first child, and she said none of her children would nurse from it. She was first seen in July, 1907, when she stated that the condition then present had begun a year previously. At that time she noticed "spots" on the skin of her chest and afterwards nodules in her right breast. Six months later the left breast became involved.

Examination.—At the time of examination she was in the fifth month of gestation. Both breasts were hard and the nipples slightly retracted. Over the front and back of the chest on the right side were scattered numerous lenticular lesions (Fig. 7), a few of which were encrusted. A lymphangitis also was present, which manifested itself by a reddish-brown discoloration of the skin, extending along the line of the lymphatics from the nipple over the shoulder, the back and down the right arm (Fig. 8).

Later History.—She was not seen again for two months, during which time the disease had made extraordinary progress. The right upper half of the chest was red and the seat of numerous encrusted lesions; the lower half showed a linear lymphangitis and lenticular nodules. The surface temperature was not elevated as in ordinary septic lymphangitis. Both

nipples were retracted, the glands were board-like and her right arm was edematous and red. She was markedly cachectic and had great difficulty in breathing, partly on account of the infiltration and partly on account of the extreme weakness from which she was suffering. She gave birth to a healthy child on November 12 and died on December 26 from exhaustion. In the meantime the tumors of the breasts had increased in extent, many of the nodules in the skin of the chest had undergone ulceration, and the edematous condition of the arm had increased. The lymphangitis which had been present only on one side involved the left side also, but was more accentuated on the right side.

Examination of Specimen.—Specimen removed from the skin over the site of lymphangitis (Fig. 9) showed the lymphatic channels of the corium filled with large epithelial cells containing hyperchromatic nuclei and numerous mitotic figures. The central portion of many of these cellular collections had undergone a fatty degeneration and resulted in cyst-like cavities containing epithelial detritus. The collagenous tissue immediately about the lymph vessels was reinforced by a new growth of fibrous bands with numerous spindle cells. Beyond an acanthosis and edema, the epidermis participated only passively, the growth from the subepidermic lymph spaces extending into the epiderm and forming cysts at its expense. As can be seen from the illustration, the cancer cells have entirely displaced those of the rete and so compressed them that they form the walls of more or less rounded or irregular spaces containing tumor cells, degenerated protoplasm and chromatic particles. It was over these areas that ulceration took place and it was the contents of these small cysts which could be extruded on pressure.

In his chapter on Paget's disease Ribbert⁵ states that the edematous and vacuolated cells present in that condition are not the primary epidermal cells which have suffered this change, but foreign elements which have emigrated from the tumor growth beneath. This view is not substantiated by a study of the illustrations showing the invasion of the epidermis by cancer cells in the case of lymphangitis (Fig. 9) and the epidermic changes in Paget's disease of the buttock (Fig. 10) given herewith. In the former the invading tumor cells formed cysts entirely distinct from the normal cells of the part, and there was no migration of individual cells. In the latter, where early changes are pictured, it is very evident that the cells belong to the epidermis.

Histologic examination of the lenticular nodules showed the lymphatic spaces of the entire corium infiltrated with rows of epithelial cells running parallel with the epidermis. Many had undergone a fatty degeneration and stained poorly. In the connective tissue surrounding the growth a mucinous degeneration had taken place.

PAGET'S DISEASE OF THE BUTTOCK.

During the past few years a number of cases of extramammary Paget's disease have come under my observation. One of them, a lesion on the buttock, was made the subject of a report.⁶ It was cured with x-rays, no recurrence having taken place after three years. The case about to be cited is interesting on account of having the same localization and many features in common with the other.

History.—J. F., aged 68, born in Ireland, had for many years been a gardener by occupation. He had a seborrheic dermatitis and patches of senile keratosis over the face and backs of the hands, together with telangiectases and pigmentations—a condition like that met with in sailor's skin. The affection for which I was consulted had slowly progressed over a period of thirty years, was located on the right buttock and consisted of an irregularly rounded, non-ulcerating area

5. Beitr. z. Entstehung der Geschwülste, 1906.

6. Jour. Cutaneous Diseases, 1905, xxiii, 193.

4 by 5 inches in diameter, dark red in color, with a marked polycyclic margin and a slightly nodular center. The diagnosis of Paget's disease with secondary epithelioma was confirmed by microscope. The condition slowly resolved under x-ray.

Examination of Specimen.—The tissue from the margin of the lesion (Fig. 10) showed acanthosis with swollen edematous epithelial cells, vacuoles and mitotic figures. Some had lost their prickles, had a dark contour, but still retained their relation to neighboring cells; others were separated and free in spaces. There was a dense sub-epidermic infiltration of

atypical mitoses. A number were detached from their neighbors. The tumor itself (Fig. 11) extended through the corium by branching processes; its upper portion was made up of cells conforming more to a transitional type, being ovoid or more or less rounded with large, pale vesicular nuclei. The protoplasm was degenerated and here and there vacuolated, suggesting oftentimes prickles. At different points of the growth the degeneration involved a number of cells and gave rise to cysts filled with fragmented nuclei and protoplasmic detritus. In the more recent and advancing parts of the tumor the cells were smaller and the nuclei richer in chromatin. The sclerotic collagenous tissue contained an inflammatory infiltration, most marked beneath the epidermis and about the upper half of the growth.

In my experience, extramammary cases of Paget's disease do not differ essentially from those of the nipple, unless it is in degree. In both instances the process is the same, namely, edema, vacuolation, loss of prickles,

Fig. 1.—A cancer of the leg following an injury. The tumor had a very malignant clinical course, its activity being evident histologically by the formation of numerous "pearls."

Fig. 2.—Cylindroma of the scalp. The cells of this tumor are small and closely resemble those of the basal layer of the epidermis. Inside the cellular aggregations hyalin degeneration has taken place in the form of small cylindrical areas, from which the growth derives its name. Immediately surrounding the cell masses the connective tissue has likewise undergone a degeneration, appearing as a narrow homogenous band of hyalin.

plasma cells and lymphocytes. Sections from the center of the lesion consisted of an intact slightly verrucose epidermis with striking changes in some of the rete cells. They were likewise edematous, contained hyperchromatic nuclei and many

Fig. 3.—Marjolin's Ulcer. Burn of hip and side of trunk when 3 or 4 years old. It healed and broke down several times. Now has ulcer 3×4 in. on hip; surface knobby; here and there islands and streaks of new epidermis. Biopsy showed typical squamous-celled cancer, sharply limited by dense mass of sclerotic tissue beneath.

separation of the cells and erosion, but in the mammary region it apparently attacks the epidermis with greater severity. Tissue from the edge of ordinary ulcers has shown that the cells are sometimes swollen and vacuolated or ballooned. This change, which is due to the simple imbibition of fluid, does not, however, follow the pathologic cycle of Paget's disease.

MULTIPLE EPITHELIOMATA OF THE ARMS AND CHEST.

The clinical picture of epithelioma of the skin includes a number of types which are not generally recog-

nized as belonging to this group, but are sometimes mistaken for infective granulomata or other diseases of the skin. They are best illustrated by examples which have fallen under my observation in which the diagnosis has been confirmed by the microscope. Sometimes there is little new growth, but a slowly spreading dark red patch, slightly scaling on the surface, with irregularly rounded outline, which probably developed on areas of chronic seborrheic dermatitis which later as-

Fig. 4.—Cancer of the skin following an old ulceration due to a burn. The photograph shows the downgrowth of the tumor sharply limited by the underlying dense connective tissue.

Fig. 5.—Nevus papillaris. Showing the styloid processes from the epidermis merging with the nevus cells. The latter are arranged in perpendicular columns, separated by collagenous tissue, and are made up of a small type of cell more in conformity with the basal cells, having deep-staining nuclei and little protoplasm. The growth is pigmented.

sumed an appearance of senile keratosis, or they are accompanied by patches of verruca senilis. These lesions are usually seen too late to follow the sequence of their development. In some of them the edges do not show

Fig. 6.—Nevus papillaris. Showing elongated and anastomosing rete pegs enclosing nevus cells which are probably of a transitional type. They are globular or polygonal, closely opposed to one another, with vesicular nuclei, clear cytoplasm and well-marked borders. Pigment is present.

Fig. 7.—Lenticular carcinoma. Hard tumor of right breast with retracted nipple. Left breast also hard; nipple slightly retracted. Multiple lenticular nodules in skin covering both. Some eroded and discharging a serous fluid. Large erythematous patches over the right shoulder, upper part of right arm and right side of back (See Fig. 8), a distinct unilateral distribution. This erythema in places shows a slight infiltration—a carcinomatous lymphangitis in the early stages.

Fig. 8.—Lenticular carcinoma. Back view of patient shown in Fig. 7.

the elevated, rolled pearly margin, but they have a depressed center with little tendency to ulceration. When situated on the face a narrow rim of ulceration is sometimes present about the margin. Seen in the typical sites of rodent ulcer these patches are not difficult to diagnose, but when met with on the extremities and trunk they are strikingly like the late syphilitic neoplasms. An instance of multiple lesions of this character lately came under my care where a number of

he had consulted numerous practitioners since that time. When I saw him he had multiple lesions over the chest and arms, dark red in color, with scalloped margins, showing a very slight tendency to elevated or pearly rolled edge. They had persisted for a period of ten years and had progressed slowly peripherally without central involution, as seen in syphilitic infiltrations.

Examination of Specimen.—Histologically, the lesion was a superficial epithelioma (Fig. 12) situated under an intact,

Fig. 9.—Cancerous lymphangitis secondary to cancer of the breast. Showing the cancer cells in the lymphatics of the skin and the mode of extension of the disease.

Fig. 11.—Tumor in Paget's disease of the buttock. Growth consisting of a transitional type of epithelium with vesicular nuclei and well marked cell bodies devoid of prickles. A mucinous degeneration of the tumor present. The supporting connective tissue is sclerotic.

Fig. 10.—Paget's disease of the buttock, showing the changes which take place in the epidermic cells in this disease. They are very much swollen from edema, vacuolated and have their nuclei pushed to one side. Many of the cells have lost their prickles. In the corium there is a dense infiltration of lymphocytes and plasma cells.

good observers had made the diagnosis of syphilis on account of the history of such an old infection.

History.—Mr. J., aged 72. Patient had had lues over forty years ago, for which he had active treatment, and for which

Fig. 12.—Rodent ulcer. Lesion from chest showing proliferation from basal layer of epidermis budding of the cell masses and cystic degeneration.

flattened epidermis, with which it was connected here and there by a delicate pedicle or in relation with a hair follicle. The cells were of the cylindrical type, being typical and uniform at the periphery of the cell masses, but showing a polymorphism in the center on account of pressure. Here many of them were elongated and spindle-shaped, like connective tissue cells.

Fig. 13.—Tricho-epithelioma. Fine strands of epithelial cells ramifying through the derma. To the right of the center of illustration is seen a follicle in process of budding. The intervening connective tissue shows a partial band-like sclerosis and a mucoid degeneration.

Fig. 14.—Tricho-epithelioma, showing proliferation of basal layer of hair follicle and extension of growth through the cutis.

Fig. 15.—Rodent ulcer. Growth began near left ala of nose and gradually extended until it involved half of the upper lip and almost all of the lower lip. Loss of tissue was so extensive that canine and bicuspid teeth were exposed. Tumor of rodent ulcer type was confirmed by microscope.

Fig. 16.—Rodent ulcer. After treatment with x-ray. Same patient as shown in Fig. 15.

A cystic degeneration had taken place at various points, the spaces containing emigrated leucocytes, protoplasmic detritus and some blood pigment. In areas in which this degeneration was beginning the cells were markedly swollen, the nuclei had lost their chromatin and appeared like shadows. In the corium lymphocytes and polynuclear leucocytes were present.

TRICHO-EPITHELIOMA.

In the following case the diagnosis was only made after a microscopic examination. There was nothing in the clinical appearance of the lesion to suggest an epithelioma.

History.—Z. L., female, aged 80, had on her upper lip near the left ala of the nose a small mole, which had broken down some months previously, and under the microscope proved to be a rodent ulcer with marked degeneration. Situated near this lesion on the lip was a small, circular, sharply limited sclerous area about $\frac{3}{8}$ in. in diameter, which the patient said had existed for only about two years. It bore the clinical features of a morphea, but histologically proved to be a basal-celled epithelioma having its origin in the lanugo hair follicles.

Examination of Specimen.—Histologically the tissue presented an intact epithelial covering with the new growth occupying the entire cutis. It was made up of fine strands of epithelium, which on longitudinal section showed two or more layers branching in various directions and occasionally swelling out into cyst formation; on transverse section there was an alveolar arrangement of three or more cells. The cells themselves were small, with deep-staining nuclei and a narrow rim of protoplasm. The follicular origin was demonstrable in their connection with the lateral buds given off by lanugo hairs (Fig. 13) in the derma and the offshoots from the superior part of the follicle (Fig. 14). A number of horny cysts were present with two or more peripheral layers of cells and central lamellated corneous material. The collagenous tissue immediately surrounding some of the alveoli had undergone a hyalin change or was condensed like a basement membrane. Between the epithelial processes were long sclerous bands and between these a mucoid degeneration of the collagen. The sweat and sebaceous apparatus showed no appreciable abnormality. A moderate round cell infiltration was present throughout.

This growth was interesting for several reasons, in that there was no suggestion of a neoplasm, on account of its close proximity to another basal cell tumor of different conformation and because of the development so late in life of this particular kind of growth. It belongs to the class designated by a varied nomenclature, but the name tricho-epithelioma, given by Jarisch, Hartzell and others, is perhaps most fitting, as it is simple and indicates a certain structure in contradistinction to others having their origin in the basal cells of the epidermis or follicles.

COINCIDENCE OF PRICKLE CELL EPITHELIOMA AND RODENT ULCER.

It is unusual to meet with instances of prickle cell epitheliomas and rodent ulcers in the same individual, as multiple epitheliomas usually preserve the same type throughout. I have, however, had a case in which there was a typical rodent ulcer of the temple and a prickle cell growth over the bridge of the nose. Similar epitheliomas in the latter locality have been met with in several instances in my experience. In the early stages it was impossible to differentiate them clinically from the ordinary rodent ulcer. In one case, although the growth was healed under x-ray, it was soon followed by metastasis of the lymph node at the angle of the jaw, which recurred several times after operation and finally led to a fatal result.

History.—Mr. G. M., aged 43, had a small, pearly, nodular lesion of twenty years' duration, situated in the temporal

region. It began to grow two or three years before I saw him, and slowly spread with central depression and elevated pearly margin. It was typical of an early rodent ulcer. The lesion on the nose had been present a year and a half and clinically was somewhat similar to the one on the temple. In addition he had another small, pearly growth beneath the right eye of seven months' duration. The lesions were enucleated, followed by the galvano-cautery, a cure resulting.

Examination of Specimen.—Microscopically the lesion on the temple was a rodent ulcer of usual type. The one on the nose was of the pure spino-cellular variety, consisting of alveoli of prickle cells showing no tendency to pearl formation. It may have had its origin in the follicular cells as the alveoli seemed to be connected with a proliferating follicle.

EPITHELIOMA OF THE LIP IN A WOMAN.

Epithelioma of the lip, although of frequent occurrence in the male, is seldom met with in woman. The following case is, therefore, of interest:

History.—M. S., aged 48, said that the tumor began sixteen years previously from a small lesion near the ala of the nose. It spread very slowly until it involved by direct extension the vermilion border of the lips. The ulceration at the left angle of the mouth and adjoining upper lip was so extensive as to expose the bicuspid and canine teeth (Fig. 15). The growth yielded to x-ray (Fig. 16). Under the microscope it was seen to be made up of basal cells with a delicate tubular arrangement, conforming with the rodent ulcer type of growth.

In regard to x-ray treatment of epitheliomas of the skin, a longer experience with it has led to much greater conservatism in recommending it. I have found that better results have been obtained from the use of Roentgen rays after the new growth has been thoroughly curetted. In this way it is possible to apply the rays directly to the more actively developing cells. It has also been my experience that the prolonged use of the ray in malignant cutaneous growths is as a rule contraindicated, especially its persistent use after repeated relapses, for it is believed that in certain cases it renders the tissue more vulnerable to the malignant process. I have in mind a number of cases of epithelioma and several of Paget's disease in extramammary locations in which no relapses have occurred after from two to four years. On the other hand, I have seen a number of instances in which x-ray had been employed to treat recurrences, each one of which involved greater areas of skin, both as to depth and circumference. In using the Roentgen treatment in such cases, one should not forget that excellent results may be obtained in many of these forms of minor epithelioma by the use of the curette followed by a chemical or the actual cautery. X-ray or radium should be reserved for epitheliomas of the inner canthus, the eye-lids or locations not accessible to other methods of treatment.

ABSTRACT OF DISCUSSION.

DR. JOSEPH ZEISLER, Chicago: Dermatologists have become convinced of the non-contagious nature of carcinoma, and I believe that a discussion along those lines between dermatologists, general surgeons and pathologists would be fruitful. We see cases of xeroderma pigmentosum, with the development of carcinoma under the influence of only one physical factor, namely, the influence of light; we see the development of carcinoma under the influence of the x-ray; we see it in the senile skin, simply due to dryness of the skin similar to that produced under the influence of the x-ray; we see it develop from nevi; so that we have sufficient proof from our own clinical observations to become convinced of the non-parasitic nature of carcinoma. I believe that this would be a wholesome lesson for the dermatologist to teach the general surgeon.

Dr. M. L. HEIDINGSFELD, Cincinnati: The distinction which Dr. Fordyce made between the prickle and the cylindrical cell type appealed to me very strongly. In my own work I have realized, both in a pathologic and clinical way, the difference between the so-called cylindrical and the prickle-celled epithelioma, and my observation has led me to a different idea as to the development of these tumors. The so-called basal-cell epitheliomata, I believe, are derived from the ducts of the sudoriferous glands, and are more of an endothelial character. In malignancy all neighboring structures such as the hair follicles, sweat glands, etc., are much stimulated to overgrowth and development by the abnormal conditions, and it is difficult to infer what elements of tissue are chiefly involved in the pathologic process. I believe the origin of these growths to be in the lower structures, the ducts of the sudoriferous glands, rather than the basal cells. In regard to the possibility of the mesoblast being involved in some of these processes, I will cite a case I saw a year or two ago. The lesion was a rapidly growing mole on the chin of a young man. I excised this and found that there were sixty or seventy small osteomata of laminated bone type, containing bone cells, corpuscles, Haversian canals and canaliculi, situated in the cutis and entirely isolated from all the underlying tissues. This case clearly indicates that mesoblastic tissue in the epiblast must owe its origin to misplaced embryonic development, and I heartily concur with Dr. Fordyce as to this probable form of origin of many new growths.

TRAUMATIC RUPTURE OF THE UTERUS AND BLADDER DURING LABOR AT FULL TERM. HYSTERECTOMY, REPAIR OF BLADDER AND RECOVERY.*

JOHN YOUNG BROWN, M.D.

Chief Surgeon to St. John's Hospital.

AND

PERCY H. SWAHLEN, M.D.

Resident Obstetrician, St. Ann's Maternity Hospital.
ST. LOUIS.

The object of this paper is not to review the literature of rupture of the uterus, but to report an interesting case.

Patient.—A Russian by birth and of Jewish extraction, at the age of 16, left her native home and came to America. She had ten brothers and sisters, nine of whom were dead; the causes of death were unknown. The father and mother are living and well. The patient knows of no particular disease in family and stated that her mother and sisters experienced no particular difficulty in giving birth. She had no drug or tobacco habit. Menstruation was regular and without pain.

History.—The patient had only done housework since her marriage to her first husband, at the age of 22. One year after her marriage she gave birth to a 6 months' child, which died within a few hours. She was divorced from her first husband after two and a half years and remarried at the age of 26. One year later she was delivered of a full-term, seven-pound child, which is still living and well. She was in labor about thirty hours, but no obstetric interference was deemed necessary.

Present Pregnancy.—The woman had reached her twenty-ninth year. Pains began normally at what she had estimated was the end of the ninth solar month of pregnancy. She was attended by a midwife for the first thirty hours and then two physicians were called in and endeavored to extract the child with forceps. When they failed in their efforts, a specialist was summoned who, with the patient anesthetized, found the vagina wide, the cervix almost completely dilated, the fetal head apparently transverse above the pelvis in beginning engagement, but caput succedaneum made an exact diagnosis

impossible. The lower uterine segment was very thin and reached the umbilicus. The physician then made a futile attempt at delivery with forceps and advised sending the patient to a hospital. The family refusing, version was decided on. No pelvic measurements were made and the fetal heart sounds were not to be heard. The back of the child was to the mother's right. A hand was inserted into the uterus with great difficulty and the lower uterine segment was found to be very tense but by pressing on the fundus with the other hand the left leg was finally brought into the vagina. At this stage another physician endeavored to complete the version and neglected holding the fundus. Suddenly he announced that the head had slipped upward. Abdominal palpation revealed the well-contracted uterus to the right and the easily defined fetal head to the left. The hemorrhage from the vagina was very small in amount. Owing to the obstinacy of the family, the patient could not be brought on the operating table till almost an hour and a half had elapsed. On admission to the St. Louis City Hospital, the patient was in profound shock. The pulse was 130, respirations 28 and temperature subnormal. Immediate preparations for operation were made.

Operation.—Ether anesthesia was given, the abdomen opened through a median incision. On entering the peritoneal cavity, a large quantity of blood and amniotic fluid escaped. The back of a well-developed, full-term child presented in the incision. The child was quickly delivered through the abdominal wound. The rent in the lower uterine segment extended two-thirds around the cervix and well up into the body of the uterus, being T-shaped in character. Both uterine arteries were bleeding. The bladder was ruptured. The broad ligaments were quickly clamped, and the uterus caught and amputated above the cervix. The rent in the bladder was repaired with two layers of silk suture. The uterine stump was covered with peritoneum. The abdominal cavity was flushed with hot saline solution and closed with through-and-through sutures of silkworm gut. Anesthesia was begun at 12:28 a. m. The patient was returned to bed at 1:05 a. m.

Postoperative History.—Convalescence was uneventful, the patient being discharged from the hospital March 3, 1907, twenty-eight days from the date she was received.

As rupture of the uterus is divided etiologically into two classes, spontaneous rupture and rupture by violence, it is of interest to consider here if there were any predisposing causes in this case. Contracted pelvis among us are not so common as in Europe. It is an accepted fact, according to Bandl, Baisch and other investigators, that an abnormally small pelvis is the chief cause of spontaneous rupture of the uterus—not a rachitic pelvis, in the majority of cases, but a generally contracted flat pelvis—and pelvimetry, as performed during the convalescent period of our patient, proved that she came under that class, the measurement being:

Interspinal	24.5 cm.
Intercristal	26.5 cm.
Intertrochanteric	28.0 cm.
External conjugate	17.0 cm.

The writers tell us that an edema of the cervix, due to the pressure between the head of the fetus and the pelvis of the mother, is an accompanying factor in this condition, as a result of which the resistance of the tissues is lessened. In this case the edema was practically absent.

We notice, moreover, that the woman was a multipara, and that she had been in labor thirty hours (the time of rupture of the membranes not obtained) before a physician was called. As the woman, at the age of 27 years, had been in labor twice as long as the average primipara, and as it is a fact, according to Döderlein, that the children increase in size up to the fifth, a birth along normal lines was hardly probable.

* Read in the Section on Obstetrics and Diseases of Women of the American Medical Association, at the Fifty-ninth Annual Session, held at Chicago, June, 1908.