No. 1.]

[Price 1s.

TRANSACTIONS
OF THE
GLASGOW GEOLOGICAL SOCIETY,
BEING
A PAPER
ON THE
GEOLOGY OF THE CAMPSIE DISTRICT.

BY
JOHN YOUNG,
VICE-PRESIDENT.

GLASGOW :
PRINTED FOR THE SOCIETY,
BY ROBERT ANDERSON, 85 QUEEN STREET.
1860.

OFFICE-BEARERS
OF
THE GLASGOW GEOLOGICAL SOCIETY.

President.

JOHN SCOULER, Esq., M.D., LL.D.

Vice-Presidents.

THOS. STRUTHERS, Esq. | JOHN YOUNG, Esq.

Honorary Secretary.

JAMES HORNE, Esq.

Council of Management.

MESSRS. E. WUNSCH.

JAMES ARMSTRONG.

JAMES THOMSON.

JOHN DENNISON.

WILLIAM CAREY.

H. W. CROSSKEY.

MESSRS. WILLIAM JOHNSTON.

JAMES STEWART.

J. C. DOUGLAS.

JAMES FARIE.

MARK FRYAR.

JOHN GOODALL.

TRANSACTIONS
OF THE
GLASGOW GEOLOGICAL SOCIETY,
BEING
A PAPER
ON THE
GEOLOGY OF THE CAMPSIE DISTRICT.

BY
JOHN YOUNG,
VICE-PRESIDENT.

GLASGOW :
PRINTED FOR THE SOCIETY,
BY ROBERT ANDERSON, 85 QUEEN STREET.
1860.

The Secretary of the Glasgow Geological Society, in offering to Members and the Public the following pages, embraces the opportunity, on the part of Mr. YOUNG, of acknowledging his obligations to Mr. STRUTHERS, his colleague in the Vice-Presidentship, for having kindly undertaken the necessary editorial labours.

The list of fossils, with the sections and tables of strata at the end of the Paper, it is hoped, will enhance its value to the student of Geological Science.

121 DOUGLAS STREET,
GLASGOW, *November*, 1860.

GEOLOGY OF THE CAMPSIE DISTRICT.

IN compliance with the wish of the Society, I have now to offer a few remarks on the Geological features of the Campsie district, chiefly for the guidance of those members who have not yet visited that interesting locality, but who may find an opportunity of doing so during the course of the summer.

The parish of Campsie is situated in the south-west of Stirlingshire; and is bounded on the north-west by Killearn; on the west by Strathblane; on the north-east and north by Fintry; on the east by Kilsyth; and on the south by Kirkintilloch. Stretching east and west eight miles in length by five in breadth, it embraces a considerable strath, with the adjacent elevations on the north and south; the former being usually called the Lennox hills or Campsie fells, and the latter the South hill. Strictly speaking, the Lennox hills embrace the heights between Stirling and the valley of the Blane; at the same time forming a portion of the great chain of trappean heights which extend across Scotland in a north-easterly direction from Ardrossan to St. Andrews. The middle division of this chain lies between the towns of Dumbarton on the south-west, and Stirling on the north-east, and embraces the Campsie fells, and the Killearn and Fintry hills in the centre, with the Strathblane and Kilpatrick hills on the west and south-west, and the Kilsyth, Gargunnoch, and Touch hills on the east and north-east. The length of the Lennox hills, from east to west, is 25 miles, their greatest breadth being about 10 miles, and their average height from 1000 to 1200 feet; but Ben Carron, or the "Muckle Ben," the highest elevation of the range, attains an altitude of 1500 feet above sea-level.

Before entering upon a description of the igneous and stratified rocks exhibited in the various romantic glens, and natural sections along the hill-sides, and noticing the different organic remains which the fossiliferous deposits have yielded, let us glance for a moment at

the more prominent physical aspects of the district, which cannot fail to interest the geological observer as he traverses its length and breadth in pursuit of his favourite study.

Col. Imrie, in a paper on the Campsie hills, read before the Wernerian Society, Edinburgh, in 1812, gives the following graphic description of their general configuration:—"The general surface of the Campsie district may, in the strictest language, be denominated highly undulated. Its surface majestically portrays the Bay of Biscay after a storm or western gale, and its high undulations are of great length from south to north, and nearly in that direction the most of them stretch from side to side of the district. The general contour of these lengthened hills individually is somewhat uncommon, although not peculiar to that part of the country. Each hill has a considerable degree of curvature, the convex side of which is uniformly presented to the west; the central part of that curve forms the highest point of their elevation, and they gradually slope towards the extremities of the segments which their curvatures form. Upon their eastern sides they rise with an equal acclivity of from 20° to 30° , the rapidity of which in some degree diminishes as it approaches the summits, where they are somewhat round-backed. Their central or highest points seldom or never exceed an elevation of from twelve to fifteen hundred feet above the sea-level, at which height their western faces very generally become abrupt and broken, and continue to be precipitous for a considerable depth under the lip of the hill; but these broken and abrupt faces seldom show more of the stratification than that of the trap, which in such situations shows its strong tendency towards the columnar form. At the base of these precipices a long and rapid slope of *debris* succeeds, which is frequently covered by vegetation. The troughs or hollows between the undulations generally form narrow dales or glens, but some few of them have a sufficient breadth to entitle them to the denomination of valleys; the most extended of these are the vales of Campsie and Fintry." This account of Col. Imrie is so far interesting as being, to the best of my knowledge, the earliest relating to the Geology of the district; but his descriptions of the strata are very meagre, and he takes no notice of the remarkable sections seen in several of the glens towards the western extremity of the range, and known as the "Ballagan beds." Belonging to the Wernerian school of geologists, he tells us that he saw no effects produced by the trap on the sedimentary rocks that he could attribute to the action of heat, and that any phenomena observable

at their junction might be referred to atmospheric agency. The now almost exploded theory of the aqueous origin of greenstone, basalt, &c., will find but little support here; while the modern or Huttonian doctrine, which attributes the formation of such rocks to the agency of fire, receives confirmation in numerous localities throughout the Campsie district, which Col. Imrie seems not to have visited, and where the masses of trap have left unmistakeable evidence of their volcanic origin in the altered condition of the sedimentary rocks with which they have come in contact. The only other point in Col. Imrie's paper to which I shall at present allude is his statement that the sources of the rivers Carron and Endrick are on the Muckle Ben. This is an error which will be apparent to any one who examines the district for himself, and induces me to believe that many of the Colonel's descriptions were drawn from a hurried visit, and afterwards written from memory; for the true source of the Endrick is in the Earl's hill to the north-east of Fintry, and no stream, taking its rise on the Ben, can possibly get into the river Endrick, as the valley of that river is divided from the valley of the Carron by a low ridge of hills; while at the same time the Carron sweeps round the base of the Ben on its northern and eastern sides, receiving as tributaries all the streams which descend from that hill, and flowing eastward to the Frith of Forth. On the other side of the dividing ridge which separates the Carron from the Endrick, the tributary streams of the latter river all descend to the Fintry valley, and discharge their waters into Lochlomond. The two valleys are thus shedding their waters in opposite directions; that of the Carron towards the east, and of the Endrick towards the west. Col. Imrie represents the Ben as the saddle of this part of the country; but as it has been shown that neither the Carron nor the Endrick has its source in this hill, we are inclined to regard, as the true saddle of the district, the elevated ridge immediately to the west of the road leading from Campsie to Fintry, and of which the Earl's seat is the highest point; for from this ridge three sets of streams derive their origin,—those on the north and west, which flow into the Strathblane and Fintry valleys, and become tributaries of the Endrick,—those on the south and south-east, which flow into the Campsie valley and form the river Glazert, a tributary of the Kelvin,—and those on the east, which flow into the Carron valley, and are the main feeders of the river Carron. We have thus on these heights, at no great distance from each other, the sources of streams that discharge their waters on two opposite sides of the

island, one set flowing to the German Ocean, and the other to the Atlantic.

The Campsie hills are in general more rugged along their outer borders than in the interior, their more precipitous faces for the most part looking to the west and north-west, while the opposite sides tail or taper off with a gradual slope. This phenomenon of "crag and tail," which is strikingly exemplified among the trap hills of the central districts of Scotland, I need hardly observe, has been attributed by geologists to the action of strong ocean currents setting in from the west and north-west during the "Pleistocene" or "Boulder drift" period, when the greater portion of the land which now constitutes the northern half of our globe stood at a lower level than it does in the present day.

On more closely examining the structure of the Campsie hills, we find that, with few exceptions, they are composed, in their upper stages, of tabular masses of trap separated by argillaceous deposits, which in some places assume the appearance of a sedimentary rock; and in others, by their amygdaloidal structure, present an analogy to rocks of igneous origin. These bands of looser material have apparently been ejected in the form of volcanic mud; or, it may be that they consist of *debris* derived, by the action of the water, from each successive flow of rock-matter, and gradually accumulated in the more depressed portions of the sea-bottom; in the latter case indicating long intervals of repose.

In their general relation to the stratified deposits with which they are associated, the igneous rocks of these hills seem to be for the most part overlying, especially towards their western extremities; and the various strata upon which they rest, and which form the base of the hills, having in many places been unable to resist the disintegrating effect of the drift currents, the superincumbent masses of trap, thus undermined, have been precipitated into the hollows below. These huge blocks of fallen rock give a rugged aspect to the hill-sides in numerous portions of the range: striking examples of this feature present themselves in the immediate neighbourhood of Lennoxtown. Nor are effects of the degrading agencies now in operation less apparent; for, although no icy sea now lashes these rocky scarps, from the rudely columnar form of the trap along the terraced steepes, water percolates into innumerable crevices, and, expanding when congealed by the frosts of winter, wedges off huge fragments, which tumble into the valleys below, with a noise like thunder. Above

Lennoxton, we have observed many of these detached rocks several hundred tons in weight.

Although, for the most part, the summits of the Campsie hills are more or less flat, many tracts presenting the appearance of an elevated table-land with steeply sloping sides, several isolated heights present themselves, as for example the Muckle Ben with the conical elevations in the Strathblane and Fintry valleys, as well as in the Kilpatrick division of the chain; these probably indicating the vents from which many of the tabular masses of trap were poured forth in molten streams over the bed of some ancient sea, in whose waters, and on whose shores lived and flourished organic forms of existence now enclosed in the solid rocks.

As we proceed in an examination of the physical features of the Campsie district we cannot but be impressed with the important extent to which these have been affected by denudation. It can hardly be doubted that Campsie valley owes its origin to this process, having evidently been scooped out by a powerful ocean current which has washed away strata belonging to the lower Carboniferous System to the extent of several hundred feet in thickness. On each side of the valley the nearly horizontal strata crop out abruptly, and are identical in their lithological character and order of sequence.

We have the same coals, limestones, and alum-shales, with their characteristic fossils; while, in the valley, bores have been put down in search of economic strata, and no trace of the missing beds has been found. By the action of these ocean currents that swept through this valley, as well as through that of the Kelvin, the strata of the North and South hills have been cut off from the main body of the great central coal field of Scotland, and now form outliers on its north-western margin. The sections across the valley, given at the end of the Paper, will convey to the reader a better idea of the amount of denudation than can be given by any written description.

Very little level ground is to be met with in the district, the surface of the valley being highly undulated, especially towards its eastern extremity, this circumstance being, no doubt, due to the unequal disintegration of the different strata during the denuding process.

In the narrowest part of the valley stands Dunglass, an isolated mass of trap, like some huge watch tower. This rock presents a good illustration of the columnar form of basalt, and the probabilities of its history are suggested to our mind when we direct our attention to the stratigraphical peculiarities of the immediate neighbourhood.

To the period of igneous activity in our island at the close of the Carboniferous era, the origin of this rock may be referred; and, although it is now exposed to the light of day, it may be supposed that the originally molten rock-matter of which it is composed struggled unsuccessfully to find vent through the Ballagan beds, and cooled under the pressure of the strata of which they consist. After the lapse of incalculable ages, the land settles down below sea-level, and the fierce currents of the "drift" period carry away the looser materials of the Ballagan beds, while the more compact core of basalt stands fast, and subsequently raises its head as a portion of the newest creation, with which we ourselves are associated. That the stratified rocks were borne upward to a considerable extent by the pressure of this mass of igneous matter, is evident when we remark that at the "Spout" of Ballagan, where the beds have been protected by the overlying trap of the hill, they crop out at an angle with the horizon; and, that, were they continued across the valley, they would obtain a high elevation over the remarkable height in question. This will be seen by looking at the section of the Ballagan beds at the end of the Paper, which represents them cropping out on the North hill towards Dunglass. Not only have the sedimentary strata of the district suffered from the effect of denudation, but even the harder trap has been much worn by the force of the watery element, and many of the boulders that occur in the drift of Campsie valley, as well as on the hill-side, have evidently been detached from the trap-rocks of the immediate vicinity. In the Strathblane district, there are several isolated areas of trap *in situ*, resting upon stratified rocks. These are now separated by considerable hollows, from the main body of the rocks to which they are allied, and with which they were, no doubt, formerly connected. A fine example may be seen at the south-eastern corner of the Mugduck reservoir, where the water is let off to supply the city. A patch of greenstone rests on sandstone, and forms the crown of a little eminence through which a wide circular opening has been cut in a vertical direction for the apparatus connected with the sluices. It will be observed that the greenstone has not been erupted through the sandstone, but merely overlies it, and is evidently an outlying patch which had been cut off from the principal mass of some trappean outburst, when the Mugduck valley was scooped out by the denuding current.

The diversified character of the Campsie district is chiefly due to the numerous flows and upthrows of trap that have taken place, and

even the verdure with which the hills are clad is attributable to the decomposition of the different rocks of the trappean family that so abundantly supply the inorganic materials necessary to the support of vegetable life; and, for grazing purposes, these hills are almost unrivalled in Scotland.

Trappean heights impress their own peculiar features on the landscape, and, although surpassed in wild grandeur by elevations of primary and transition rocks, their undulating and conical heights, and terraced fronts, possess a charm inseparable from themselves, while isolated outbursts, and traces of denudation cannot fail to be specially interesting to the geological observer. The characteristic features of trap scenery are well exhibited in the neighbourhood of Edinburgh and Stirling, as well as in the Bowling braes and Dumbarton rock in our own neighbourhood. In the Strathblane valley and surrounding country, the character stamped upon the scenery by the trap is particularly interesting. Standing on the western extremity of the South hill, as we have often done, let the geologist, or even the mere scenery-hunter, turn his eyes to the inviting prospect around him. To the north-east, he marks the long unbroken ridge of the Campsie range, with its boldly terraced fronts rising abruptly from the undulating tract of country on the south. Before him, in the middle of the valley, stands the conical hill of Dunglass, with its well defined columns of basalt. On the opposite side of the valley, Ballagan glen presents itself, with its many-ribbed strata capped by sandstone and trap. At its upper extremity may be seen the mountain stream, which gives rise to the river Blane, forming a cascade, (the "Spout") then leaping from ledge to ledge in its descent into the valley of Strathblane, through which it wends its way to mingle its waters with those of the Endrick. Farther west, in the valley of the Blane, the conical heights of Dunfoyne, Dungoyne, and Dungoich stand boldly up, like stepping-stones across the valley, while the lofty terraces of trap on the north hill, with the less elevated ridges opposite, contribute to impart a charm to the landscape not easily effaced from the memory.

Before closing our description of the Physical Geology of the Campsie district, we may notice another agent to which, no doubt, some of its more striking peculiarities are owing, viz,—Glacial action. All over the western portions of the South hill, at an elevation of 600 feet above the sea, large areas of sandstone-rock present a smooth and polished appearance, while, in many places there may

also be seen those grooves or furrows commonly attributed to the passing iceberg of the drift period, with its base of boulders ploughing the shallower sea bottoms, that now appear as elevated portions of land. These glacial *striæ* may be observed on the sandstone tracts on Craigmaddie moor around the "Auld wives' lifts," three ponderous blocks of the sandstone of the hill, one resting a-top of the two others, which are firmly imbedded in a marshy basin. Local tradition gives to three old wives the credit of having lifted, or transported, these remarkable stones to their present position—hence their designation. They are pronounced, however, by antiquaries versed in the pre-historic annals of our country to be *Cromlechs* marking an ancient burial-place; while not a few believe that they are Druidical altar stones. The markings to which we refer are also seen in Craigend moor, a little farther to the north-east, as well as on the flat surface of the trap-rock at the western extremity of the South hill, on the old road leading from Strathblane to Milngavie. The general direction of the channelled markings is from west to east, or from a few degrees north of west, according to the deflection of the hill-top, and in this respect they coincide with the course which a current would pursue in passing through the adjacent valley. Resting on the sandstone-rock, and perched at high levels on the hill-sides, solitary blocks of trap, and other rocks found *in situ* to the west and north-west, are often to be met with, and there can be little doubt that they played their part in this furrowing process, as they drifted along attached to masses of floating ice, while the low, flat summit of the South hill was still under the waters of a frigid sea.

Near the summit of Dunglass, a travelled boulder of quartzose metamorphic rock may be seen,

"Like a sea-beast crawled forth, which on a shelf
Of rock or sand repositeth, there to sun itself."

It has probably been dropped by some old-world iceberg that struck upon this sunken rock, as it was borne along by the current that rolled through the valley. Striated boulders of a large size are also to be met with in the drift in various localities over the district, especially in the Western-burn at Corrie, near Kilsyth.

As formerly stated, little level ground exists in the Campsie district, except some narrow strips along the margin of the Glazert and Kelvin. These *haughs*, or low tracts of land, are very little elevated

above the river-beds, particularly on the banks of the Kelvin, where the land is, not unfrequently, flooded for miles; and from the number of places termed "inches" along the banks of that river, it has been conjectured that, within the historic period, nearly the whole of this low land was constantly under water by the flooding of the river, the elevated patches called inches alone appearing above the surface. The river is now confined within its channel by artificial embankments, and it is only when in a highly flooded state that it spreads beyond its bounds. Wherever this flat land has been dipped into, it is found to consist of alluvium, stratified sand and gravel, not unusually resting on the Boulder-drift formation, being thus the most recent deposit of the district.

The student of Physical Geology will find much to interest him in the surface configuration of the valleys in the neighbourhood of Campsie, where he will recognise water as the last great agent to which they owe their present form, while, if he betake himself to the hill-sides, he cannot fail to trace the operation of that not less powerful natural force—fire, in the disrupted rock-masses by which the stratified deposits have been disturbed and overlaid. Since the land attained its present elevation, the physical aspects of the district do not appear to have undergone much change. The mountain streams are, no doubt, constantly eating their way deeper into the hills, while rains and other atmospheric causes have a tendency to wear down their higher levels; but these degrading agents operate so gradually, that the lapse even of many centuries would be scarcely sufficient to render their effect upon the general landscape perceptible.

We shall now proceed to notice the various sedimentary strata of the district, beginning with the lowest, or oldest. They are first seen in Campsie glen, on the North hill at the Clachan of Campsie, and may be traced from this point westward through the several glens in the Strathblane valley onward to the hills beyond the village of Fintry. In mineralogical character, this group of strata differs much from the carboniferous beds to the east of the Campsie valley, as well as from the subjacent Old Red Sandstone in the valley of Strathblane. It consists principally of thin layers of nodular limestone, seldom so thick or persistent as to form regular beds, except in one or two instances. These deposits alternate with thin beds of a bluish-grey marly shale, so numerous as to make up more than two-thirds of the group. Thin seams of red, and grey micaceous sandstone also occur, but more commonly in the lower division of the series. A thick bed

of light-coloured sandstone caps the whole group, and a similar rock is found also at the base between the limestones and the Old Red Sandstone. On the south-western flank of the Kilpatrick range, in Auchenreoch and Dumbuck glens, near Dumbarton, strata presenting the same distinctive features as the Ballagan beds are seen cropping out, and forming some very interesting sections. They are also found to be overlaid by the igneous rocks of the Lennox hills along their western extremities, so that they must, at one time, have covered a considerable surface of sea-bottom, before our submarine trap-rocks were spread over them. They form the base of the hill all around the Strathblane valley, and, from their thin-bedded character, present several interesting examples of stratification in the different glens in which they are exposed on the hill-sides.

In Campsie glen, the most easterly point where they are observed in the Campsie district, they are seen nearly in a horizontal position in the bed of the stream at the lower end of the glen, where they assume a yellowish hue from exposure to the weather, owing to the presence of a little iron in their composition. The limestones are here thicker than in any of the other localities in which we have observed them; and they alternate with thin beds of blue shale, and seams of red micaceous sandstone of no great thickness. The strata in this glen have been crossed by several trap-dykes nearly in a line between south-west and north-east; and they have had the effect of producing occasional faults and distortions of the strata. These may be observed in one of the sections on the east side of the glen below the first fall, the stratified rocks, at the line of junction with the trap, being, to a great extent, altered by heat; while several fragments of the limestone and shale are embedded in some of the dykes. Above the first fall, the strata appear on the west side of the glen, in a section of considerable thickness, forming the lofty bank below Craiglea Spout, and known as the "Lovers' leap."

At this point, overlying the strata, may be observed an immense mass of trap intercalated with bands of the reddish earthy deposit already alluded to. These trap rocks constitute the higher reaches of this glen, as well as the upper stages of the hills on both sides, and occur mostly in tabular masses of greater or less thickness, each successive bed forming the breast of one of the several waterfalls that may be seen in the course of the stream. These ledges consist principally of greenstone, in some cases porphyritic; but a soft claystone, not unusually of amygdaloidal structure, also occurs, and where this

softer rock is exposed to the recoil of the water under the falls, it is hollowed out into cave-like recesses. The line of contact between the igneous, and sedimentary rocks is clearly seen at this particular spot, where the deep brown hue, and scorched appearance of the shale, normally bluish-grey, are evidently owing to its contact with the originally heated mass of trap with which it is associated. The strata in this locality may be observed dipping into the hill in a north-western direction at an angle of about 10° , and, from the circumstance that the trap dips in the same direction, we may infer that the forces by which the strata have been disturbed from their originally horizontal position were not exerted till after they were overlaid by these outbursts of submarine lava.

The Campsie hills, at this point, appear to have been built up by successive overflows of igneous matter, consisting generally of alternate layers of compact greenstone, and the looser material, the peculiar characteristics of which have been already noticed.

Upwards of twenty of these trap beds can be counted in following the course of the glen from its lower levels to the hill-top, all differing more or less from each other in mineral character, and mechanical structure.

As Campsie glen is the last place to the eastward where the Ballagan beds are seen in the main valley, they, in all probability, underlie the higher carboniferous beds which appear in the valley still farther east. We are not aware of the Ballagan beds being represented in any other districts, with the exception that, in the Merse of Berwickshire, strata of a similar character, and holding the same relation to the Old Red Sandstone and superincumbent Coal Measures are said to occur. They seem to be wanting all along the southern borders of the Lanarkshire coal field, where we find the thick-bedded limestones of the Carboniferous System coming close upon the horizon of the Old Red Sandstone, without any great thickness of intervening strata, as exists along the north-western border. The next place to the westward, in the Campsie valley, where we observe these beds is Fin-glen. The lower reaches of this glen are wholly composed of trap rocks, which descend low into the valley, and consist chiefly of felspar-porphry, and compact greenstone, in thick masses. Proceeding upwards above the first considerable waterfall, we observe in the bed of the stream the thin yellow limestones of the Ballagan series cropping out; but the section of these strata in this glen is by no means extensive. They are intersected

by one or two narrow trap-dykes, presenting a similar appearance to those in Campsie glen, and running nearly in the same course. Near the second large fall, the sedimentary beds are cut off abruptly by a great mass of fine-grained greenstone. This small patch of strata seems to have been completely locked up in the trap, and but for the mountain stream that has cut its bed through that hard overlying rock, the position of these beds, in this locality, would have been unknown. From Fin-glen to the Spout of Ballagan, there is no section where the Ballagan strata are exposed; but it cannot be doubted that they constitute the base of the hill in many places between these two points. The finest section of these thin-bedded deposits to be met with in the district is the lofty escarpment in Ballagan glen, where upwards of 230 beds of limestone, sandstone, and marly shale are seen lying, tier upon tier, in a slightly inclined position, and capped by massive beds of white sandstone, and trap, the whole presenting one of the grandest specimens of stratification to be witnessed in this country-side. The beds dip into the hill in a north-westerly direction; and in the course of the stream, a little below the Spout, one or two trap-dykes may be seen intersecting the strata, while a fault occurs in the middle of the section. As in the localities already noticed, the overlying trap at the Spout has evidently flowed from some ancient volcanic crater far in the interior of the hill, without producing much disturbance of the strata; but here, too, the agency of heat may be traced in the indurated condition of the sandstones with which the trap has come in contact. Farther west in the Strathblane valley, the Ballagan beds are seen in the different glens that intersect the Strathblane hills, and in Spittal glen, a few minutes' walk to the west of the village of Strathblane, we first observe the junction of these beds with the Old Red Sandstone, which forms the floor of the valley. It would be difficult to point out the exact line of demarcation between the Old Red Sandstone and the strata in question, as the one formation passes almost imperceptibly into the other by various gradations without any change in the angle of inclination; and as it is here, so is it in Auchenreoch glen, where we can again trace the inoculation of the two formations.

Where the Ballagan beds approach the inferior formation they assume a reddish hue, and alternate with layers of yellow, and grey sandstones, and greenish shales; these being apparently the lowest members of the series. In the vicinity of Dunfoyne, one of the Strathblane hills, the Ballagan beds are again seen tilted up into various

positions by that isolated height of disrupted trap; while fine sections appear in the Corrie of Balglass, near the Earl's seat, on the Fintry side of the range, where they form the entire base of the hill towards the river Endrick. To the north of the river, they are again observed cropping out in the banks of Lernoek burn on the borders of Balgair moor; and they probably form the base of most of the elevated ground on the north side of Fintry valley in the direction of Stirling. From the thin-bedded character of the strata of this group, they are readily recognised wherever they are seen cropping out, and their thickness has been estimated at about 1000 feet; but this is possibly above the mark.

Auchenreoch glen is the last place to the westward where these beds appear, and this seems to be their western limit in Scotland, as far as yet observed; for the strata immediately beyond this point belong to the Old Red Sandstone formation, and occupy the vale of Leven.

Up to the last few years, the Ballagan group in the west of Scotland was considered to be non-fossiliferous; but in one or two localities fragmentary remains of plants, and scales of fishes, have been found in a grey, micaceous sandstone of the group, in which annelide tracks have also been discovered.

The fish-remains detected consisted of single scales of one or two ganoid species. These are very rare, and have hitherto been found only in Auchenreoch glen. Specimens were submitted to the late Professor Fleming of Edinburgh, who could not identify them with any known species, but conjectured that they might belong to the genera *Palæoniscus* and *Amblypterus*, both carboniferous fishes. The plant remains have been found at the Corrie of Balglass, the Spout of Ballagan, and in Auchenreoch glen. One species appears to have been a reedy plant, striated longitudinally like a calamite, but not jointed, and sending off spikes, or branching stems, at intervals. It was shown to the late Hugh Miller, and other geologists, at the last meeting of the British Association in Glasgow; but none of those gentlemen could determine the genus to which it belonged. Since that time the remains of other plants have turned up at the Spout of Ballagan; one of these is a long narrow stem of a species of *Lepidodendron*, and was to be seen embedded in a block of sandstone in the bed of the stream: another plant resembled the *Stigmaria*, but was in a less perfect state of preservation.

Two varieties of gypsum occur at Ballagan; one of a red colour in

veins intersecting the strata nearly at right angles; the other of a yellowish colour, and fibrous structure, appearing in laminæ between the beds. The fibrous variety of gypsum is also found in the same position in Auchenreoch and Dumbuck glens, where it is of a pure white colour; and it is said that ores of antimony and nickel have been discovered in very small quantities at the Spout; but we have never met with any in all our searches. The only mineral of a metallic nature which we have yet found is iron pyrites in crystallized nodules. The Ballagan limestones have never been successfully employed for any economic purpose, nor is it likely they will ever be wrought, so long as richer and more workable beds are to be found in the district. Several attempts have, at different periods, been made to find coal in the Ballagan strata; but, as might have been anticipated, invariably without success.

These strata lie considerably lower in the Carboniferous System than the Campsie coal and lime, which are understood at present to be the lowest workable coal and lime in the Lanarkshire basin, and we think there is little likelihood of any coal of available thickness existing in the beds in question to the west or north-west of the Campsie hills.

We have thus, at some length, endeavoured to describe the leading features of the Ballagan series, and to point out the localities in which they occur; and we shall now briefly consider the circumstances under which these strata appear to have been deposited, and the place they occupy in the geological scale. Taking the group as a whole, the Ballagan beds are a fine-grained deposit; the clays and sands of which they consist having evidently been produced by the gradual disintegration of the older rocks, and precipitated in tranquil waters without any paroxysmal effort of nature. Many of the sandstones are very finely laminated, while others exhibit a ripple-marked appearance, with the supposed worm-tracks previously alluded to. The undisturbed character of the deposit appears throughout its entire thickness, which, as has already been stated, is made up of numerous alternations of limestones, sandstones, and friable shales, the last constituting the great bulk of the formation. Setting aside the conjectural evidence of the worm-tracks, these rocks have yielded no fossils to prove their marine origin, so that it cannot be averred that they are not fresh-water deposits.

The Ballagan beds are interesting to the geologist from the circumstance that they are, undoubtedly, the oldest carboniferous de-

posits that we have in the west of Scotland; and while the fine views of stratification they afford in the different glens where they have been exposed cannot fail to delight his eye, his imagination cannot but be excited by the thought that he is looking upon the depository of the first traces of that wondrous vegetation that culminated in the later stages of the true carboniferous era, and now, in a mineralized form, exerts so important an influence on the economy of the world.

There are no sections in the valley of Campsie to the eastward, where we can trace the Ballagan beds upwards into the true fossiliferous strata of the Coal Formation; but their presence in the lower levels of the valley is proved by bores put down by the Alum Work Company several years ago in search of coal.

They were led to believe that some of the more valuable seams of coal which are found in the upper parts of the Carboniferous System might be found in the locality referred to; but had they paid the least attention to the outcrop of the strata on either side of the valley, they would have found sufficient evidence that the rocks into which they bored lay under the Campsie main coal and lime; and no coal of workable thickness is known to occur in the lower strata. A journal of one of the bores put down in the valley at Burnhouse, taken from the last Statistical Account of the parish, is given at the end of this paper, and it will be seen from it that, at the depth of about 33 fathoms under the surface of the valley, nearly 60 thin beds were penetrated. Of these, two were coal seams, one eleven, and the other fourteen inches in thickness. The beds under the second coal were of so unpromising a character that the search was abandoned at the above-mentioned depth. An adjacent bore near the level of the Glazert penetrated a series of porous beds, which, on being first tapped, discharged 120 gallons of water per minute; but after it had run for some time, it decreased to about 90 gallons, at which, I believe, it still continues to flow.

This Artesian well may be accounted for by supposing a fault or dyke crossing the strata of the valley, and intercepting all the water admitted at the upturned edges of the inclined beds containing it, which crop out in the upper reaches of the valley. The spring rises a few feet above the surface, and affords a permanent supply of water to the Alum Work Company. The first mentioned bore seems to have struck on the opposite side of the obstruction, for although it reaches to a greater depth, it has disclosed no extraordinary accumulation of water. These strata lie under the level of the river Glazert, and are

not exposed in any natural section, so that we know not to what extent they are fossiliferous. From their thin-bedded character, they seem to occupy an intermediate position between the Ballagan beds, and the strata overlying the thick layers of sandstone, which form the floor of the valley to the eastward. This sandstone rises along the South hill to the westward, becoming immensely developed in Craigmaddie moor, where some of the beds pass into a conglomerate consisting of quartz-pebbles loosely cemented in a matrix of fine sand. This sandstone, which crowns the South hill towards the west, is in many instances of a fine white colour; and, being compact and silicious, it would, no doubt, be useful for building purposes were it more accessible. In some localities, it contains numerous casts of the *Calamite*, *Sigillaria*, *Stigmara*, and *Lepidodendron*. These were formerly to be obtained at Calside quarry, as well as at the quarry under the Church-hill, and towards the western extremity of the South hill, at the quarries on the road leading from Strathblane to Milngavie; but as these workings are not carried on to any great extent at present, specimens of plants are now rarely obtained. The strata that lie immediately over, and under this sandstone partake much of the character ascribed by Mr. Page to the lower Coal Measures of Scotland, consisting, as they do, of beds of fresh-water or estuary limestone, with *Cypridinidæ* and remains of fishes; thin layers of bituminous shale, containing plants in addition to fishes; dark shales with bands of clay-ironstone, and two or three thin seams of coal. Associated with these deposits are also some inconsiderable seams of impure limestone and shales. These are of marine origin, and contain fossils characteristic of the lower Coal Measures. At the Linn of Baldernock, where one of the estuary limestones of the group makes its appearance, a thick bed of greenstone may be seen immediately above it in an interstratified position, and forming the roof of the limestone workings. The upper portion of the stratum of limestone contains a considerable quantity of iron pyrites, which appears to owe its origin to the presence of the overlying trap.

As there are few localities in the valley of Campsie where the rocks are sufficiently exposed to exhibit the ascending order of the strata immediately under the main coal and limestone of the district, it would be difficult to construct a section giving their true relative order; for, although they are found cropping out in the banks of the different streams, their original stratigraphical position has evidently been much disturbed by several faults which intersect them. In attempt-

ing, therefore, to point out the localities in which the lower fossiliferous beds of the recognised coal-strata are exhibited, and to present them in their relative ascending order, we cannot vouch for the absolute correctness of our conclusions.

CRAIGENGLLEN BEDS.

Next in order above the strata penetrated by the bore near the level of the Glazert, we may place the Craigen Glen beds. This glen is situated on the southern flank of the South hill of Campsie, nearly in a direct line south from Lennoxton, and the strata which are found cropping out in the banks of the rivulet that flows through it present an abundance and variety of organic remains. Indeed, the fragmentary stems of *Encrinites* were at one time so plentifully strewn on the bed of the stream that it was known among the old people of the district as the "Witch-whorl burn;" but we rather suspect that the sound of the geologist's hammer has broken up the wierd establishment in which these "wheel-stones" were supposed to have been manufactured, for their remains are getting rather scarce. The strata exposed in Craigen Glen consist of thick-bedded white sandstone, one or two beds of a silicious sandstone containing plant-remains, and dark-blue shale interspersed with bands and nodules of rich clay-ironstone, which were formerly worked to some extent. Bands of impure limestone, some of them of marine origin, and others of a somewhat oolitic structure, but destitute of fossils, also occur in this series, with a thin bed of highly bituminous shale containing the remains of plants and fishes. The sections in this glen are by no means extensive, and the organic remains are found principally in loose blocks of limestone and shale which had been tumbled into the bed of the stream during the working of the ironstone. The strata dip to the south-east, and a dyke of a purplish-coloured claystone seems to cut them off in the lower part of the glen, as we do not find any of the fossiliferous beds exposed beyond this point. This dyke is seen in the bed of the stream near the middle of the glen, but it cannot be traced to any great distance beyond. A little farther down the hill to the westward, in the bed of Glenwyne burn, we also find the Craigen Glen strata cropping out, and dipping at a high angle to the south-east. Fully half a mile more to the west, they are again exposed in the banks of the same burn; but we are not aware of any other locality in the district where

they make their appearance. The Craigenglen beds are interesting chiefly for the abundance of their organic remains. Many of the shales are, for the most part, composed of crushed shells, corals, and stems of Crinodea. In these shales, however, shells are often found in a very perfect state, their original texture being wonderfully well preserved; but they never occur in the form of casts, as they do in the limestones and ironstones.

The bivalve shells are generally separated into single valves, but complete specimens of some species are frequently discovered in the strata. Univalve shells, belonging to several genera and species, occur here in greater numbers than in any of the other beds of the Campsie district. Several of the Craigenglen shells still retain traces of colour, one species of *Orthoceras* exhibiting well-marked *chevron*-shaped bands traversing its surface. Since the working of the ironstone was given up, specimens of fossils are not so easily obtained in Craigenglen as formerly, and, owing to the frequent visits of many of the members of our society, the banks have been pretty well searched: still, good specimens may be procured by digging out, and splitting up, some of the slabs of soft shale. From the list of the various organic remains found in the Campsie district, given at the end of this Paper, it will be seen that the Craigenglen beds have yielded a greater number and variety than any of the other beds in the district.

BALGLASS BURN BEDS.

Passing over to Balglass burn, on the north side of the South hill, we discover a variety of organic remains in blocks of dark-grey shale and nodules of clay-ironstone lying in the bed of the burn, which is so shallow that no section of the strata from which these fossiliferous rocks have been derived is exposed to view. It is evident, however, that we are here close upon the outcrop of the beds, as the blocks of shale and ironstone found embedded in the clay of the banks of the stream are not at all water-worn. From the shale, have been obtained *Trilobites*, teeth of the *Psammodus porosus*, finely preserved arms of the species of *Encrinure*, figured by Ure in his "Natural History of Rutherglen and Kilbride," Plate XVIII., Figure 8; while both the ironstone and shale have yielded the remains of *Serpulites*, and various other shells—chiefly Brachiopoda. There is no other place in the district where the same strata crop out; but we are inclined to think, from the position of the shale on the hill-side, that the beds are a little higher than the Craigenglen strata on the other side of the hill.

MILLBURN BEDS.

Next in ascending order may be placed the Millburn beds, which appear on the north side of the Campsie valley to the east of the church, and consist of a series of fossiliferous shales, clay-ironstones, and limestones, cropping out in the bed of the burn. One of the limestones of this series seems to be of fresh-water origin, containing, in addition to plants, the remains of *Oypris*, a genus of extremely minute crustacean, many species of which inhabit our modern ponds and lakes; while another, of little thickness, is evidently marine, being charged with shells and corals. We lately discovered the *Productus giganteus* in the latter limestone; and from the circumstance that that shell is not found in any other locality throughout the district, it becomes comparatively easy to indicate the relative position of the Millburn beds. The *Productus giganteus* is a good index-shell in the Lanarkshire coal-field; and from its occurrence in the beds in question, they may be pronounced to be about 22 fathoms under the Campsie Main Coal and Limestone. Teeth, spines, and scales of fishes are sometimes to be met with in these beds; and in one of the calcareous shales, the *Lingula squamiformis* is abundant. *Rhynchonella pleurodon*, and *Athyris ambigua* abound in the clay-ironstone; and *Streptorhynchus crenistria*, *Pinna flabelliformis*, with one or two species of *Aviculopecten*, are common in the shale above the limestone. No economic value is at present attached to any of the limestones of this locality. In other parts of the Lanarkshire coal-field, however, the *Productus giganteus*-limestone is a good workable stratum; but here it has thinned out to not much more than a foot in thickness.

BALGROCHAN BEDS.

Near the village of Lennoxton, on the road leading to Fintry, in the banks of Balgrochan burn, another bed of limestone, associated with fossiliferous shales, may be seen cropping out. These strata also underlie the Main Limestone of the Campsie district, which makes its appearance at a higher level upon the hill-side. The limestone of Balgrochan burn is made up of a vast accumulation of crushed shells; *Spirifera duplicicosta*, and *Productus semireticulatus* (var. *Martini*), being the most common. The material which cements the shells together is a dark-grey clay containing very

little calcareous matter; so that the rock is not at present valued as a limestone. Many of the clay-ironstone nodules found in the shale under the limestone have yielded very perfect specimens of *Athyris*, *Rhynchonella* and *Discina*, as well as various other species of Brachiopoda. Although the limestone of this locality is made up of shells to so large an extent, very few complete specimens can be obtained. The *Producti* and *Spiriferæ* are represented by single valves, often much crushed; and from the shale above the limestone, *Fenestella plebeia*, and *Cerriopora interporosa* have been obtained. Encrinal remains, which are common in the Craigenglen, and Balglass burn beds, are exceedingly rare in those of the Millburn, and Balgrochan.

It is very probable that the shelly limestone of the Lanarkshire coal-field, which ranks about 16 fathoms under the Main Limestone, and about 6 fathoms above the *Productus giganteus* limestone, is the equivalent, or representative of the Balgrochan beds.

CAMPSIE MAIN COAL AND LIME.

Proceeding in ascending order, the next strata in the Campsie district to which we come, and of which we have any certain knowledge, embrace the Main Coal and Limestone, and form a well-defined group in the Lanarkshire coal-field; both the coal and limestone being very persistent over a great part of the basin. With them a rich bed of alum-shale is frequently associated, and has been long worked with profit at Hurlet and Campsie.

In the table of strata appended to Mr. William Moore's lately published pamphlet on "The extent, and supply of the minerals in the coal-fields of the west of Scotland," the Main Limestone is placed 702 fathoms under the Upper Coal; and if we take into account the thickness of strata above that seam, with that of the beds known to occur under the Main Limestone, adding the Ballagan series, we shall probably have not less than 6,000 feet of deposits representing the Carboniferous System in this part of the island.

In the Campsie district, immediately under the Main Coal, there is a bed of limestone between four and five feet in thickness. It has been termed the "White Limestone," being of a light cream-colour after exposure to the weather, and, like the limestone at the Linn of Baldernock, formerly mentioned, is apparently also of fresh-water, or estuary origin, abounding with the fossilized shields of *Cypripis*. This limestone is said to contain about 83 per cent. of car-

bonate of lime; but it has never been much worked in the district. Three or four beds of this Cypris-limestone are found at intervals among the marine strata under the Main Limestone of Campsie; and they have hitherto yielded no salt-water fossils, which are common enough in the strata immediately above and below them. We have not been able to discover evidence of the existence of any of these estuary or fresh-water limestones out of the Campsie district; and it is possible that they are merely a local deposit, soon thinning out. Alternations of fresh-water, and marine strata are said to be of common occurrence in the Lower Coal Measures of the east of Scotland; but the Campsie district is the only locality in the west where we have yet been able to trace, in these deposits, such indications of a diversity of origin. It is more than likely, however, that they occur also in the Kilpatrick district, a little to the south-west of the Lennox hills, where similar strata are found.

The alterations in the character of the strata seem to indicate successive elevations and depressions of the area of deposit—at one period the bed of an estuary or lake, and at another a portion of the sea-bottom; and if the fresh-water limestones could be traced gradually assuming the character of marine deposits in other localities, interesting evidence of their estuarine origin might thus be obtained. One of the Cypris-limestones in the Mill-burn contains a slight percentage of iron, and is in close contact with another bed of limestone of the same colour and composition, which contains *Producti*, *Spiriferæ*, &c. May not these two impure limestones have been derived from the same source, but deposited under such a difference of conditions as we have suggested? We can imagine the Cypris-limestone forming in fresh-water at the *embouchure* of a river, or in a lake little elevated above the sea-level. A slight depression of the land takes place: the briny waters of the ocean rush in: fresh-water forms of life die out, and a marine *fauna* takes their place; while the river, deriving its spoils from the same source, continues to discharge the sedimentary matter of which these limestones equally consist.

The Main Coal of the Campsie district is separated from the White Limestone beneath by about seven inches of “underclay,” and from the overlying Main Limestone by the alum-shale, which is about two feet thick on the South hill, while on the North hill, on the opposite side of the valley, it attains a thickness of about fourteen feet; but in the latter locality it is rich in alum only in the lower portion of the bed, which is contiguous to the coal. The coal averages three feet

eight inches in thickness, but is not of superior quality, especially along the North hill, where it is much deteriorated by thin bands and nodules of iron pyrites. It was formerly much used in the district for household purposes; but now that the finer coals of the upper division of the Carboniferous System are imported by the railway, the local deposit is drawn upon chiefly for use in the public works, and for burning the limestone.

The Main Limestone of the Campsie district has long been famous as a building material when calcined; and if we may judge from the numerous old workings along the out-crop of the bed on the North and South hills, it must have been quarried for many years. In the parish there is a tradition that the lime used in the erection of the High Church of Glasgow was taken from the South braes of Campsie.

As previously stated, the South hill is an outlier of the lower beds of the Lanarkshire coal-field. Nearly all round the hill the coal and limestone crop out, and these, with the alum-shale, have been worked from both flanks in the line of strike of the beds, so that the hill is now undermined from side to side. A greenstone dyke traverses the South hill from east to west, and is probably a spur or branch of a dyke of the same composition, which extends from south-west to north-east, and intersects the strata at Milton-of-Campsie.

The sedimentary rocks on each side of the dyke are not much displaced, but the coal in contact with it is greatly altered,—evidently by the agency of heat.

Near Barriston, on the southern slope of the South hill, extensive faults occur in the strata; “downthrows” at several stages causing the coal and limestone to occupy a low position on the hill-side.

Many patches of the coal in this locality are converted into anthracite, called by the miners “humph,” or “blind” coal; and on the North hill, at Schiliengow quarries, where a narrow dyke of a light coloured trap passes through the strata, the coal in contact with it has been burned into a cindery mass, and in various other localities along the hill-side where these igneous intrusions have taken place similar effects have been produced.

The principal fossils of the Campsie Main Limestone are *Productus*, *Spirifera*, and *Athyris*; but it cannot be pronounced a rich fossiliferous deposit. As far as has yet been discovered, it contains certain shells peculiar to this division of the Campsie strata, viz.:—*Productus mesolobus*, *P. aculeatus*, and *P. fimbriatus*.

Nautilus ingens, and *Actinoceras giganteum* are sometimes met

with, as also fragments of a large crustacean—*Eurypterus Scouleri*. In the ironstone and shale under the limestone, *Spirifera Uri* and *Productus longispinus* are very abundant; and in the shales above the limestone, remains of *Trilobites*, shells, and *Placoid fishes* have been found. Specimens may be picked up wherever the limestone has been quarried along the hill-side.

The shale above the Main Limestone is 22 fathoms in thickness, and parted at intervals by thin beds of clay-ironstone. It contains in its upper parts very few organic remains; and is overlaid by another bed of shale, associated with three or four workable beds of clay-ironstone, and a stratum of coarse limestone about six feet in thickness, supposed to be identical with the “Hosie” Limestone, which crops out along the southern limit of the Lanarkshire coal-field. This limestone has never been much worked in the Campsie district; but the ironstone which underlies it was mined to a considerable extent at Barriston and some other points on the South hill. The clay-bands at Hurlet, and Crossbasket seem to hold nearly the same stratigraphical position as those found in this division of the Campsie deposits, and some of the shale-beds contain organic remains nearly identical. The shale immediately above the coarse limestone of the Campsie section abounds with various species of fossils in an excellent state of preservation. Some of these are peculiar to this stage of the Campsie beds, as will be seen from the list; *Orthoceras*, *Pleurotomaria*, *Nucula*, *Leda*, *Spirifera*, and *Chonetes* being the most common; but *Dentalium*, and a small shell, which appears to be the cast-off integument of a large species of *Cythere*, are also found in the same deposit. The shales accompanying the ironstone under the limestone contain few fossils; but on the upper surface of one of the ironstone-bands, numerous specimens of fish-remains are to be seen. These consist of scattered fragments of scales, spines, and teeth, with one or two species of *Cypris*. From the absence of marine shells in these beds, we are inclined to consider them of fresh-water origin. They are the highest strata of the Campsie series in which we have detected remains of *Cypridinidæ*. The fossiliferous strata of the district above these ironstones are more decidedly of marine formation, and belong to the Mountain Limestone group of the Scotch Coal Measures. During the sinking of a pit at Barriston, a few years ago, another bed of limestone was discovered near the surface, and does not seem to have been noticed in this district before. It is of marine

formation; but its fossils differ from those of any other limestone in the district. It contains a small percentage of iron, is very hard, and the upper part of the bed, to some extent, takes the form of the "Cone-in-cone" Limestone. The relative position of this bed is fully 30 fathoms above the Hosie Limestone. The strata at Barriston, although at a lower level, are higher in the series, than those on any other part of the South hill. This circumstance is evidently owing to the "upthrows" and "downthrows" that accompany the faults in this locality, as well as to the more extensive denudation to which the higher portions of the hill appear to have been subjected; and from the same combination of causes, it also happens that in the denuded "upthrows," lower beds are brought nearer the surface; for, in the pit where the new limestone was discovered, the Campsie Main Coal is fully 60 fathoms under-ground, while in another pit in an adjoining field, and nearly on a corresponding surface level, the same coal was worked at a depth of not much more than 12 fathoms. These displacements of the strata, although common in our coal-fields, are seldom indicated by corresponding irregularities of the surface, the more elevated beds having been reduced by subsequent denudation. The annexed section will illustrate such a condition of the strata as that referred to. A fault is represented, accompanied by a downthrow of the beds, *a*, and an upthrow of those marked *b*. The strata above the dotted curve being denuded, the deposit, *c*, is brought nearer the surface on the side of the upthrow, as shown by the perpendicular lines, *d*; while the beds of the downthrow are higher in stratigraphical order.

As has already been observed, the limestone found at Barriston is fully 30 fathoms higher in the series than the Hosie Limestone. Between them a bed of blue shale intervenes, accompanied by one or two bands of clay-ironstone, and a thick-bedded yellow sandstone. These deposits form the most elevated portion of the South hill towards Mount Hooly. A portion of the same bed of sandstone underlies the trap of the North hill to the east of the village of

Lennoxtown. At Ferrot's quarry, this sandstone is thrown into a vertical position; and a little farther westward the beds are much crushed and contorted by their proximity to the igneous rocks.

CORRIE-BURN BEDS.

To the east of the quarry referred to, the strata which form the base of the hill take rank above the Main Limestone, and below the yellow sandstone which underlies the trap, but they are not exhibited in any section worthy of notice. Still farther to the east, we arrive at Corrie, a ravine terminated by a *cul de sac*, as the term implies. It is threaded by a stream, the Western-burn, which enters it by a precipice of considerable height. In this neighbourhood we find some very interesting sections of strata exposed in the banks of the streams that descend from the hill. The beds here are very different in their character from those in the valley of Campsie to the westward, and seem to belong to a higher stage of the Carboniferous System. They are the best example of the Mountain Limestone group we have in this part of the country.

The lowest strata of the Corrie consist of thick beds of shale, with thin bands of impure clay-ironstone, and these may be observed cropping out at the entrance of the glen. They are succeeded by a thick-bedded yellow sandstone, which is, perhaps, a continuation of the yellow sandstone immediately under the trap in the Campsie valley. Beds of shale succeed, with thin bands and nodules of clay-ironstone, and a coal-seam of inferior quality, which is wrought and applied in burning the limestone. Above the coal we have again beds of shale, with clay-ironstone, overlaid by a bed of coralline limestone about three feet in thickness. The corals of which the great bulk of this stone is composed have evidently lived and died upon the spot they now occupy. We are here looking upon a portion of the bed of an ancient sea, on which myriads of these beautifully organized forms were rooted, to live their little day and perform the humble functions assigned them in the scheme of nature. Gradually and silently, however, they are enveloped in a muddy sediment, and await their higher mission to speak to the earnest student of geological science in a language which he alone can interpret. The clay which fills up the spaces between these zoophytes is of a greenish colour, containing little calcareous matter, and the corals themselves are the *Lithodendron fasciculatum*, of which the stratum is mainly composed, and

small masses of the *Lithostrotion Portlocki*, with one or two species of *Cyathophyllum*. Overlying the coralline limestone, we find a bed of light-coloured friable shale, abounding in several species of the *Productus* and other genera of Brachiopoda, with stems of Encrinites and fragments of corals. A little higher on the hill-side these beds are succeeded by thick beds of hard calcareous shales, and several bands of limestone, one of which, about five feet thick, is being extensively wrought. The calcareous shales contain several species of fossils. *Aviculopecten* is very abundant in some of the beds, and in various thin limestones, large specimens of *Cyathophyllum* may be found, while *Lithodendron junceum* is also common. In some of the finely laminated shales which alternate with the limestone, very good specimens of various species of *Fenestella*, *Glauconome*, *Vincularia*, &c., are found in an excellent state of preservation. Fish-remains are rare at Corrie, the teeth of *Cochliodus*, *Helodus*, and *Petalodus* being all that have been detected in the strata up to the present time: *Trilobites*, *Serpulæ*, and *Serpulites*, are sometimes met with. The marine limestone and calcareous shales are overlaid by a very thick bed of dark-grey shale containing numerous bands of clay-ironstone-nodules arranged in regular layers coincident with the plane of stratification. These nodules are finely exhibited in an interesting section at the Eastern-burn in a cliff nearly 100 feet in height. This deposit is of marine origin, and contains, thinly dispersed through the strata, remains of *Orthoceras*, *Goniatites*, *Nucula*, *Spirifera*, &c. The beds in the Eastern-burn are the highest in the Corrie section. The general dip of all the strata in this locality is from the hill toward the south-east, the beds flanking the hill, lying at a high angle against the trap which forms its summit. The junction of the trap with the sandstone may be seen in the bed of the Western-burn above the first waterfall, as well as at several other points along the hill-side between Corrie and Campsie. Above the waterfall at the upper extremity of the glen, and at a high level on the hill-side, there may be observed a considerable body of the Corrie strata lifted up and surrounded by the trap; and it is interesting to notice the altered condition of the various stratified rocks in the more immediate neighbourhood of the igneous mass. The fossiliferous limestones are changed into a mottled marble, which retains no trace of organic remains; the shales are converted into jasper or flint, and in some places into a black hornstone. Of all the strata we have observed in the district in contact with trap, these are the most altered.

In the banks of the Western-burn, several trap-dykes are seen crossing the strata; and near one of these, a bed of impure limestone, thrown up on edge, has been long known to have yielded small crystals of galena. Farther down the stream, on the western bank, several horizontal beds of trap occur, appearing to dip under the strata to the south-east, and consisting of a thin bed of greenstone, with beds of a reddish claystone, and tufa, which have all a stratified appearance; and it is very probable that these igneous beds may be contemporaneous with the stratified rocks they accompany. In the same stream, a very fine section of Boulder-drift appears, and is well worthy of notice. It seems to have abutted upon a great wall of trap, which forms the eastern boundary of the stream below the large fall. The stream has cut a channel for itself between the trap and the drift, so that we have now, on the one side, a lofty wall of igneous rock, and on the other, a steep, sloping bank of Boulder-drift. From the latter, have been derived the numerous rounded blocks of stone that occupy the bed of the stream. Some of these are of enormous size, and exhibit, on their flattened sides, very distinct glacial *strice*. In the trap at Corrie there occurs a vein of sulphate of Baryta of a pale reddish colour, and in it traces of copper and silver are to be found. This vein passes through the hill from north to south in a position nearly vertical; and it was found, after an attempt once made to work it, that it was not sufficiently metalliferous to be remunerative.

To the geologist, Corrie is one of the most interesting localities in the Campsie district, exhibiting, as it does, sections of Boulder-drift, igneous rocks, and carboniferous strata, all highly instructive, and worthy of a visit. We can only point out the more remarkable features of this locality without entering into lengthened details, in order to keep our Paper within due limits; and the only other beds of carboniferous strata we shall at present notice, as occurring in this district, but of which no natural sections are exposed, are those known to exist in the grounds of Balquarhage on the south-eastern termination of the South hill, Campsie.

BALQUARHAGE BEDS.

A pit was sunk in these beds a few years since for the purpose of working the coal, and the strata passed through were dark-blue shales, calm-limestones, in two beds, with a thin parting of shale between

them; shale, with ironstone nodules; bituminous shale; coal, and dark-grey arenaceous shale full of plants. The coal is about 13 fathoms under the surface where the pit was opened, and is five feet in thickness. It was not considered of good quality, being difficult to kindle, and leaving a large amount of ash of a whitish colour; but it is much less sulphurous than the Main Coal at Campsie. The working of the coal was soon abandoned, as it could not compete in the market with that of the upper part of the system, but it is possible that it may yet be wrought when coal becomes less abundant in this country. The Balquarhage beds dip slightly towards the valley of the Kelvin, and the coal may possibly be found of better quality farther from the hill, as, in the opinion of some of the miners, the pit was sunk not far from a "trouble" or fault in the strata. The sinking of this pit was very much retarded by the great quantity of water that poured into it from the limestone strata passed through; and from the same cause, the expense of sinking was so great that the lessee soon became bankrupt, and no one has had the courage to make any farther investigation of these beds. Some of the shales, and the limestone above the coal, are very rich in shells, stems of *Encrinites*, remains of *Trilobites*, &c. In their general character, and fossil contents, the Balquarhage deposits correspond more with the upper marine limestone series of the Lanarkshire coal-field than any other bed I have yet examined; the limestone, in its composition, and organic remains resembling that found at Moodiesburn, Bedlay, and several other places, while the coal lies about the same distance under the limestone as that found at those localities. The Balquarhage beds lie to the south of the great greenstone dyke, formerly noticed, which intersects the strata of the district from north-east to south-west, and appears to throw down the Campsie coal and limestone to a great depth—it may be upwards of 400 fathoms, if we have determined rightly the position of the Balquarhage limestone as corresponding with that of the Bedlay limestone, which is placed in Mr. Moore's section fully 430 fathoms above the Campsie Main Limestone; so that all the beds on the South and North hills, and in the valley of Campsie, must belong to a much lower stage of the Carboniferous System than any on the south of the greenstone dyke. In the railway cutting at Milton-of-Campsie, a great upthrow of the strata, occasioned by this dyke, is exposed, and if a corresponding effect has been produced along its course through the South hill, a considerable difference between the level of the strata on either side may be ex-

It is extremely difficult to fix the horizon of the Mountain Limestone group in the Campsie district; for several species which have a wide range are found passing through the whole of the strata, from our lowest to our highest marine limestones and shales. In the Lanarkshire coal-field, the middle division of the series extends from the Gare, Bedlay, and Moodiesburn limestone down to the horizon of our thickest marine limestones; and from this point, to the horizon of the Old Red Sandstone;—the strata of coal, shales, and fresh-water, and marine limestones, being included in the Lower Coal Measures.

Till the species peculiar to each group be defined, it will be very difficult to determine the order of the strata by their fossils alone, as very many species have a wide range, being found in various members of the series; and it is to be hoped that the Geological Survey will throw additional light upon the position of the carboniferous limestones of Scotland.

The igneous rocks of the Campsie district consist of several varieties of trap, such as basalt, greenstone, claystone, clinkstone, and tufa, with porphyries, amygdaloids, &c. The principal mass of trap-rock forms the upper stage of the range of hills on the north of the district, and may be studied, in all its phases, in the numerous precipices, glens, and natural sections along their borders. The relation between the igneous, and the sedimentary rocks of the carboniferous period may be conveniently studied at Corrie, Shieldsburn, Campsie glen, Fin-glen, and the Spout of Ballagan, all on the North hill; while, on the South hill, the strata may be seen in contact with the trap at Milton railway station, the greenstone quarry near Kinkeld, the Linn of Baldernock, and the Mugduck reservoir. The trap-dyke that passes through the strata of the South hill appears at the surface in Craigend moor, and from this locality it is traceable to the westward along the low, flat summit of the hill. In Craigmaddie moor, it rises above the sandstone to a height of from 20 to 40 feet, and forms a natural boundary-wall between the two moors, the softer sandstone on each side having been worn away by denudation.

The trap of the Campsie district yields fewer of the crystalline minerals than many parts of the Kilpatrick hills to the south-west, where, from veins and cavities, many fine specimens of *Thomsonite*, *Prehnite*, *Stilbite*, *Heulandite*, *Harmatome*, *Analcime*, &c., have been obtained.

In the cutting of the tunnel for the waterworks between Strath-

blane and Milngave, a very interesting series of amygdaloidal clay-stones, or felstones as they are sometimes termed, was passed through, and these have yielded many of the above minerals, with what appears to be a mineral new to the district, brought under our notice by one of the members of this society. It is of a light chocolate colour, and in its crystalline structure resembles *Laumonite*, consisting of an aggregation of small crystals radiating from a centre: it is also brittle in its composition.

We shall now shortly consider the relation which the Campsie strata hold to the other parts of the Lanarkshire coal-field. We seem to want in this district, all the upper coals and ironstones, with their accompanying strata extending from the Upper Coal to the first, or upper marine limestone series, the higher coal-beds being found to occupy the more central parts of the basin; so that the Campsie strata may be said to embrace only two of the three divisions into which the Scotch Coal Formation has been divided by geologists, viz:—Upper Coal series, Middle, or Carboniferous Limestone series, and Lower Coal series, the two last divisions including the whole of the Campsie strata from the Upper Limestone down to the base of the Ballagan series. The fossils of the Campsie beds are identical in species with those found in the corresponding strata of other parts of the Lanarkshire Coal-fields, very few, if any, being peculiar to the district, and they are all nearly identical with the species found in the Mountain Limestone and Lower Coal Measures of England and Ireland, a circumstance which will be observed by reference to the list of species found in the district. Indeed, it has very rarely happened that a species has been found, which had not been previously discovered in the Mountain Limestone, or Lower Coal Measures of other countries, so that we have in these fossils a good index to the position of the economic deposits of the Lanarkshire coal-fields, many of our most valuable coal and ironstone beds being found in strata considered the equivalents of the Mountain Limestone group. It was at one time much doubted by some geologists whether we had any strata in Scotland of the age of the Mountain Limestone series, but a belief in the absence of such strata is, I think, now nearly abandoned, besides it is evident that we have in our lower Coal Measures, a vast series of beds very much lower in the Carboniferous System than the mountain limestone, and characterised by its own peculiar strata and fossils. In Scotland, the carboniferous, or Mountain Limestone group has been split up into a number of beds of

greater or less thickness by interruptions of muddy sediment, sometimes forming shale-beds of great depth, in other instances vast beds of sandstone occur, while, at intervals we meet with a seam of good workable coal, which, perhaps, may indicate some ancient land-surface on which grew the vegetable matter forming the coal. All these alternate beds of coal, ironstone, limestone, sandstone, and shale, found in the middle division of our coal-field have, no doubt, been produced under circumstances different from those which produced the thick-bedded limestones of England and Ireland, which have been deposited apparently in seas less subject to the deposition of muddy sediment, and probably more favourable to the existence of marine animals, from whose calcareous *exuviae* so large a proportion of these limestones has been derived: yet the life of the whole period presents a striking analogy in all countries where synchronous strata have been recognised.

Jasper of a red, and yellow colour, are found in the trap of the Campsie range, as for example at Corrie, Fin-glen, the Spout of Ballagan, and in the rocky fronts of the Strathblane hills.

In the last-named locality it occurs in considerable masses in a thick deposit of earthy matter which alternates with the denser beds of trap of which the hills mainly consist. Small veins of a green jasper may be seen running through the basaltic rock of Dunglass, on various parts of the hill. Scotch pebbles, or agates, are not very common, still specimens of considerable beauty may occasionally be found on various parts of the hill. Besides Dunglass, there are several examples of the columnar form of trap. At Slough Neach, on the hill-side to the north-east of Lennoxtown, an escarpment about 100 feet in height consists of long slender columns of a buff-coloured compact claystone, in some instances assuming a greenish hue. On Craighalzean hill, in the Strathblane district, a section of well-developed columns of basalt gets the name of the "Pillar Craig;" and on Craigton hill, on the north side of the Fintry valley, are the "Pillars of Dun" consisting of the same rock.

In closing these observations on the more prominent features of the Geology of the Campsie district, I may say that much remains to be done with the view of determining the relation which some of the strata bear to other members of the great coal-field of Scotland. It is only by a more thorough examination of these beds and their organic contents that we can arrive at a knowledge of their true

bearing in the Carboniferous System; and I trust that this Paper will be the means of stimulating some of the members of the Glasgow Geological Society to a more complete investigation of the coal-fields to the north of the city.

NOTE.—The table of fossils already found, with the strata that have yielded them, will be useful to future collectors. The only class of carboniferous fossils of which the species have been satisfactorily determined are the *Brachiopoda*; and these have been fully wrought out by that distinguished member of our Society, Thomas Davidson, Esq., F.R.S., Brighton. The different genera and species of the other classes of the sub-kingdoms represented in the district are also given, as far as we have been able to make them out from the few works we have had an opportunity of consulting, and by comparing them with named species in various collections. Those which we have not been able decidedly to identify are marked doubtful (?). Palæontologists have much to do among the other groups of Scottish carboniferous fossils before a complete classification can be made; and we hope that some one able for the task may enter the field, and do for the other classes of extinct life, what Mr. Davidson has done for the *Brachiopoda* of our carboniferous deposits in his splendid monograph.

No. 1.—Journal of Strata passed through in sinking a pit near the summit level of the South hill, 450 feet above the river Glazert.—Copied from the last *Statistical Account of the Parish*.

	Fh.	Ft.	In.	Under the coal the following beds crop out on the hill above the level of the Glazert:—	Fh.	Ft.	In.
Sandstone, out of Section, thickness not known	...			Brought forward,	40	3	5
Blue shale, first bed pas- sed through	-	11	1 0	Under-clay	-	-	0 0 7
Clay-ironstone	-	0	0 7	White Limestone (fresh- water)	-	-	0 4 6
Blue shale	-	3	2 2	"Kingle," (a hard, silicious sandstone)	-	-	0 4 6
Black, or Hosie Limestone	1	0	0	Shale	-	-	1 1 4
Shale, with four bands of Clay-ironstone	-	1	0 0	Limestone (marine)	-	-	0 0 10
Blue shale	-	22	2 0	"Fakes," (a sandy shale)	-	-	0 0 10
Campsie Main Limestone (marine)	-	0	4 0	Alum shale	-	-	0 0 9
Alum-shale	-	0	2 0	Parrot coal	-	-	0 0 4
Main Coal	-	0	3 8	Common coal	-	-	0 0 10
				Fire-Clay in two beds, parted by Kingle	-	1	2 6
Carry forward,	40	3	5			45	2 5

No. 2.—Journal of Bore put down at Burnhouse near the level of the Glazert. The beds passed through in this bore lie under the Sandstone forming the floor of the valley, and are lower in their position than the Strata passed through in Journal No. 1.—Copied from the last *Statistical Account of the Parish*.

	Fh.	Ft.	In.		Fh.	Ft.	In.
Surface earth, and boulders, Sandstone forming the floor of the valley	1	0	7	Brought forward,	12	1	10
Very hard band	-	0	1 0	Shale	-	-	0 1 1
Fire-clay	-	0	1 8½	Blue fakes	-	-	0 0 7½
Fakes,	-	0	1 8	Shale	-	-	0 1 1½
White Kingle	-	0	4 6	Ironstone	-	-	0 0 3
Hard shale	-	1	5 8	Shale	-	-	0 0 4
Fakes	-	0	1 6	Coal	-	-	0 1 2
Fire-clay	-	0	0 11	Fire-clay	-	-	0 2 1½
Impure limestone	-	0	3 9	Fakes	-	-	1 0 10½
Shale with strips of fakes	1	4	6	Impure limestone	-	-	0 1 10½
Blue shale	-	0	1 7½	Shale	-	-	1 5 5
Ironstone	-	0	0 3	White kingle	-	-	0 2 10½
Shale	-	0	0 7	Hard shale	-	-	0 3 6½
Coal	-	0	0 11	White sandstone	-	-	0 2 5½
Fire-clay	-	0	1 2½	Kingle	-	-	1 0 7
Fakes	-	1	0 6½	Gray fakes,	-	-	0 1 7
Kingle	-	0	0 7	White sandstone	-	-	2 1 9½
Limestone	-	0	0 8½	Fire-clay	-	-	0 2 9
Shale with strips of fakes	1	4	10½	Light coloured fakes	-	-	0 2 0½
				Kingle	-	-	0 1 5
Carry forward,	12	1	10	Fakes	-	-	0 1 0
				Carry forward,	23	0	9

BURNHOUSE BORE—*Continued.*

	Fh.	Ft.	In.		Fh.	Ft.	In.
Brought forward,	23	0	9	Brought forward,	29	3	3½
White kingle - - -	-	2	4 6	Limestone - - -	-	0	1 8
White marble-band -	-	1	1 11½	White limestone -	-	0	1 5
Fire-clay - - -	-	0	1 5½	Light coloured fakes	-	0	1 5
Blue shale - - -	-	0	0 2½	Kingle - - -	-	0	0 4
Light coloured shale	-	0	1 3½	Light coloured fakes	-	0	0 9½
Shale with strips of fakes	-	0	1 6	Kingle - - -	-	1	4 9½
Kingle - - -	-	0	1 7½	Light coloured fakes	-	0	1 6
Fakes - - -	-	0	0 6	Light coloured kingle	-	0	3 9½
Shale - - -	-	-	1 0 0				
Shale with strips of fakes	0	0	9	Depth of the bore under			
Rock, composition not known	0	0	9	the level of the Campsie			
				valley - - -	-	33	1 0
Carry forward,	29	3	3½				

In this bore the character of many of the beds seems not to have been very clearly determined. "*Kingle*" and "*Fakes*," are terms used by the Miners, the former, to denote any hard silicious band, and the latter, Shale, with strips of impure Sandstone.

No. 3.—Journal of strata passed through in sinking a pit for Coal on the Grounds of Balquarhage on the south-eastern flank of the South hill, and to the south of the great greenstone dyke which crosses the strata of this district.

	Fh.	Ft.	In.		Fh.	Ft.	In.
Surface earth, with clay				Brought forward,	10	2	0
and boulders, - - -	-	1	1 0	Bituminous shale -	-	0	2 0
Blue shale, with marine				Dark-blue shale -	-	0	2 0
shells - - -	-	7	0 0	Shale, with large ironstone			
Shale, with <i>Trilobites</i> , <i>Cri-</i>				nodules - - -	-	0	2 0
<i>noidea</i> and shells -	-	0	3 0	Dark, soft shale -	-	0	1 0
Calcareous shale -	-	0	1 0	Black shale, with small			
Calmy limestone, with ma-				ironstone nodules -	-	0	1 6
rine shells, in two beds				Bituminous shale -	-	0	0 6
parted by a band of				Coal (not of good quality)	0	5	0
argillaceous clay -	-	1	2 0	Dark sandy shale full of			
Calcareous shale -	-	0	1 0	coal-measure plants	-	0	3 0
Carry forward,	10	2	0	Depth of pit - - -	-	13	1 0

The strata passed through in sinking this pit belong to a higher stage of the Lanarkshire coal-field than any of those beds in the Campsie district, and are probably in the same relative position as the Garnkirk, Bedlay, and Moodiesburn beds, which occur to the south of the river Kelvin.

LIST OF CARBONIFEROUS FOSSILS

FROM THE CAMPSIE DISTRICT, AND THE LOCALITIES IN WHICH THEY HAVE BEEN FOUND.

The following list of Fossils from the Campsie district shows the various beds in which each species has been discovered. Those species that have been found only in one locality may yet be found to have a wider range in the strata of the district. They have been named from the works of Davidson, D'Koninck, Phillips, Fleming, Sowerby, M'Coy, and one or two other acknowledged authorities on Carboniferous Fossils. Several species put down in the list have been marked doubtful, while one or two others have been left undetermined. They seem, however, to be nearly related to the genera or species indicated, and may in many instances be only varieties of those species whose names are given. The asterisk (*) indicates the discovery of the species in the strata above.

GENERA.	SPECIES.	AUTHORITIES.	Craigenglen Strata. Ironstone, impure Lime- stone, Sandstone, Bi- tuminous Shale, Clay- Shale, &c.	Balgassburn Strata, and Sandstone of the Campsie Valley. Clay- Ironstone, Shale, and Sandstone.	Millburn Strata. Calcareous Shale, impure Limestone, Clay-Iron- stones, and Shales.	Balgrochan-burn Strata, North-hill. Shelly Lime- stone, Clay-Ironstone, and Shale.	Main Limestone Strata. Coal, Limestone, Cal- careous Shale, Clay- Ironstone, Alum- Shale, &c.	Hosie Limestone Strata, South Hill. Clay-Iron- stones, Limestones, and dark-blue Shales.	Corrie-burn Strata, North Hill. Coralline and Fossiliferous Limestones, Clay-Ironstones, Shale, and Sandstone.	Balquharage Strata, East end of South Hill. Calmy - Limestone, Coal, Clay-Ironstone, and Shales.
<i>Plantæ.</i>										
Calamites	cannæformis	Schlotheim	..	*	...	...	...	...	...	...
Cauda-galli	(impression of fucoid ?)		...	...	...	...	...	...	...	...
Lepidodendron	elegans	Brongniart	...	*	...	...	...	...	*	...
"	gracile	Lindley	...	...	...	...	*	...	...	...
"	Sternbergii	Brongniart	...	...	...	...	...	...	...	...
"	selaginoides	Sternberg	...	...	...	...	...	...	...	...
Lepidostrobus	ornatus	Brongniart	...	...	...	...	...	...	...	...
Sigillaria	organum	Lindley	...	*	...	...	...	*	...	...
Stigmaria	ficoidea	Brongniart	...	*	*	...	*	...	*	...
Sphenopteris	bifida	Lindley	...	...	...	...	*	...	...	...
Sphenophyllum	erosum	Lindley	...	...	...	...	*	...	...	...
Trigonocarpum	(species not determined)		...	...	...	...	...	*	...	...
<i>Zoophyta.</i>										
Chætetes	tumidus	Milne Edward	...	...	...	*	*	...	*	...
Cyathophyllum	Wrighti	Milne Edward	*	...	...	...	...	...	*	...
"	(two species not determined)		...	...	...	...	...	...	*	...

GENERA.	SPECIES.	AUTHORITIES.	Craigieglen Strata.	Balglass-burn Strata.	Mill-burn Strata.	Balgrochan-burn Strata.	Main Limestone Strata.	Hosie Limestone Strata.	Corrie-burn Strata.	Balquhagge Strata.
Cyathopsis	fungites	Fleming	*	...	...	...	...	...	*	...
Lithodendron	fasciculatum	Fleming	*	...	...	...	...	...	*	...
"	junceum	Fleming	*	...	...	...	...	...	*	...
Lithostrotion	Portlocki	Milne Edward	...	...	...	...	...	...	*	...
<i>Echinodermata.</i>										
Actinocrinus	(several species undetermined)									
Archæocidaris	Urii	Fleming	*	*	...	...	*	*	*	...
Hydreionocrinus	globularis	D'Koninck	*	...	...	...	*	...	*	...
Poteriocrinus	rugosus (?)									
"	(several species undetermined)									
<i>Articulata.</i>										
Serpula (?)	hexicarinata	M'Coy	*	...	...	...	...	...	...	...
"	parallela	M'Coy	...	...	*	...	...	...	...	...
Serpulites	carbonarius	M'Coy	...	*	...	...	*	...	*	...
"	membranaceus	M'Coy	*	...	...	...	...	...	...	...
"	(species not determined)		...	...	...	...	*	...	...	...
Spirorbis	caperatus	M'Coy	*	...	...	...	...	...	*	...
"	globosus	M'Coy	*	...	...	...	...	...	...	...
<i>Crustacea.</i>										
Bairdia	curta	M'Coy	*	...	...	...	...	...	...	...
Cyclus	radialis (?)	D'Koninck	...	...	...	...	...	...	*	...
Cythere										
Dithyrocaris	Colei	Portlock	*	...	...	...	...	*	...	...
Eurypterus	Scouleri	Hibbert	...	...	...	...	*	...	...	...
Griffithides	mesotuberculatus	M'Coy	...	...	...	...	*	...	...	*
"	Eichwaldi	Fischer	...	*	...	...	...	...	*	...
<i>Bryozoa.</i>										
Ceriopora	interporosa	Phillips	*	...	*	*	...	...	...	...
"	similis	Phillips	...	...	...	...	*	...	*	...
Diastopora	megastoma	M'Coy	*	...	...	...	...	...	...	...

Fenestella	plebeia	M'Coy	*	...	+	...	*	...	*	...
"	nodulosa	Phillips	...	...	...	...	...	...	*	...
"	Morrisii	M'Coy	...	...	...	...	...	...	*	...
"	frutex	M'Coy	...	...	...	...	...	...	*	...
"	formosa	M'Coy	...	...	...	...	...	...	*	...
"	multiporata	M'Coy	...	...	*	...	...	...	*	...
"	flabellata	Phillips	...	...	...	...	...	...	*	...
"										
Glauconome	bipinnata	Phillips	...	...	*	...	...	...	*	...
"	gracilis	M'Coy	...	...	...	...	...	...	*	...
Polypora	dendroides	M'Coy	...	...	*	...	...	...	*	...
Ptylopora	pluma	M'Coy	...	...	...	...	...	...	*	...
Sulcoretepora	parallela	Phillips	...	...	*	...	*	...	*	...
<i>Brachiopoda.</i>										
Terebratula	hastata	Sowerby	*	...	...	...	*	...	*	...
"	sacculus	Martin	...	...	...	...	...	...	*	...
Spirifera	duplicicosta	Phillips	...	...	...	*	*	...	*	...
"	bisulcata	Sowerby	*	*	...	...	*	...	*	*
"	trigonalis	Sowerby	...	...	...	...	...	...	...	*
"	ovalis	Phillips	...	...	...	...	...	...	*	...
"	glabra	Martin	...	*	...	...	*	...	*	...
"	Urii	Fleming	*	...	...	...	*	*	*	...
"	lineata	Martin	...	...	...	...	*	...	*	*
Spiriferina	octoplicata	Sowerby	...	...	...	...	...	...	*	...
Athyris	ambigua	Sowerby	*	...	*	*	*	...	*	...
"	plano-sulcata	Phillips	*	...	...	...	...	...	...	...
"	Royssii	L'Eveille	...	...	...	...	*	...	*	...
Rhynchonella	pugnus	Martin	...	...	*	...	*	...	*	...
"	pleurodon	Phillips	*	...	*	*	...	...	*	...
Strophomena	analoga	Sowerby	...	...	...	...	*	...	...	...
Orthis	resupinata	Martin	*	*	...	...	...	...	*	...
"	Michelini	L'Eveille	*	...	...	...	*	...	*	...
Streptorhynchus	crenistris	Phillips	...	*	*	...	...	...	*	...
"	(variety) radialis	Phillips	...	...	...	...	*	...	...	...
Chonetes	Hardrensis	Phillips	*	...	*	...	*	*	*	...
Productus	giganteus	Martin	...	...	*	...	...	...	...	...
"	latissimus	Sowerby	*	...	...	...	...	...	...	...
"	cora	D'Orbigny	...	...	...	...	*	...	*	...
"	semireticulatus	Martin	*	*	*	*	*	...	*	*
"	(variety) Martini	Sowerby	*	...	*	*	...	*	*	...

GENERA.	SPECIES.	AUTHORITIES.	Craigenglen Strata.	Balglass-burn Strata.	Mill-burn Strata.	Balgrochan-burn Strata.	Main Limestone Strata.	Hosie Limestone Strata.	Corrie-burn Strata.	Balquharnage Strata.
Productus	longispinus	Sowerby	*	...	*	...	*	...	*	...
"	undatus	Defrance	...	...	*	...	*	...	...	...
"	costatus	Sowerby	...	...	...	...	*	...	...	...
"	Youngianus	Davidson	...	...	...	...	...	...	...	...
"	scabriculus	Martin	*	...	...	...	...	...	*	...
"	pustulosus	Phillips	...	...	...	...	*	...	*	...
"	fimbriatus	Sowerby	...	...	...	...	*	...	...	...
"	(species not determined.)		...	...	...	...	...	...	...	...
"	punctatus	Martin	*	...	...	...	...	...	*	...
"	aculeatus	Martin	...	...	...	...	*	...	*	...
"	mesolobus	Phillips	...	...	...	...	*	...	...	...
Crania	quadrata	M'Coy	*	*	...	...	*	...	*	...
Discina	nitida	Phillips	*	...	...	...	*	...	*	...
Lingula	squamiformis	Phillips	...	...	...	*	*	...	*	...
"	mytiloides	Sowerby	*	...	...	...	*	...	*	...
<i>Lamellibranchiata.</i>										
<i>(Monomyaria.)</i>										
Aviculopecten	arenosus	Phillips	...	...	*	...	*	...	*	...
"	concentricostriatus	M'Coy	...	...	...	...	...	...	*	...
"	consimilis	M'Coy	...	...	...	...	...	...	...	...
"	fimbriatus	Phillips	*	...	...	...	...	...	...	...
"	granosus	Sowerby	...	...	...	...	...	...	*	...
"	interstitialis	Phillips	...	...	...	...	...	...	*	...
"	intercostatus	M'Coy	...	...	...	...	*	...	...	...
"	leiotis (?)	M'Coy	...	...	...	...	...	...	*	...
"	plicatus	Sowerby	...	...	...	...	...	...	*	...
Pecten	Sowerbii	M'Coy	*	...	...	...	...	...	...	...
"	stellaris	Phillips	...	...	...	...	...	...	...	...
Pinna	flabelliformis	Martin	...	...	...	...	...	...	*	...
Pteronites	persulcatus	M'Coy	...	...	*	...	...	...	*	...
"	angustatus (?)	M'Coy	*	...	...	...	...	...	*	...
"	latus	M'Coy	...	...	*	...	...	...	...	...
Posidonomya	vetusta	Sowerby	...	...	...	...	...	...	*	...

<i>Lamellibranchiata.</i> (<i>Dimyaria.</i>)												
Arca	reticulata	M'Coy	*	...	...	...	...	...	...	...	...	...
" (?)	(not determined)		...	...	...	...	...	...	*	...	...	...
Axinus	axiniformis	Phillips	*	...	...	...	...	...	...	...	...	...
"	deltoides	Phillips	*	...	...	...	...	...	...	...	...	...
Cardiomorpha	oblonga	Sowerby	*	...	...	...	...	...	...	...	...	...
"	(species not determined)		...	...	...	...	...	...	*	...	...	...
Conocardium	(species not determined)		...	...	...	...	...	...	...	*	...	...
Cypricardia	cylindrica	M'Coy	...	...	...	...	...	...	...	...	*	...
"	glaberata	Phillips	*	...	...	...	...	...	...	...	...	...
"	rhombea	Phillips	*	...	...	...	...	...	...	...	...	...
"	striato-lamellosa	D'Koninck	*	...	...	...	...	...	...	...	...	...
Edmondia	quadrata	M'Coy	...	...	...	...	...	*	*	...	...	...
"	rudis	M'Coy	*	...	...	...	...	...	...	...	*	...
"	sulcata	Phillips	...	...	...	...	...	*	...	...	...	...
"	unioniformis	Phillips	...	...	...	...	...	...	...	...	...	...
Leda	attenuata	Fleming	*	...	...	...	...	...	...	...	*	...
"	longirostris	M'Coy	...	...	...	...	...	...	...	...	...	...
Leptodomus	costellatus	M'Coy	...	...	...	...	...	*	...	...	...	...
Mactra (?)	incrassata	M'Coy	*	...	...	...	...	...	...	...	...	...
Modiola	elongata	Phillips	*	...	...	...	...	...	...	...	...	...
"	lingualis	Phillips	*	...	...	...	...	...	...	...	...	...
"	Macadami	Portlock	*	...	...	...	...	...	...	...	...	...
"	(species not determined)		*	...	...	...	...	...	...	...	...	...
Myalina (?)	(not determined)		...	...	...	...	...	...	...	...	*	...
Mytilus	Flemingii	M'Coy	...	...	...	...	...	*	...	...	...	...
Nucula	gibbosa	Fleming	*	...	...	...	...	*	*	*	...	...
"	lineata	Phillips	*	...	...	...	...	...	*	...	...	...
"	luciniiformis	Phillips	...	...	...	...	...	...	*	...	...	...
Sanguinolites (?)	plicatus		...	...	*	...	...	...	...	...	...	...
Sedgwickia (?)			*	...	...	...	...	...	...	...	...	...
Venus (?)	elliptica	Phillips	*	...	...	...	...	...	...	...	...	...
<i>Pteropoda.</i>												
Conularia	quadrissulcata	Sowerby	...	...	...	...	...	...	...	*	...	...
<i>Gasteropoda</i>												
Dentalium	inornatum	M'Coy	...	...	...	...	...	...	*	...	...	...
Euomphalus	acutus		*	...	...	...	...	...	...	...	...	...
"	carbonarius	Sowerby	...	...	...	...	...	...	*	...	...	...

GENERA.	SPECIES.	AUTHORITIES.	Craigenglen Strata.	Balglass-burn Strata.	Mill-burn Strata.	Balgrochar-burn Strata.	Main Limestone Strata.	Hosie Limestone Strata.	Corrie-burn Strata.	Balquharage Strata.
Euomphalus	pentangulatus	Sowerby	*	...	...	...	*	...	*	...
"	radians	D'Koninck	*	...	...	...	...	...	...	...
"	tuberculatus	Fleming	*	...	...	...	...	...	...	...
Loxonema	rugifera	Phillips	*	...	...	...	...	...	*	...
"	scalaroidea	Phillips	*	...	...	...	...	...	...	...
"	Lefebvrei (?)	L'Eveille	...	...	...	...	...	...	*	...
"			*	...	...	...	...	...	...	...
Macrocheilus	acutus	Sowerby	*	...	...	...	...	...	...	...
"	imbricatum	Sowerby	*	...	...	...	...	...	...	...
"	Michotianus	D'Koninck	*	...	...	...	...	...	...	...
"	ventricosus	Goldfuss	*	...	...	...	...	...	...	...
"	(two species not determined)		*	...	...	...	...	...	...	...
Murchisonia	quadricarinata	M'Coy	*	...	...	...	*	...	...	...
"	striatula	D'Koninck	*	...	...	...	...	...	...	...
Natica	elliptica	Phillips	*	...	...	...	...	...	...	...
"	elongata	Phillips	*	*	...	...	...	...	...	...
"	Omaliana	D'Koninck	*	...	...	...	...	...	...	...
"	variata	Phillips	*	...	...	...	...	...	...	...
"	canaliculata	M'Coy	*	...	...	...	...	...	...	...
Pileopsis	carinatus	M'Coy	...	*	...	...	*	...	...	...
Pleurotomaria	contraria	D'Koninck	*	...	...	...	...	...	...	...
"	expansa	Phillips	*	...	...	...	...	...	...	...
"	Galeottiana	D'Koninck	*	...	...	...	...	...	...	...
"	monilifera	Phillips	*	...	...	...	...	...	...	...
"	Sowerbyana	D'Koninck	...	...	...	...	...	...	...	...
"	(undetermined)		*	...	...	...	...	*	...	...
Trochus	lepidus	D'Koninck	*	...	...	...	...	...	...	...
<i>Nucleobranchiata.</i>										
Bellerophon	decussatus	Fleming	*	...	...	...	...	*	...	...
"	Dumontii	D'Orbigny	*	...	...	...	...	...	...	...
"	striatus	Fleming	*	...	...	...	...	*	...	...
"	Oldhamii	Portlock	*	...	...	...	...	*	...	...
"	Urii	Fleming	*	...	...	...	...	*	*	*
Porcellia	armata	Verneuil	*	...	...	...	*	...	...	...

<i>Cephalopoda.</i>										
Actinoceras	<i>giganteus</i>	Sowerby	...	...	...	...	*	...	*	*
Cyrtoceras	<i>Gesneri</i>	Martin	*	...	...	...	...	...	...	...
Goniatites	<i>diadema</i>	Goldfuss	*	...	...	...	...	...	...	...
"	<i>micronotus</i>	Phillips	...	...	...	...	*	*	*	...
"	<i>Listeri</i>	Martin	...	...	...	...	...	*	...	...
Nautilus	<i>ingens</i>	Martin	...	...	...	...	...	*	...	*
"	<i>tuberculatus</i> (?)	Sowerby	...	...	...	...	*	...	...	...
"	<i>subsulcatus</i>	Phillips	...	...	...	...	*	...	...	...
"	<i>sulcatus</i>	Sowerby	...	...	...	...	*	...	*	...
Orthoceras	<i>attenuatum</i>	Fleming	...	...	...	...	*	...	*	...
"	<i>cinctum</i>	Sowerby	*	...	...	...	...	*	*	...
"	<i>cylindraceum</i>	Fleming	...	...	...	...	...	*	...	...
"	<i>dilatatum</i>	D'Koninck	*	...	...	...	...	...	*	...
"	<i>lineale</i>	D'Koninck	...	...	...	...	...	...	*	...
"	<i>Muensterianum</i> (?)	D'Koninck	...	...	*	...	...	...	...	...
"	<i>pygmæum</i>	D'Koninck	...	...	...	...	...	*	...	...
"	<i>rugosum</i>	Fleming	*	...	...	...	...	*	...	...
"	<i>undatum</i>	Fleming	*	...	*	...	*	*	*	...
"	<i>unguis</i>	Phillips	*	...	...	...	...	...	...	...
<i>Pisces.</i>										
Cladodus	<i>mirabilis</i>	Agassiz	...	...	...	...	*	...	...	...
"	(species not determined)		...	...	...	...	...	*	...	...
Cochliodus	<i>magnus</i>	Agassiz	...	...	...	...	*	...	*	...
Cœlacanthus	<i>lepturus</i>	Agassiz	*	...	...	...	...	*	...	...
Ctenacanthus	(species undetermined)		...	...	...	...	...	*	...	...
Ctenoptychius	<i>serratus</i>	Agassiz	...	...	*	...	...	...	*	...
Holoptychius	<i>Portlocki</i>	Agassiz	*	...	...	...	...	*	...	...
"	<i>minor</i>		*	...	...	...	*	*	...	...
Homacanthus	(species undetermined)		...	...	*	...	...	...	...	...
Helodus	<i>lævissimus</i>	Agassiz	...	...	...	...	*	...	*	...
Megalichthys	<i>Hibberti</i>	Agassiz	*	...	*	*	...	*	...	...
Palæoniscus	<i>Robisoni</i>	Hibbert	*	...	...	...	*	...	...	...
Petalodus	<i>Hastingsiæ</i>	Owen	*	...	...	...	*	...	*	*
Pœcilodus	<i>obliquus</i>	Agassiz	*	...	...	...	*	...	...	...
Polyrhizodus	<i>radicans</i>	Agassiz	*	...	...	...	...	...	...	...
Psammodus	<i>porosus</i>	Agassiz	...	...	*	...	...	...	...	...
Rhizodus	<i>Hibberti</i>	Agassiz	*	...	...	...	...	*	...	...

Section across the Valley of Campsie, looking toward the West, showing the general position of the strata at the Spout of Ballagan.

a a, Yellow and Gritty Sandstone of South hill; b, Dunglass, columnar basalt; c c, Trap-dykes crossing the strata; d, Ballagan beds, thin-bedded limestone and shales; e, Fault in the strata; f, Yellow Sandstone; g, Overlying trap of North hill; h, Steep sloping bank of debris.

Section across the Valley of Campsie, looking towards the East, showing the amount of denudation in the Valley, and the general position of the strata on the North and South hills.

a, Overlying trap of North hill; b b, Yellow sandstone, forming the highest bed under the trap of North hill, and crowning the highest summit of South hill; c c c, Hosie limestone series; d d d, Main coal and limestone series; e e e, Sandstone series forming the floor of the Valley to the East; f f, Trap-dykes crossing the strata; g, Faults in the strata of North and South hills.

Section along the North hill, Campsie, showing the general succession of the strata from the Strathblane valley eastward to Corrie-burn, near Kilsyth.

Old Red Sandstone. Strathblane Valley.

Ballagan series, and overlying trap-rock.

Termination of Ballagan series.

Campsie Main Coal and Limestone series.

Corrie-burn series.

a, Overlying trap; b, Large veins of trap running from the hill toward the valley, and cutting off the strata at several points; c, Interstratified greenstone, trap-tufa, and claystone, Corrie-burn.