cholepsy. We must not forget that we find comparable phenomena in the normal and especially in the pathologic forms of sleep, in dreams, in exhaustion, and particularly in the emotional states. The psychologic and physiologic study of these facts is far from finished, and it appears to me indispensable for the explanation of the phenomena presented by our patients. Hence if there be anything of importance in what I have brought to your attention this evening, it is the clinical description of certain curious mental troubles which I have endeavored to compare and connect, in order to show you that the mental degradation which paves the way for the obsessions and which must even play a part in the genesis of certain deliria, is not always effected gradually, but may come on abruptly and constitute a genuine psycholeptic crisis.

REFERENCES.

- ¹ Ottolenghi: *Epilepsia psychica*. Riv. sper. di freniatria, 1892.
- ² Lad. Haskovec: *Contrib. à la connaissance des idées obsédantes*, xiiième Congrès de Médecine à Paris, 1900; section de Psychiatrie, p. 121.
- ³ Pick: *Brain*, 1903, p. 242.
- ⁴ Les Obsessions et la Psychasténie, 1903, vol. i, p. 504.
- ⁵ A more complete study of these two and other similar cases may be found in my work, "Obsessions et la Psychasténie," vol. i, p. 506, and vol. ii, p. 54. I have also recently presented my observations on an analogous trouble in an epileptic before the Société de Psychologie: *Bulletin de l'Institut Psychologique*, 1904, p. 215.
- ⁶ Ball: *Revue Scientifique*, 1882, ii, p. 42.
- ⁷ Névroses et idées fixes, 1898, ii, p. 61.
- ⁸ Ségas: *Leçons sur les maladies mentales*, 1895, p. 70.
- ⁹ Obsessions et psychasténie: Les périodes psychasténiques, i, pp. 439, 441.
- ¹⁰ *Op. cit.*, ii, p. 424.
- ¹¹ Névroses et idées fixes, ii, pp. 63, 71. Obsessions et psychasténie, i, p. 319.
- ¹² *Op. cit.*, i, p. 252.

GASTRO-ENTEROSTOMY.

(A PRELIMINARY NOTE.)

BY JAMES G. MUMFORD, M.D., BOSTON,
Assistant Visiting Surgeon to the Massachusetts General Hospital.

OPERATIONS on the stomach were undertaken fearfully and carried out timidly for many years. We have not yet outgrown that mental attitude. Apparently the feeling has been, if you open the digestive tract you are doing a dangerous thing; do as little damage as possible and get the visceral wounds closed as soon as you can.

Experience with operations for malignant disease of the stomach first opened our eyes to the fallacy of this over-cautious view. There are two operations usual in the case of gastric cancer, — partial gastrectomy and gastro-enterostomy. At first sight one would suppose gastro-enterostomy to be far the more formidable, but operators have learned that in the long run, gastro-enterostomy for malignant disease results in a higher mortality than does gastrectomy.

Setting aside the consideration of, and in the case of "benign" disease, looking for a radical operation for stomach drainage, for an operation which shall return the parts to an anatomical condition essentially normal, one sees that some form of gastro-enterostomy must be done. Finney's operation is valuable, for it returns the parts to a condition approximating the normal, but Finney's operation is not always applicable. Rodman's partial gastrectomy, combined with

all cases of gastro-enterostomy, returns the parts essentially to the normal, but Rodman's operation seems needlessly radical for routine practice.

None of the other operations, as ordinarily performed, regularly restores the parts to anything like a normal condition.

The object of this communication is to urge an operation I have practiced of late, an operation elaborated out of the well-known procedure of Chaput. As originally done by Chaput, that operation consisted of:

(1) Posterior gastro-enterostomy with the long loop; (2) Entero-enterostomy; (3) Section of the afferent loop between the two anastomoses.

So far as it goes that operation is very satisfactory, but it does not restore the parts to their normal relations. Vicious circle sometimes ensues, the pylorus may resume its functions; the new gastric stoma may close.

I advocate a fourth step, — invariable, routine, (4) Section of the pylorus. That adds nothing to the risks; vicious circle is rendered impossible; the duodenum, now side-tracked, becomes a mere duct, a continuation of the common bile duct.

Doubtless this method of operating is not original, and it would be preposterous to describe it as a novelty; but since it offers a certain solution of the vicious circle problem, since it leaves the parts much as nature intended and since it is easy of accomplishment, I am urging it as a routine measure.

THE PROPHYLAXIS OF SYPHILIS.*

BY WM. G. MACDONALD, M.D., BOSTON,
Dermatologist to the Carney Hospital.

AT the last Charitable Conference held at Boston, one of the speakers said something to this effect: "We will never accomplish much along some lines of charitable work, until we recognize the fact that certain diseases exist, and are willing to discuss them and their consequences as openly and as freely as we now argue other and kindred matters." These were not his exact words, but they express the lesson he intended to teach.

Public health and public charity cannot be disassociated. They walk hand-in-hand, and deal always with the same individual. Almost the whole of charitable work comes down to the preservation of health, the care of the sick, or the burial of the dead. Our hospitals are managed by health experts, but are founded, endowed and supported by charity. Yet, on the one question of venereal disease, charity closes its delicate ears with its daintily gloved finger tips, looks inexpressibly shocked and a little bit bored, and refers the whole matter to public health. The excuse is that these diseases are self-inflicted and can only come about by the depravity of the individual. Even if this be true, what matters it? When the Samaritan saw a stranger lying by the roadside, naked, bleeding and senseless, he did not ask who started the fight. Not at all; his every effort was devoted to the care of the sick man. But, in point of fact, the charge is not

*Read before the Catholic Charity Conference, Oct. 29, 1904.

FIG. 1. — Operation of Chaput: (1) Gastro-enterostomy with the long loop. (2) Entero-enterostomy. (3) Section of the afferent loop between the two anastomoses.

FIG. 2. — Operation recommended: (1) Posterior gastro-enterostomy with long loop. (2) Entero-enterostomy (3) Section of afferent loop between the two anastomoses. (4) Section of the pylorus.

true. The cases of syphilis innocently contracted are almost innumerable.

Many epidemics of syphilis have been reported, in which the first offending cause was most innocent. Dr. Commenge has tabulated a large number of such cases, of which a few may be cited.¹ For instance, Riccordi reported in 1863 the case of an abandoned infant taken to nurse; this child communicated syphilis to twenty-three individuals. At Uboldo, in the same year, another abandoned child transmitted the disease to nineteen persons, including its nurse. At Capistrello, syphilis was propagated by a child at the breast, and through lack of knowledge on the part of the physicians and victims, lasted for eight years. At last, in 1867, it was recognized and investigated, and found to have already afflicted more than three hundred persons. In our own country, Dr. Buckley published in 1894 a treatise called "Syphilis in the Innocent," in which he cites a large number of cases, and places especial stress on those which resulted from vaccinal accidents. In these, however, the remedy was easy, and the practical disuse of humanized virus has eliminated them. In France, Fournier, probably the best syphilographer in the world, concluded that at least one in ten cases of syphilis was extragenital,² and this did not include the large number of married people who might contract the disease genitally but innocently. Fournier collected a number of cases caused by unclean instruments in the hands of physicians.³ He cited one instance in which a specialist of Paris had contaminated as many as seventy persons. Tattooing has had many authenticated victims. Razors, lead pencils, pipes and cigars, many others. The practice of passing around water in the theaters is an abominable one. I have seen an usher, whose mouth was full of mucous patches, passing about these glasses. He probably sipped when he was thirsty. Last summer I treated a young girl who had a chancre in the mouth, innocently contracted. She was a bundle girl in one of the large stores, and presented herself with a chancre on the tonsil, probably caused by virus received into the mouth from the edge of a drinking vessel, and caught by the protruding gland on its way down the throat. She had well-defined secondary symptoms when I saw her. I warned her very carefully about kissing, about using the same cup as others, etc., but did not name the disease to her mother or herself. Some weeks after, she reappeared with an infected friend—a chancre on the tongue, and no history. After some groping, my first friend asked innocently if lending her chewing gum would be any harm, and said that I had neglected to warn her about that!

In the face of such facts, can anyone doubt that there should be some education in syphilis? that there should be, even among ordinary people, a clear understanding of the nature and dangers of this disease? Why should not philan-

thropists discuss it openly in their conferences, even though the audiences were of both sexes? A disease so insidious and so treacherous, and so capable of entering into and contaminating the purest and best-behaved of families, sparing none from the grandsire to the swaddled infant, should be at least understood, so that its first advances might be recognized and checked. Once understood, and discussed without fear or shame, effective measures for its suppression will be taken.

Some few months ago, Professor Lesser of Berlin reported the second conference, held March 6, 1904, of the German Society for Combating Venereal Disease.⁴ This society was founded in October, 1902, and already numbers some four thousand members. At this conference, it was particularly noticed that, notwithstanding the great strides made by the medical profession in the treatment and cure of venereal diseases, yet on account of the reluctance of the people at large to speak of them, but little had been done for their prophylaxis. It was then decided that, in the German Society Educational Pamphlet, there should be set forth, in a simple and comprehensive form, the nature of venereal diseases, their dangers, and the methods of guarding against them. In this connection it was voted that there should be set down the methods of protection which might be employed in extra-conjugal intercourse. This latter recommendation would, of course, be at utter variance with our line of thought.

The sanitation of prostitution was also considered, and Professor Lesser urged strongly the founding of free institutions for the treatment of venereal diseases. In conclusion, he very wisely counselled a slow and careful advance. Old prejudices and beliefs were not to be pushed away, but rather so worn down by gentle, constant friction that the light behind might be continually more perceptible.

In France, the inscription of prostitution is an old story. With the inscription, comes the regular physical examination, and removal to a hospital, if venereal disease of any kind be found. This works fairly well, but does not, by any means, cover the whole ground. In the first place the number of uninscribed prostitutes is very large, and secondly, a perpetual warfare is being waged against these venereal hospitals because they are prisons. Some of our most eminent authorities on syphilis have constantly objected to them on purely ethical grounds.

In the British Islands the so-called Lock Hospitals may be found, scattered here and there, in garrison towns. As far as I have seen, they are of little service in the prevention of syphilis, since few if any of the inmates are in the acute, contagious stages of the disease. Those that I have noticed were soiled and dishevelled old remnants that were long ago taken from the bargain counter. There is no inscription in British cities, at present, and, therefore, no official way of discovering the diseased prostitute. If

¹ La Prostitution Clandestine à Paris, 1897.

² Traité de la syphilis, Fournier, 1898.

³ Bulletin Médical, 15 et 19 mai, 1895.

⁴ Journal of Cutaneous Diseases, June, 1904.

there were, I question whether the English public would allow compulsory detention on account of bodily illness.

In a conversation the other day with Mr. John B. Martin, Penal and Charitable Commissioner of Boston, a man of much experience in these matters, and one of broad and well-balanced views, he suggested a plan which would well cover a part of the field, and which is not as yet in use so far as I know. His suggestion was as follows: that, whenever, hereafter, a man or woman be convicted of an offence which may call for either fine or imprisonment, he or she should be physically examined, and if venereal disease be found, then the punishment must be by imprisonment. This would be farther-reaching than would seem at first glance, since a large number of those arrested for petty offences are not only the very ones who are liable to be suffering from syphilis, but, also, the ones most careless about its transmission.

It should also be the routine duty of every physician of an institution, whether it be jail, almshouse, or prison, to thoroughly and carefully examine each new inmate so that proper isolation might be secured if necessary. This examination should take in every part of the body, and the trustees or superintendents of such institutions should provide suitable rooms, instruments and attendants for the purpose.

The standing armies of Europe have proven to be about the only communities from which comparative statistical estimates could be made, and Dr. Commenge has considered this subject rather carefully.⁵ He finds that, for instance, in the five years from 1888 to 1892 inclusive, there was a yearly average in the English army of about two hundred in the thousand taken into hospital for venereal diseases; while in the French army the proportion was but forty-four to the thousand, and that there were about thirty-seven English syphilitics to the thousand, averaged yearly, with only eight or nine French. Researches into the statistics of the German, Russian, Austro-Hungarian and Roumanian armies show that the proportion of disablements under the British flag was greater than that of any other European nation.

From these premises, Dr. Commenge draws the conclusion that the regulation of prostitution is the great safeguard against syphilis, but he forgets many important accessories. The English army is drawn from a hard-headed Saxon or Celtic yeomanry, not much given to immoral practices. The young men are kept in as much ignorance of venereal disease as the young women. Suddenly they enlist, are sent to some of the English colonies, and are at once launched into semi-savage communities either without perceptible morals, or worse still, debauched by the traders of all nations who have preceded the army. Is it any wonder that these young men contract syphilis? Is it any wonder that, having contracted it, a sense of shame compels their silence until, unwittingly, they have transmitted the disease to

one or more of their comrades? In other words, the English youth suffer on account of a lack of proper education in matters which are only too familiar to their continental brethren.

At the time of Dr. Commenge's article, the American army was not to be considered. We had a handful of sturdy fellows, living in a few scattered garrisons, and subject to all the inclemencies of our northwest winters. Immoralities had little chance among such folk. Now, however, we are facing a different problem. With the acquisition of new territory comes the demand for an enlarged army, and instead of the quiet, moral influences of the northwest, we must send our youth to tropical countries, to come in contact with indolent, semi-civilized, unclothed peoples. We should not send them uneducated.

Now, what should we do for those at home? We should teach them, in as kindly and proper way as possible, the dangers, near and remote, of improper sexual intercourse, and not leave this important part of their education to bitter experience and the lying pamphlet of the traveling quack. Suppose, for instance, the emergency association included some consideration of syphilis in its regular lectures given to policemen, firemen and others. If to this were added an enumeration of the many innocent ways in which syphilis might be contracted, together with an insistence that false shame should be thrown aside, and that the sufferer should apply at once for medical aid, much good might be done. It is of the utmost importance that lectures on this subject should be given to our militia by their regular medical officers, and that a thorough understanding of their teachings should be demanded of the men. And this, especially, before any regiment could be considered eligible for foreign service.

As to the question of proper isolation, each community must look out for itself. In Boston, a few years ago, the city government appropriated a small sum for the building of a hospital for chronic, contagious diseases. Nothing was done, however, toward the actual, practical pushing of the work, and our present mayor dropped the whole scheme because he very wisely considered that the sum appropriated was too insignificant to be of any use, and he did not care to plunge into unknown financial depths for the sake of a new enterprise, when so many of the existent departments and institutions needed much more money than he was able to spare.

Long Island, to be sure, has a well-furnished hospital, but Long Island's institutions are primarily and always almshouses, and its inmates must be stigmatized as paupers. It is impossible that an institution working under laws which are intended to discourage applications for admission should be of much service in the controlling of contagious disease.

No one can enter the Long Island Hospital unless he or she has a pauper settlement in Boston, either acquired from parents, or through a residence of five years consecutively, during which time at least three years' payment of all state,

⁵ *La Prostitution Clandestine à Paris*, 1897.

county, city, or town taxes must be made.⁶ This, naturally, bars out the great majority of active syphilitics who are essentially birds of passage, and destroys the whole object of a contagious hospital, namely, the protection of the community. As a matter of fact, like the lock hospitals of England, the Long Island hospital treats, for the most part, those who are beyond the active and contagious stages of the disease.

It would seem possible, however, to reconstruct our present system so as to separate absolutely the hospital from the almshouse. They could exist as two separate and distinct institutions, under absolutely different names and managements, even though they occupy the same island.

In point of fact, according to the law of 1902,⁷ no person under treatment for syphilis, or other chronic contagious disease, and, consequently, a source of expense to town or commonwealth, can be deemed a pauper. Under this law, then, the pauper settlement act would be of no consequence to the individual. Circulars might be addressed to physicians requesting them to send such of their patients as might be a menace to the public health, to this hospital. Its geographical situation might do much toward answering objections as to bodily detention, as, on an island, there might be much bodily freedom with little danger of absolute departure.

The general public would need to be educated to the fact that we here possessed a desirable place for treatment, but as in the case of the South Department of the City Hospital, this knowledge would come by a little gentle and judicious management, and a real advance would be made in the prophylaxis of syphilis in this community. Other communities would soon profit by our example.

THE ETIOLOGY OF PULMONARY TUBERCULOSIS CONSIDERED IN RELATION TO ITS THERAPEUTICS.*

BY LOUIS F. HIGH, M.D.,

Medical Director, Pineshire Sanitarium, Southern Pines, N. C.

OUR acceptance of the theories with regard to the causative relation of certain germs in the production of particular diseases has led us too far afield in some instances. The exciting cause has been exalted almost to the point of sole cause in cases in which the predisposing factors are most important. This is particularly true of tuberculosis.

While most disease germs require none or but indifferent modification in the physiological processes of the body for their growth, the tubercle bacillus is a notable exception to that rule. There are few known pathogenic micro-organisms that require such definite conditions for the maintenance of their vitality and proliferative capacity. Though the limitations of favorable environment are narrow, clinically, it appears that a slight

physiological deviation of a particular kind is sufficient to meet the requirements of the parasite. Proof of this is seen in the large per cent of recoveries found post-mortem in which the disease had not been suspected during life. The favorable nature of their environment having changed the bacilli had perished leaving a cicatrix as evidence of the feebleness of their struggle against an approximately normal resistance.

It is asserted by various observers that from 50 to 98% of all persons have tuberculosis at some period of life. As the mortality rate of the disease is about 15% of mankind's total death-rate it is at once seen that by far the larger part of those in whom it develops recover without treatment or without any directed to the real cause of the disorder. Again, it appears that the tubercle germ has a very slight hold upon life, and can only reproduce itself under very certain and particular conditions. It has been experimentally demonstrated that the greater number of bacilli contained in phthisical sputum and in the cavities are incapable of further growth.¹

The difficulty of growing vigorous cultures from bacilli contained in pieces of old tissue infested by the organism indicates that many of those so situated have lost their vitality. Even when cultivated under favorable conditions and at a proper temperature upon dead media and shielded from all deleterious influences, they die within a relatively short time.

Furthermore, it has been shown that the bacilli of greatest virulence are more difficult to cultivate and that their growth is scantier than in the case of bacilli of less virulence.²

Finally, sufficient has come to light from the investigation of the bio-chemistry of the bacillus and analysis of the therapeutic resources that have shown decided influence over its control, to prove that the conditions for its multiplication are limited and transitory.

That so large a per cent of recoveries follows the simple procedure of keeping patients constantly surrounded by an air supply of absolute purity is evidence which, in the present state of our knowledge, admits little doubt that the consequences of impure air as a result of deficient ventilation is the underlying cause of pulmonary tuberculosis.

It has been demonstrated again and again that the remedy which dispels the sallow skin, pale face, cold feet and hands, weak pulse, feeble power of digestion and assimilation with the attendant loss of nerve-energy, all of which are prime characteristics of deficient oxygenation, is the same that arrests tuberculosis when applied sufficiently early and continued under proper conditions sufficiently long.

Herein have therapeutics cast a ray which has illumined and shown the primary, the pregermic cause, the physio-chemical condition without which tuberculosis practically does not exist. The catabolism of suboxygenation is resolved from dyscrasia into disease by such even and con-

* Revised Laws, Chap. 80.

⁷ Acts of 1902, Chap. 213, Section 2.

⁶ Read before the Medical Society of Virginia at Richmond, Oct. 18-21, 1904.

¹ Nothnagel's Encyclopedia, p. 49.

² Journal of Medical Research, July, 1901.