witness, and the clerk shall thereupon issue a subpoena as hereinbefore provided. If the parties do not agree that a medical expert witness shall be summoned in the case, the court or judge, upon motion of either party and upon hearing, may determine the question, and may designate the person to be summoned, if any, as hereinbefore provided.

Section 3. Such witness shall be paid for his attendance, travel, and services a reasonable compensation, to be allowed by the court, and paid out of the treasury of the county. In all civil actions and proceedings the defeated party shall be liable to refund the amount so disbursed, and after final judgment an execution may issue against him therefor in favor of the county commissioners, or, in the county of Suffolk, the city of Boston.

Section 4. In any case the court, upon its own motion or for cause shown, may order more than one, and not exceeding three persons, to be summoned as medical expert witnesses, and such additional witnesses shall be designated and summoned, and shall perform the same services and receive the same compensation as hereinbefore provided.

[Section 5. In any criminal proceeding the defendant may call and examine other medical expert witnesses in addition to those hereinbefore provided for, but at his own cost, and in such case other medical expert witnesses may be called and examined in behalf of the Commonwealth.]

Section 6. No medical expert witness shall be admitted to testify before any court or magistrate except as hereinbefore provided.

OTIS'S TREATMENT OF STRICTURE OF THE URETHRA.¹

BY GEORGE W. GAY, M. D.
Surgeon to Boston City Hospital.

OTIS's method of treating stricture is a combination of division and urethrotomy. It is designed for all classes of the affection located anterior to the membranous portion of the urethra, and aims at a radical cure.

The operation is performed by putting the stricture on the stretch, slightly cutting it with a concealed blade, and then at once dilating it to a little more than the full size of the normal canal, this having been previously determined by the use of the urethra-meter. The calibre of the canal is to be maintained by the passage of steel sounds till the rent or incision is healed, when, if the operation has been properly performed, and the after treatment faithfully carried out, the stricture, in nine cases out of ten, is permanently cured.

The operation may usually be done without an anæsthetic. The patient is confined to the house but a few days, and in the lighter cases not at all. Dr. Otis claims that accidents are comparatively infrequent, and the danger slight. He has operated in over seven hundred cases "without a death or any permanent disability." But it should be noted that in several of his cases the constriction was so slight that the necessity for an operation may well be questioned. It will be very difficult, for example, to convince practical surgeons that a urethra which admits a No. 36² steel sound requires cutting so that a No. 38 will pass; or in

more general terms that a urethra which takes a No. 30 or over requires any operation to allow of the passage of a larger instrument.

Although our experience in this method of treating stricture is limited, yet it is quite sufficient to justify the opinion that in *deep* strictures it is by no means free from danger, and should be resorted to only after other and safer modes of treatment have failed.

CASE I. A robust man, twenty-three years of age, came under my care in October, 1879, with an obstinate stricture of the urethra of four years' duration. It followed gonorrhœa, and was located in the bulbous portion. The symptoms were: frequent and painful micturition, constant urethral discharge, a heavy dragging sensation over the bladder, and urine loaded with pus and triple phosphates.

The urethra was extremely sensitive and the stricture unyielding, so that after six months' constant efforts, twice a week, at gradual dilatations, I could only pass a No. 22 sound. As the symptoms were only partially relieved and his nervous system was breaking down, a more radical treatment was indicated.

In June, 1880, I cut and dilated the stricture till a No. 34 steel sound readily passed through it, after the meatus had been divided.

The second day after the operation this patient had chills followed by partial suppression of urine, fever, emaciation, and loss of strength. During his convalescence he had an attack of epididymitis, and subacute inflammation of the bladder, so that it was six or seven weeks before he regained his usual health. He now, at the end of a year, passes No. 24 sound once a fortnight; micturition is free and easy, and his only trouble is a little of the old dragging sensation in the region of the bladder.

CASE II. Mr. B. had gonorrhœa twenty-five years ago, but had no trouble in passing water till within five years. He was admitted to the hospital in September, 1880, for complete retention, which was relieved by aspiration, catheter, and hot bath. A week later the patient was etherized, the urethra dilated sufficiently with Gouley's sound to allow the use of Otis's urethrotome. A stricture two inches from the meatus, and another in the bulbous portion, were then cut and dilated till a No. 25 readily passed through them.

The patient soon began to have chills and high temperature, followed by loss of strength, and emaciation. The sound could be used but a few times, as an abscess formed in the perinæum, which required opening; fortunately it did not communicate with the urethra, and soon healed.

He was laid up three months, but now, eight months after the operation, uses a No. 24 sound, has no trouble in passing water, and considers the operation a great success.

CASE III. Mr. C. entered the hospital in October, 1880, with a stricture in the bulbo-membranous urethra which admitted only a small bougie. It followed a gonorrhœa about seven years ago. Dr. Thorndike performed Otis's operation on the 18th of October, cutting and dilating the stricture up to No. 26. Operation followed by chills, some hæmorrhage, retention, and high fever. Pyæmia set in and eight large abscesses were opened within three weeks. They were located on the arms, legs, and back; none were in the vicinity of the stricture. The patient was delirious, lost flesh and strength, and came very near losing his life. There were no other cases of pyæmia in the hospital at the

¹ Read before the Boston Society for Medical Observation.

² The numbers refer to the French scale.

time; on the contrary the sanitary condition of all the wards was good. Three months after the operation he was occasionally using a No. 25 sound, micturition was normal, and his general condition was satisfactory.

In October I operated on another patient, who had two strictures admitting, respectively, a No. 9 and 12 sound. No. 28 passed easily after dilatation. He had a chill within twenty-four hours, but no further trouble while under observation. The patient having been discharged from the hospital for misconduct at the end of ten days the case is incomplete.

The above cases include all that have been treated by this method in the City Hospital thus far, and the experience is not such as to encourage further trial. The strictures were fair samples of those most commonly met with in practice, but the accidents were much more serious than those which usually follow internal urethrotomy or divulsion as formerly practiced.

The grave complications in the above cases were undoubtedly due to the great violence done to the deep strictures of the perinæum. The walls of the urethra were widely torn, the nerves and blood-vessels were stretched and lacerated, and the cellular interspaces were opened up, thereby furnishing depots for the collection of urine, blood, and pus. These fluids rapidly decompose, infiltrate the tissues, and furnish all the conditions necessary for septic poisoning.

Although our patients are at present in a very satisfactory condition as regards their strictures, it is as yet too soon to determine whether the improvement will be permanent. But even if a radical cure should be the result in all of them, we cannot but feel that the danger attending this operation, when performed as directed by its distinguished author, is too great to justify its use in ordinary cases of deep strictures of the urethra.

SOME OBSERVATIONS ON THE SURGICAL TREATMENT OF RECTAL AFFECTIONS.

BY WALTER ELA, M. D., OF CAMBRIDGE, MASS.

THE frequent occurrence, pain, and discomfort of rectal diseases makes any contribution tending toward curing or alleviating them more speedily, thoroughly, or certainly, not wholly unwelcome. Several cases of rectal diseases which had been unsuccessfully operated on have lately come under my care, and as the writer has had some opportunities to compare the methods and results of the surgical treatment of these affections here and elsewhere, he has deemed it worth while to note the advance that has been made in this particular branch of surgery.

Said a well-known local surgeon in an amphitheatre before a class of students, "the operation for fistula is simple, any one can do it. Mr. — can do it. Here, slit up this fistula," after he had introduced a director through the external and internal openings. The rapid placing of a pledget of lint and a T bandage made the performance seem simple enough. But this very patient soon after leaving the hospital returned with a recurrence of his malady, which had been superficially but not thoroughly operated on. So it was not soundly healed by so simple an operation as we were led to believe it would be.

Syme's¹ doctrine, that in performing the operation

for anal fistula it is sufficient to divide the parts between the external and internal openings, regardless of farther-reaching sinuses, is not followed by beneficial results, except in a few persons of a strong, healthy constitution. His theory, which has been entertained by some authors, that a fistulous sinus running up above the internal opening can be safely left untouched, is not borne out by experience. Failure of the operation thus performed is the result in the majority of cases. The inference to be drawn is obvious.

As a reward of an imperfectly-performed operation for anal fistula we have often seen a fresh sinus appear directly beneath the one operated on, and perhaps nearly cicatrized, or the wound not heal, owing to a branch or farther-reaching sinus which had not been properly laid open.

There is a distinction to be made whether a discharge comes from a healing, pus-secreting wound, or whether, coming from another source, it courses over a healing surface. In the latter case the discharge may catch under the healing edge of the wound, burrow, and finally form a sinus beneath the healing surface.

The old and unjustifiable process of excision of the tube-like fistulous track, as practiced now by an eminent professor of surgery in Paris, and also by a noted surgeon in this neighborhood, will often be followed by failure if a lateral branch sinus exists, which has to be cut across in the operation; and this method cannot be commended, since it shows an ignorance of the pathology of anal fistula, and non-success is the usual result.

In a healthy subject an uncomplicated anal fistula occasionally heals spontaneously, or with the simplest treatment; so it does sometimes by various caustics, injections, ligatures, elastic or otherwise, but the result by these means is always doubtful. In a complicated case, as frequently occurs, the free use of a cutting instrument is our main-stay, and we can never make sure of the result of a case unless we use it. To guarantee a cure the main and branch sinuses must be freely laid open, and the edges of the wound liberally trimmed off.

If severe hæmorrhage is apprehended after incising tracks reaching far up the bowel, the screw tourniquet with twine, as Luke's, or, better, the elastic ligature, may be used for cutting through the cul-de-sac above the internal opening. The solid elastic ligature, with a metallic ring, is quicker in action, easily applied, and absolutely painless. The most successful prefer usually to cut, and depend on efficient plugging to arrest the hæmorrhage.

Beyond a thorough and solid packing of the wound with absorbent cotton (not lint as is used here), which serves greatly to consolidate the lax tissues, the wound requires but little active after-treatment, but jealous care and watching must be observed to detect early any small or perhaps unnoticed branch sinus.

The after-treatment is often delegated to a ward tender, a green medical officer, or may be to an inexperienced practitioner, and an unsuccessful result is sometimes due to this, much to the patient's disappointment and surgeon's chagrin.

Personal supervision for three or four weeks is necessary, then it may be entrusted to attendants, who are to notify the surgeon on the occurrence of any pain about the wound.

The discharge of pus in a case which has been properly operated on is insignificant in comparison with the

¹ Diseases of the Rectum, third edition, page 43.