hand, if the child became worse, the parents and friends would attribute the worse condition to the early interference, and it would be better to wait twelve months at least.

DR. T. G. WEISENBURG, Philadelphia, stated that the first case mentioned he saw both before and after the operation, and there was a very marked improvement. In the second case the improvement was not so great.

DR. HOWELL T. PERSHING, Denver, Colo., said that while in a disease that leaves such sad results as this one must be on the lookout for every improvement in treatment, it seems that the results of such an operation would have to be very critically examined before one could assign to it any definite value; certainly more than one or two cases would be necessary. Although most of the improvement usually occurs within three months, or at the most within six months, yet occasionally under the use of massage and electricity some improvement occurs in a muscle that has long been out of use, and after such an operation treatment would be apt to be applied with especial care. Dr. Pershing asked how the axis cylinders of the nerve are supposed to act on the muscle. He disapproved of the treatment of all cases of poliomyelitis in the early stages with electricity and massage. In any case of inflammation affecting the nerve centers the indications for rest ought to be observed until the inflammation has completely subsided.

DR. ALBERT E. STERNE, Indianapolis, stated that he sees a large number of these cases and sees them early, and that in those cases where early remedial measures are taken the most excellent results are achieved. It is an absolute error to wait in these cases. No possible damage can be done with ice bags on the spinal column; it is only unfortunate that they exert so little action as they do. They are used in other conditions of inflammation; not to use them here is equivalent to cutting down the wires if the telephone is out of order. Electricity can do no possible harm in these cases seen early after the beginning of the inflammatory process. In these cases active measures do good, and waiting for the process of regeneration is not permissible.

DR. W. G. SPILLER, Philadelphia, stated that they are not at all radical in advocating this method of treatment. They should use it with a great deal of caution, and have said that the cases should be selected. He thinks that neurologists whose attention is directed to the subject will acknowledge that those cases of poliomyelitis in which the paralysis is limited to a small number of muscles are numerous; not a majority, but still numerous. In answer to Dr. Pershing's question, he stated that the subject of regeneration at the present day is a matter of great dispute. A few years ago most neurologists regarded this question as settled, and believed that the axis cylinders grew out from the central stump, and a regeneration from the peripheral portion of the nerve fiber. Later investigators have shown that the peripheral stump takes part in the regeneration. It is probable that the regeneration which occurs in the peripheral stump is not permanent, unless there is union with the central stump. The method recommended depends on the union of peripheral portions of diseased nerve fibers with healthy nerve cells and their axis cylinders. The period that should elapse before the operation is done is, of course, a matter of uncertainty. The longer the operation is postponed the less likely is it to be successful. It seems that six months would be the proper period; perhaps others may prefer a longer period. When this operation is done after a sufficiently long period, when further restoration of motor power is improbable, it can not in any way injure the child. The muscles have already become paralyzed, and cutting the nerve fibers supplying these muscles can in no way injure the child. The inserting of these diseased nerve fibers into the healthy nerve does not injure the nerve if properly done. Not all cases will be successful, but when the operation is successful decided advantages will have been gained over the method of tendon transplantation, because in tendon transplantation the muscle or muscles that have been diseased are thrown out of function entirely and use other

muscles to perform a function which they have not been called on to perform heretofore. Nerve anastomosis restores the function of the paralyzed muscles where it is successful. The operation which Peckham performed in 1900 is remarkable on account of the long duration of the paralysis before the operation was performed and the rapid restoration of power. It differs from the operation here suggested in that healthy nerve fibers were cut and anastomosed with diseased nerve fibers. This Drs. Spiller and Frazier have not proposed, as by this operation, in most cases, one certainly causes paralysis in the region in which these healthy nerve fibers are distributed, and the operation is of very limited utility. Dr. Spiller is in the habit of using massage in poliomyelitis. He believes in all cases of paralysis it is our duty to keep the muscles in as good tone as possible until by natural processes their function is restored. Massage and electricity, he thinks, are useful in all cases of paralysis, especially in cases where recovery may be hoped for by natural means.

INTESTINAL OBSTRUCTION IN CHILDREN.*

JOHN F. ERDMANN, M.D.

Visiting Surgeon to Gouverneur Hospital and Clinical Professor of Surgery, University and Bellevue Hospital Medical College.

NEW YORK CITY.

I shall take up in this paper the chief symptomatic factors in intussusception, strangulated hernia, post-operative bands, etc., Meckel's diverticulum, devoting the major portion of my time to intussusception.

INTUSSUSCEPTION.

This disease, barring a few accidental cases, is one that demands prompt surgical interference. Each successive case I see impresses me with the necessity of making operations of urgency in these cases, and of not temporizing with air inflation or water injections. The diagnosis is not difficult, although the symptomatology, as given in former text-books on diseases of children, etc., should be rewritten with a view of placing all stress on blood or bloody mucous stools, and not on the presence of a sausage-shaped tumor.

In over 60 per cent. of a series of 28 cases seen by me, 24 of which I operated on, no tumor of any kind was palpable per rectum or through the abdominal wall. I do not find, on searching my histories, a single acute case in which blood, bloody stool, bloody mucus or bloody serum was not found, either on the diaper, in the commode, or expelled from the anus after digital examination.

Symptoms.—Sudden pain is almost invariably the first symptom that the child manifests, then a degree of shock varying in intensity, and efforts at defecation take place; these are followed by manifestations of spasmodic pains; a peculiar whining cry or fits of crying. In the first hour or two the abdomen is lax, and no marked evidences of pain are produced on palpation; tumor is not felt unless the intussusception is of considerable extent and is taking place rapidly. After a few hours, depending entirely on the degree of obstruction, abdominal distension and evidences of complete obstruction occur. Although some feces and gas may be expelled with the bloody rectal contents mentioned, they are due entirely to the fact that these contents were below the point of obstruction. When a tumor is felt it is not usually the proverbial sausage-shaped tumor of our didactic days, but is more apt to be round or nodulated and moderately movable. Recently, in oper-

* Read in the Section on Diseases of Children of the American Medical Association, at the Fifty-fifth Annual Session, June, 1904.

ating on a subacute case of six days' duration, a tumor was found that in every possible manner simulated a freely floating kidney.

There is invariably an accelerated pulse and an elevation of temperature of from one degree to two degrees and a half. I feel that the temperature noted in the majority of my cases has been due to absorption of some of the decomposed, foul-smelling, bloody rectal contents that were and are present in every case of over six hours' duration. In two instances the symptoms were of a subacute variety and rather puzzling, no blood having been observed in either case, only a little mucus, and in neither was there complete obstruction noted.

Subacute Intussusception.—Case 20.—(Operative Case 19.) Operated on after a period of six days duration with recovery. I am indebted to both Francis and Joseph Huber.

DR. JOSEPH HUBER'S REPORT.

"N. U. was seen for the first time Sunday, Jan. 25, 1903 (the prior history having no bearing on the present illness, is therefore omitted). The child looked well; temperature was 99. He complained of pain in the abdomen; the pain was paroxysmal in character and not localized in any special area. The mother was positive in her statements that the bowels had moved twice daily up to the previous twenty-four hours.

Codein, gr. 1/6 hourly, high injections, and cold or warm applications were advised.

January 26, restless night; temperature 99, same paroxysmal pain, slight resistance of right rectus, bowels moved twice after injections, a. m. and p. m.

January 27, temperature 96.6, condition the same; restless, with pain in paroxysms all night.

At 11:30 a. m. I saw the boy in consultation for the first time.

Examination.—General condition excellent, pulse good, about 80 to the minute; respiration natural and not accelerated; tongue clean, no odor to breath; heart and lungs negative.

Abdomen: both recti tense, no pains in right iliac fossa; palpation negative, as was also examination per rectum; no retention of urine or delay in the urination; no evidence of shock or collapse. The patient insisted on getting out of bed to use the chamber. Between the paroxysmal pains he would play with his toys and took great pride in blowing a fish horn. No blood or mucus in the stools at any time.

Symptoms.—Pain referred to the umbilical region and tenderness in this area were constant. Pressure over McBurney's point negative. Colicky pains, recurring at irregular intervals and severe in type, caused the boy to roll about and cry out. Both recti tense. There was some interference with the functions of the bowels, as they did not act as readily as in health. No urgent symptoms being present, it was decided to await further developments. If no improvement occurred in twenty-four hours an exploratory laparotomy was suggested, the suspicion tending in the direction of a mild inflammation of the appendix.

January 28: Temperature 100, pulse 104; p. m. temperature 99; restless all night, pains in paroxysms continued. Two good fecal movements followed injections, made by nurse. Dr. John F. Erdmann was now requested to see the patient in consultation, and suggested an early exploratory operation.

January 29: Temperature 99, pulse 110; very restless and in pain all night. Bowels moved well (distinct fecal masses) after injection. The necessary arrangements having been made, the child was sent to Gouverneur Hospital and operated on by Dr. Erdmann at 3:30 p. m.

DR. ERDMANN'S REPORT.

When I first saw the patient, with Dr. Joseph Huber, he had all the manifestations of an appendicitis without any shock. He had had some so-called results from enemata. I suggested to them that the boy be submitted to operation as soon as possible.

Examination.—He was admitted to the hospital the next

morning, Jan. 30, 1903, for operation, and when under ether, on abdominal and lumbar palpation no tumor was found in the appendicular region; but on subcostal palpation, a tumor about the size, and absolutely the shape of an infant's kidney was displaced and could be brought down into the right iliac region. This mass would slip back into the loin exactly as a floating kidney would, but it was evident that the tumor was intra-peritoneal.

Operation.—The abdomen was opened, with a view of exploration, through the right rectus. Digital examination detected a tumor involving the cecum and colon with a slight portion of the ileum. This tumor proved to be an intussusception, which was readily reduced; the appendix was removed, and the abdomen closed without a drain. The patient made a perfect recovery and was discharged to his home in ten days.

This case, without question, was one of those varieties, subacute in character, no adhesions interfering between the separate layers of the intussusception, in which the bowel was not sufficiently constricted to produce the characteristic discharge or to induce complete obstruction. When I saw the child I spent quite a length of time at the history, with the possibility of an intussusception in view; but at this time all manifestations were those of appendicitis, except the constitutional manifestations due to pus, which one would naturally suspect at the end of five or six days, and I therefore suggested, as above stated, exploratory laparotomy. The question of appendicitis was readily eliminated when the child was under the anesthetic.

Movements of the bowel occurred with mucus, spasmodic pains with intervals of from one-half hour to an hour, a tumor not palpable until anesthetized, and then simulating in size, shape and motility a floating kidney. The condition only demonstrated by exploration.

CASE 28.—(Operative Case 24.) The second case was one of eight days duration in a girl 22 months old, seen on the third day by one of our leading pediatricians. A diagnosis was not made. The case was seen by me on the eighth day, with the following signs and symptoms.

Symptoms.—A history of spasmodic pains, the child getting in the knee-chest position and crying out at intervals of thirty minutes. In the interim the child was either quiet or whining. Mucus was present, but no blood. Movements of the bowels occurred daily, but were due to rectal feeding. There was a movable tumor in the right side, but having no distinct outline. The child was in shock.

A diagnosis of intussusception with subacute symptoms was made.

Operation.—At the operation a gangrenous, irreducible, ileocecal intussusception was found. The appendix and ten inches of the ileum were excised. End to end suture anastomosis was employed. The patient died in six hours.

In this case I was satisfied that an intussusception existed although an appendicular mass had been suspected by the attending physicians.

Spontaneous Reduction.—Case 26.—A short time ago a case was referred to me by Dr. Francis Huber at about 11:30 in the morning. As I was unable to reach the house before 2 o'clock, I sent my nurses and assistants to prepare for operation so that no time should be wasted. On their arrival at 1:30 o'clock, they were told that the child had just had a distinct fecal movement, so that operation would not be necessary. The history of the case is as follows:

History.—Baby, 5 months old; was well up to 6 p. m. of the preceding day, when suddenly it was taken with a rectal discharge of pure blood, followed by vomiting. The mother said the child had agonizing pains, occurring at frequent intervals. When seen by Dr. Francis Huber at 11 o'clock of the following morning, i. e., the morning after the onset, the child was greatly depressed; the abdomen greatly relaxed; no stools since the preceding day; no tumor was palpable externally;

rectal examination was followed by the escape of about one ounce of offensive bloody discharge; no mass to be felt in the rectum or bimanually. The history further showed that an enema had been given some time before midnight, but nothing in the line of injections had been used following this time until the patient was seen by Dr. Huber. He, feeling that the symptoms were so pronouncedly those of intussusception, deemed it inadvisable for further interference other than surgical.

I was then called and ordered all preparation, as above stated, when, at 1:30 I was telephoned of the distinct fecal movement having taken place, and that the child's general condition was that which we see after a reduction of intussusception.

This spontaneous reduction evidently took place somewhere between twelve and fourteen hours after the injection had been used, and followed within an hour after the abdominal and rectal examination made by Dr. Huber. The mother said that from the time of the examination the vomiting and pain ceased, and that between 1 and 1:30 p. m. the bowels began to act.

The after-history of this patient was also typical of the cases we have reduced by operative procedure—that is, there was a foul-smelling, greenish diarrhea, which is almost invariably present, and is due to a superficial necrosis of the mucous membrane. In a comment by Dr. Huber on this case to me later, he said that in view of the history, symptoms and appearance of the baby with its foul enteritis (characteristic), he concluded that this was a fortunate case of spontaneous reduction, feeling that there could be absolutely little doubt on this score. I, personally, feel that this case was reduced largely through the manipulation, abdominal and rectal palpation, which simulated a massage. I certainly do not feel that the enema given about midnight could have had any influence whatever at so remote a time.

In the 28 cases seen by me twenty-four were operated on. One operative treatment was refused. Two are recorded as reduced shortly after the use of large enemata with inversion. One was recorded as a case of spontaneous recovery (see case No. 26) twelve or fourteen hours after injections had been used.

TREATMENT.

Rectal enemas in the first few hours are not productive of harm, but may, though rarely, be followed by reduction. This occurred but three times in the 28 cases seen by me.

Admitted that one meets occasionally with a reduction by the use of enemas, nevertheless this very important fact must be remembered, i. e., that the whole mass may be reduced but the ileocecal junction and one or more inches of the ileum. This being so, all the symptoms clear up for a time. They again return and necessitate operative interference at a time when the conditions of this region are not nearly so favorable to reduction, and may even require excision. This condition was well demonstrated in my sixteenth case, a male child, seven and one-half months old. Duration eight and three-fourths hours, injection used by the father, a well-known pediatrician, Dr. H. M. Silver, and myself was followed by a perfectly tranquil picture, but realizing the possibility of this condition of incomplete reduction, we all felt that the risk of exploration would be far less than that taken by leaving matters rest for farther manifestation. That our fears were not groundless was proved by the evidence of a mass at the cecum, which consisted of almost three inches of the ilium and the entire appendix. Reduction was readily accomplished, the appendix was removed, and recovery followed.

I should, therefore, not feel satisfied that reduction

had taken place, even if the child should have no further manifestations of pain, etc., unless this was followed in a short time by a movement of the bowels that we could feel satisfied had come from beyond the supposed site of the previous intussusception; and that in the waiting interval for a movement we should be prepared to proceed at a moment's notice with the operative measures.

The use of enemas previous to operating is offered as a suggestion, and I can say that clinically it is of very valuable assistance, as there is no doubt that the injections reduce a portion of the intussusception, and any agency or procedure that limits operative time must of necessity diminish shock in direct proportion.

Although these little patients bear operative interference quite well, it is quite significant that my greatest mortality rate occurred in those cases which required the longest time for reduction, and that these cases were also the ones of longest duration. Enemas as a means of reduction are not advised after six hours time has elapsed from the first positive symptoms, as, after this period, much valuable time is lost by such procedure.

Anesthetic.—Chloroform has been the anesthetic used in all cases operated on by me, for the theoretical reason that vomiting may be a causative factor in producing this condition; therefore, the anesthetic that is the least liable to induce postoperative vomiting, which might reproduce the intussusception, should be used unless its danger to life in other ways is greater.

Operation.—The abdomen is prepared as in all emergency operations. The incision is invariably made by me in the middle third of the right rectus, and does not usually exceed two inches in length; the index finger explores the abdomen and then, if the tumor can not be delivered on the abdominal wall or into the incision, the latter is enlarged. Should the tumor be of considerable size, it is advisable, and even necessary, to reduce it in part as it lies in the abdomen, by compressing the distal end of the tumor in the proximal direction, i. e., so that the escaping intussusceptum can be seen or felt escaping toward the cecum in colonic or ileocecal varieties, and toward the duodenum in the enteric variety.

Traction as a means of reduction should never be used except in the lightest possible manner and then only with a view of directing the expulsion of the intussusceptum in its proper course.

Sometimes reduction occurs with a distinct pop; this is seen in the colonic and enteric varieties, but when the ileocecal valve is the part of the mass, great difficulty is often met with in reduction of the final portion. A little patience and gentle compression, aided with a backward and forward motion of the unreduced mass, allied to attempts at reduction and production of the condition is usually followed by success.

Occasionally a blunt instrument introduced between the intussusciens and the intussusceptum is of benefit in breaking up any adhesions, and may also be employed to dilate the tonic, contracted intussusciens.

When, as reduction proceeds, the tumor advances toward the right iliac region, it is advisable to bring the mass out of the abdomen before finishing the reduction. In each case, twelve in number, that I have operated on since January, 1902, I have removed the appendix before closing the abdominal incision, and in all but two of these the appendix was a portion of the intussusception. My reason for making the incision in the right rectus is due to the fact that the majority of cases

are those invading the cecum and colon, and such being the case, the most frequent site of final reduction is that of the right lower quadrant of the abdomen. Even if the case is one of the sigmoid or enteric variety, this incision can be enlarged sufficiently to approach the most remote parts of the abdomen in children, without subjecting the patient to any danger of a hernia.

In case of inability to reduce the intussusception or when gangrene of the gut exists, excision with end to end anastomosis in the enteric or colonic varieties, and end to side anastomosis in those cases requiring excision of portions of both large and small intestines is necessary. Although occupying a few moments more time, this should be done before a patient is taken from the operating table.

In one case of over five days duration, in a male four months old, I was compelled to excise for gangrene and irreducibility six inches of the ileum and all of the large intestines to within four inches of the rectum, making an anastomosis between the ileum and the lower portion of the sigmoid. Death followed in a few hours.

Before closing the abdomen care should be taken to examine thoroughly the abdominal contents for multiple intussusceptions; the term multiple is used in contradistinction to the terms double, triple, etc., given by Edred M. Corner, M.B.,¹ to those cases in which double, etc., invagination takes place in one intussusception.

After reduction, the abdominal incision is sutured in the usual manner and a very tightly applied abdominal dressing is used.

After-treatment.—The after-treatment is of great importance. Purgative, in sufficient doses to control peristalsis, is to be administered every three hours; barley water and milk, proportions varying with the child's age, are employed, and a small amount of bismuth, etc., is used.

The first day or two the child may have all the evidences of a foul, putrid enteritis with a temperature of from one-half a degree to two degrees elevation, the latter being thoroughly consistent with primary union in the wound.

My work for the past fourteen months consists of six operations with two deaths. One of these patients was doing nicely for twenty-four hours when he suddenly died. Dr. Francis Huber, to whom I am indebted for the case, was of the opinion that death was due to status lymphaticus. The second case was that cited as Case 28, operative case No. 24, where I excised ten inches of the ileum, etc.

I can not but feel that the citation of the two cases, one of spontaneous reduction several hours after enema, and the one of a subacute variety of over six days' duration with recovery after operation, are of sufficient interest to repay us for the time occupied in recording them in detail.

STRANGULATED HERNIA.

In a recent paper² on strangulated hernia, in which I reported 58 cases, two were in children under five years. Since reading the paper I have had another, occurring in a male six weeks old, son of a physician. Two occurred on the right side, while the third was on the left. All were inguinal and occurred in males.

The left-sided hernia occurred in an infant not quite 13 days old. The symptoms expressed were those of irritability, crying, restlessness and shock. The pro-

trusion, readily discoverable, is usually found by the mother while giving the child its bath. No differential diagnostic difficulty is experienced in these unless the case be one of atypical Richter hernia as occurred in the patient of 5 years of age. In this case obstruction was evident for several days. A small nodule in the groin, said by the mother to have been there for a considerable time, never varying in size, was taken for an enlarged gland. An abdominal section was made, and the protrusion found to be a small portion of a knuckle of gut (Richter's hernia), strangulated and gangrenous, requiring a resection. The case of a male, aged six weeks, was reduced after manipulation, no operation being necessary.

Since recording the 58 cases of all varieties of strangulated hernia, I have had 7 more, including the child of six weeks, cited above. There were 5 inguinal and 2 femoral, making a total of 65 cases, giving us about .046 per cent. occurring in children under 5 years in strangulated hernia of all varieties. If the inguinal variety alone be considered, and this should be, as I do not know of any strangulations recorded in the literature of femorals, ventrals, etc., in children of this age. I then have 31 previously reported and 5 additional cases in this paper, making a total of 36 cases in my recent practice, of which 3 were in children.

We then find .083 per cent. of all inguinal strangulations occurred in children under 5 years. The treatment in these cases resolves itself into: 1, Reduction, if possible, and in recent cases when the child is under 2 years the use of a properly fitting truss, relying on the possibility of a spontaneous cure. The six-weeks-old child has never had a recurrence in the six months past since reducing the hernia. 2. Operation with radical cure in all irreducible cases and all cases over 2 years, as the chance of spontaneous recovery with truss is relatively infrequent after this period.

POSTOPERATIVE BANDS, ADHESIONS, ETC.

It has fallen to my lot to have 2 cases of obstruction due to the above causes—one in a boy 6 years old after an operation for appendicitis in a Brooklyn hospital, and the other in a boy 10 years old following a drain operation for appendicitis.

Symptoms.—The symptoms of obstruction are the usual ones—constipation, meteorism, singultus, increased pulse, bad facies, temperature, vomiting as time goes on becoming fecal.

These symptoms are readily assigned to their proper source by fact of the history containing evidence of a previous operation. Both of these cases occurred after a period of from five to seven months had elapsed since the time of the original operation. This period of time corresponds to that cited in quite a number of cases reported in the New York Surgical Society, May 11, 1904.

Treatment.—The treatment in the cases occurring early, i. e., in from two to four weeks after operation, consists of elevation of the hips and body, high injections, etc. Should these not accomplish the result desired, operative means must be undertaken.

In cases occurring late, i. e., after from four to six weeks, precious time is lost by any attempt at mechanical relief, and operative procedures should be promptly instituted.

Prophylaxis.—As a means of preventing adhesive obstruction, I deem it essential to have a movement from the bowel of the patient within from twenty-four to thirty-six hours after the operation and one or more

1. Pathology and Classification of Intussusception, *Annals of Surgery*, 1903.

2. Medical Record, March 12, 1904.

daily after that, and careful watching for not less than three weeks.

MECKEL'S DIVERTICULUM.

These abnormal conditions can produce obstruction in one of three ways, an instance of each one having been reported by me in past years.

First, the diverticulum may become inverted and act as a drag producing an intussusception. Case 2 in my list of intussusceptions.

Second, by strangulation by cords or band-like prolongations; two cases in young adults.

Third, by inflammation simulating appendicitis and producing obstruction either by means of parietal or visceral adhesion of the distal end, thereby forming a loop for strangulation; one case of a young adult reported by me; or by producing a general peritonitis with subsequent obstruction.

Symptoms.—The symptoms of obstruction by Meckel's diverticulum in types two and three are very similar to an appendicitis plus in a day or so a profound obstruction with or without a generalized peritonitis; while in appendicitis obstruction is not evident except in those cases accompanied by general peritonitis or in those exceptionally rare cases where the appendix overlies a portion of the intestine and acts as a band-producing or cord-producing obstruction.

I feel safe in saying that close observation of the literature of this subject will show that the great majority of cases have heretofore been operated on for appendicitis, and the actual condition was only found on exploration.

OBSTRUCTION FROM OTHER CAUSES.

I do not feel that the subject of obstruction in children has had proper attention without considering the obstruction peritonitis seen occasionally in pneumonia cases in which no visible lesion of the appendix or other abdominal viscus is evident. These cases simulate very closely those of intussusception or general peritonitis from appendicitis, and are usually differentiated by the fact that a pneumonia exists; that respirations are increased; that the appearance is of a livid or red-cheek character instead of pale and anxious, as in acute obstruction; temperature is higher than usual in obstruction; that no blood has been passed per rectum; that no mucus is present in the stool; that no mass is found in the right iliac region.

At the same time it must be remembered that a sufficient amount of fluid may be present in the peritoneal cavity to give a sense of fluctuation per rectum. The citation of the following case will impress itself on all: Recently I saw a case with one of our pediatricians for intussusception, in which I made a diagnosis of pneumonic peritonitis.

This patient was about 8 months old, had a lobar pneumonia, and with it a profoundly distended and rigid abdomen, complete obstruction, rapid and feeble pulse, and temperature of 102.5. At the time of the consultation I suggested exploration, although I felt that neither intussusception nor appendicitis was present. On returning in two hours with my assistants I found the child practically moribund, and refused to operate. Death followed in about two hours. Autopsy, abdominal only, revealed a general peritonitis; no other visceral involvement; appendix healthy.

The onset in these pneumonia peritonitis cases resembles an appendicitis, especially if the lower lobe be involved, pain is reflected down into the abdominal wall through the eighth, ninth, tenth and eleventh inter-

costal nerves, giving us rigidity over the spot seen at times in appendicitis cases.

It must be also admitted that a pneumonia and an appendicitis can occur at the same time. Recently I have seen such a case; fortunately the appendicitis complication became an enclosed abscess, and was successfully operated on after the right lower pneumonia had subsided.

60 West Fifty-second Street.

DISCUSSION.

DR. A. JACOBI, New York City, asked Dr. Erdmann the age limits of the patients on whom he had operated; if he *always* removed the appendix, and, if so, why.

DR. ISAAC A. ABT, Chicago, believes that all these cases are not due to mechanical obstruction of the bowels; there are cases of paralytic obstruction which deserve consideration. We have all seen examples of complete intestinal obstruction in pneumonia and in the course of severe intestinal infections which proved on autopsy to be paralytic in nature. Two years ago a colleague asked Dr. Abt to see his child. She was suffering from an acute gastrointestinal infection. The child vomited and had loose stools for a time, later on this was followed by a condition of absolute obstruction. There was complete prostration; the face was ashen; the child refused all nourishment; the temperature was normal—at any rate, less than 100—the pulse and respirations were very rapid. The abdomen was enlarged and tympanitic. A surgeon was called in during the night and pronounced the case one of intussusception. He was mistaken, however, as the autopsy a few hours later showed. During the earlier stage of the child's illness the stools had contained no blood; simply large quantities of mucus. The child died, and at the autopsy every inch of the intestine was carefully gone over. It was much distended, but no obstruction was found. The liver was fatty, and on section resembled that found in fat necrosis. Dr. Abt has seen two cases recently that were absolutely similar to this one. The striking features were, first, the symptoms of intestinal infection; then the absolute obstruction of the bowels, the low temperature, the rapid pulse and respiration, the fatty degeneration of liver and other viscera, and death. He believes that these cases belong to a class which has not been sufficiently described in literature, and that they are the result of ptomain poisoning, depending on the ingestion of some kind of poisonous food. He does not believe that such cases of obstruction can be helped by surgical measures.

DR. LOUIS FISCHER, New York City, said that the results that he has seen in operations on children have not been so good as those reported by Dr. Erdmann, and he attributes this to the fact that children do not bear the shock of laparotomy well. He has seen a series of cases in which laparotomy was done, and all the patients died. The fatal result in these cases might be attributable to several causes, one of which was a slow operation and shock. He would like to know whether rapidity of operating is not an important factor, so as to lessen the shock. The following case came under his observation recently. A child, apparently well in the morning, was taken suddenly sick in the afternoon with symptoms of constipation and considerable abdominal distension. In the evening the abdomen was still more distended. He saw the child at 9 o'clock, 12 hours after the first symptoms appeared, pronounced the case one of intestinal obstruction and suggested an operation. The operation was delayed till the following day and resulted fatally. The child was moribund at the time of operation.

DR. S. W. KELLEY, Cleveland, gave it as his opinion that spontaneous reduction occasionally occurs and may take place under the use of opium. If there is any time at which opium is useful in intussusception, it is in the beginning, but it should not be depended on. There are cases in which the use of air or water as a means of reduction is useful, and sometimes it must be adopted as a matter of necessity, as a laparotomy is not always permitted by the parents. The introduc-

tion into the bowel of air or water, when properly done, is sometimes followed by reduction, and Dr. Kelley does not think the method should be altogether dispensed with. Very little is said in regard to the way in which it should be applied, the amount of pressure that should be used, nor the length of time the pressure should be maintained. Dr. Kelley thinks the figures that Morris gave some years ago placed the pressure at 8 or 9 pounds, which appears to be too high. He believes that air pressure is preferable to water pressure in very many cases, but it is so difficult to measure that he finds it succeeds best by using air first and then following it with water, which is more easily handled, and by means of which the air pressure can be measured. He prefers a pressure of four or five pounds, keeping it up for a period of 15, 20 or 25 minutes, and believes that this is more effective than a higher pressure for a shorter time.

DR. SAMUEL MCC. HAMILL, Philadelphia, said that it has been shown that irritation of the rectum produces reversed peristalsis, and this fact may have had some bearing in producing spontaneous reduction in the case reported by Dr. Erdmann.

DR. A. JACOBI, New York City, considers that the phenomenon mentioned by Dr. Hamill would not be of any practical value in the treatment of these cases, because the reverse peristaltic wave would undoubtedly be interrupted at the point of intussusception.

DR. JOHN LOVETT MORSE, Boston, reported that the experience at the Infants' Hospital in Boston is that babies not only bear laparotomy but other severe operations very well, providing they are kept warm during the operation. All the patients who are to undergo an operation are wrapped in cotton and placed on a special operating table, which is kept warm.

DR. JOHN F. ERDMANN said that eight or ten years ago Dr. R. T. Morris produced reversed peristalsis by irritating the rectum by the use of chlorid of soda, to determine what portion of the intestine he was dealing with—whether the afferent or the efferent portion. Dr. Erdmann considers that it is not only time lost, but that it is very apt to prove useless, because the colon and the rectum will not hold the amount of pressure we wish to use. The sphincter will not hold the water in the bowel. With both air and water the danger of producing a rupture of the bowel is very great. If Dr. Kelley could see the difficulty Dr. Erdmann has experienced in reducing an intussusception with the fingers after the abdomen is opened he would never again resort to the introduction of air or water. The coats of the intestine are soggy and edematous, and unless great care is exercised in the reduction, a rupture of the serosa and musculosa will occur. While Dr. Erdmann does not favor the use of injections, he would not object to their use during the first six hours. Among his last six cases there were two deaths. In one, the cause of death was given as status lymphaticus; the other patient died after an extensive resection, with end-to-end anastomosis. There is no question that the more rapid the operation, the less the shock, and the latter is frequently the cause of death. The youngest child he has operated on was 14 or 15 weeks; that patient recovered. Children, as a rule, bear operation well. He has recently operated on 22 cases of appendicitis, all under 5 years of age, with a mortality of 9.9 per cent., and at least 14 of these cases were of the septic type. In the majority of his cases of intussusception, the operation was done within 15 or 20 minutes. If those cases that he operated on were not mechanical, then he does not know what constitutes a mechanical obstruction. An intussusception will occlude the caliber of the gut as completely as a wad of cotton, and even more thoroughly. The paralytic type of obstruction does occur, particularly in pneumonia in children, when there may be a general pneumococcus peritonitis, with paresis of the bowels, and death. The majority of his patients were under 1 year old: two of them were over 5, and the rest under 3. He removed the appendix because it could be done without complicating or lengthening the operation to any extent. He thinks it wise to do so because it prevents the formation of subsequent adhesions and a second abdominal section.

CHOICE OF A TIME OF ELECTION IN MASTOID OPERATIONS.

SOME CONSIDERATIONS ARISING FROM THE DIFFICULTIES OF THE CHOICE—PROSPECTIVE RESULTS.*

D. A. KUYK, M.D.

Lecturer on Diseases of the Nose and Throat, Medical College of Virginia.

RICHMOND, VA.

Although voluminous articles, reports and discussions have appeared on the subject of mastoiditis, the mastoid operation and its ultimate results; indeed, experiences so exhaustively detailed that it would seem the searcher after truth might find his way without groping, yet perplexities that defy satisfactory decision so constantly confront the otologist, that I have chosen to place before you some such perplexities as have arisen for me in the recent past; and especially the choice of the proper moment for most advantageous operation.

This paper deals with well-to-do private patients, with whose intelligence it is undoubtedly more difficult to co-operate than in the case of the ordinary grade of dispensary subject. For, adding, in the former instance, to the difficulty and confusion which the conscientious worker experiences at times in his own mind, and at a moment, too, when most he needs to be unhampered in power to decide, there are presented to him by relatives and friends such questions and demands for positive answers that the picture of uncertainty is complete for him. While in the latter instance with the usual dispensary patient, one may follow a more independent and assured attitude, more easily ignoring inconvenient demands for enlightenment on the problems involved until one has arrived at some satisfactory solution of the best course to pursue.

So varied, too, are the final chapters in the histories of these cases, even after the otologist has had freedom in following his best intelligence, that they are at times quite inexplicable to him. Although reports of cases, like statistics, may be made to serve most contradictory ends, and are therefore not so useful as they might be, yet I wish to enforce and illustrate my theme by a few experiences which I will relate as briefly and relevantly as possible.

For the sake of logical discussion I have grouped these cases as follows:

Group 1. Features obtaining in mastoiditis as a complication or a sequel of the exanthemata.

Group 2. Features obtaining in mastoiditis as a complication or a sequel of influenza.

Group 3. Features obtaining in mastoiditis as occurring in the course of chronic suppurative otitis media.

As an object lesson looking toward the more intimate discussion of each of the above-mentioned groups it becomes necessary to study briefly the histories of cases apparently similar, yet with results widely divergent.

REPORTS OF CASES IN GROUP 1.

CASE 1.—Female child, age 6 years. Scarlet fever of severe type of six weeks' duration. Subsidence of all symptoms except slight fever never ranging above 100.5 degrees. Without premonition, many days after apparent convalescence set in, pain began in the right ear.

Examination.—This revealed a highly congested, slightly bulging drum membrane, no sagging of the posterosuperior canal wall, mastoid sensitive to pressure.

* Read in the Section on Laryngology and Otology of the American Medical Association, at the Fifty-fifth Annual Session, June, 1904.