SUGGESTIONS FOR MEDICO-PAROC- CHIAL REMUNERATION.

By F. H. SANKEY, *Esq., Wingham.*

I AM inclined to think, that a payment of a fixed sum per case, is the most eligible way, as then the amount of remuneration will always bear some proportion to the amount of services rendered. I am not prepared to say what that sum should be, as it should of course vary with the density of the population, and the greater or less distance at which the patients are situate from the medical man. The knowledge that a certain expenditure is caused by every order that is given, acts as a check on the relieving officer or overseer, and prevents that indiscriminate giving of orders, which was the great cause of driving all the poor to the "parish-doctor," under the old system, while the smallness of the amount does not prevent the order being given to the real pauper as soon as he is taken ill;—the difficulty is to fix the amount, but I think, in general, ten shillings per order would be sufficient remuneration, provided that, in every case of illness of long standing, the order should be renewed every three months. This, I think, would be better, than fixing a larger sum, as, in some small parishes, there might be severe chronic cases, requiring a good deal of medicine (if not of attendance), and, perhaps, very often, few acute or epidemic cases to counterbalance them; on the contrary, if the sum per case were high, one family labouring under scarlatina or measles (as they would have an order for every individual, however slight the disease might be), would amount to what the guardians would consider too high a remuneration.

I do not think that the plan of providing drugs from a different source would be any improvement, unless in towns. I am satisfied that, in remote country districts, it could not always be accomplished.

We have two *well* regulated medical Provident-Societies in this neighbourhood,* *not* established according to the recommendation of the Poor Law Commissioners, and I am satisfied, in spite of all that has been said against them, that they are acting well for the medical man and also for the poor: the numbers who belong to them are increasing every year.

* The subscriptions of the poor to these institutions are three, four, or even five times the amount of those to the Medical Club established by the Poor Law Commissioners.

ON THE INFLUENCE OF GRAVITATION ON THE CIRCULATION OF THE BLOOD IN HEALTH AND DISEASE.

By FRANCIS MOSLEY, *Esq., Surgeon, London.*

ALTHOUGH the presiding and controlling influence of a vital principle, renders the actions, movements, and effects of living bodies, at variance with the laws governing inanimate matter, under very many circumstances this control is not paramount.

We know that the influence of gravitation is never for a moment suspended, either with regard to living or inert matter: we can conceive no state of organization capable of maintaining an existence independent of its power; motion is the effect which renders its operation, as a cause, perceptible to us.

Inert matter moves in obedience to the impulse communicated, without offering any assistance of its own.

Living matter, when moved under certain circumstances, possesses the faculty of modifying the impulses according to the organization peculiar to its mode of existence.

Thus each organ finds in this agent an assistant, or antagonist, according to the necessity of the action to be performed.

As the true antagonist to the soaring eagle's wing is gravity, so the circulation through all parts of the body is, unquestionably, affected by the influence of the gravity of the blood, but in different degrees, according to their situation and position; and, as a general truth, we may assert, that whenever the venous circulation is favoured by the gravity of the blood, under ordinary circumstances, there congestion will be apt to take place by any change of the accustomed position, such veins having but little contractile power to aid in propelling their contents.

Whenever the upright posture is changed for another, say the horizontal, the circulation is very perceptibly affected; the veins of the neck become swollen and livid; the carotid and temporal arteries pulsate with greater force, and headach and confusion of thought are often induced. These occurrences are still more rapidly and strikingly developed if the head be placed lower than the rest of the body.

Some of the severest cases of cephalic disorder have come under my observation, emanating solely from the head being unsupported when in bed by the usual way. This, though apparently a trifling circumstance, I am led to consider, merits closer attention from us in practice than is usually bestowed; an opinion which a moment's physiological view can but confirm.

It is not unfrequently observed, in elderly patients who have been, from whatever