IX—Landry's paralysis.	} Microbe analogous to	} Marinesco. Marie.
	} the bacteria of car-	
	} buncles.	

KRAUSS.

Epithelioma of the Cranial Vault. In the *Gazzeta degli Ospedali*, Dec. 21, 1895, Dr. Buel reports a case of epithelioma of the cranium in a man sixty-nine years old, who at two years of age was badly burned about the head leaving a large partially healed cicatrix. During his whole life the patient presented a large superficial ulcer the size of the palm of the hand, covered with crusts. In scratching the head the scabs were removed, granulations appeared with ichorous secretions presenting all the characteristics of an epithelioma, the glands becoming enlarged, cachexia, etc.

The cerebral symptoms present were, cephalalgia, vertigo, convulsions of the lower extremities, then paresis of the left leg and complete hemiplegia sinistra. Death occurred two months after entrance into hospital. Autopsy showed perforation of the cerebral theca at level of the middle part of the superior longitudinal sinus. The meninges were perforated and the mass extended into the right frontal ascending convolution and into right lateral ventricle, in which was found some pus. Histological examination confirmed the diagnosis of epithelioma. The bibliography shows but ten cases of this nature.

KRAUSS,

THERAPEUTICAL.

Acromegaly Treated by Thyroid. At the Berlin Medical Society, Nov. 27, 1895, Professor Mendel presented a typical case of acromegaly, in which the only peculiar symptom was typical double temporal hemianopsia. This hemianopic condition, which is a not common complication of cases of acromegaly, was considered by Professor Mendel the result of pressure exerted by the pituitary gland upon the optic chiasm. The thyroid gland tablets producing no effect, the patient was given one gramme of pituitary gland daily for a fortnight. The time was too short to judge of the results, but it was stated that the patient could close her mouth better than before, and the knee-jerk, which had been lost, reappeared upon the left side.—*Sem. Med.*, Dec. 6, 1895.

MITCHELL.

Insomnia and Its Treatment. Maurice de Fleury (*Gazette des Hôpitaux*). In all cases of insomnia not arising from acute pain or irritation of the meninges (tumor or meningitis), it is possible to employ dynamic treatment, which will be efficacious and easy of application, instead of hypnotics. A study of arterial pressure at the radial pulse in those suffer-

ing from insomnia demonstrates that this pathological phenomenon coincides either with a state of marked hypertension or excessive hypotension. Without the aid of drugs, and by the employment of simple physical procedures, it is easy in the great majority of cases to restore the arterial tension to normal and at the same time to cause sleep. It is always useful to add to this physiological treatment, a psychological one, and cause a habit of sleep. There are two kinds of insomnias,—the neuro-pathic, in which suggestion is applicable and the more serious form where the treatment by opium or chloral is justifiable. The insomnia of neurasthenia, that occurring during convalescence from typhoid fever, and that accompanying a condition of low arterial tension, all have the same pathogeny. When a patient with asystole cannot sleep, it is because the arterial pressure is low. If his feeble heart be relieved with digitalis, sleep at the same time becomes possible. In a great number of cases one can substitute the dynamic agents, such as massage, frictions, douches and transfusions, and at the same time caffeine and digitalis may be employed as heart tonics.

FREEMAN.

**Idiocy Without Myx-
œdema Improved by
Thyroid Treatment.**

Dr. Netter, at the Biological Society at Paris, Nov. 30, 1895, read a note on a case of idiocy without myxœdema treated successfully by thyroid. The patient was a child six years of age, progressively cachexic, with puffiness of the face, and thickening of the skin of the legs. The cachexia, the œdema, the skin thickening and the trace of a special albumin, which was found in the urine disappeared under the thyroid treatment. No trace of the thyroid gland could be detected in this child.—*Sem. Med.*, Dec. 6, 1895.

MITCHELL.

**Methods of Prepar-
ing Thyroid Extract.**

At the Société de Thérapeutique, on the 11th of December, 1895, there was a discussion as to the value of the different methods of preparing thyroid extract. M. Vigier related his plan, which was as follows: He uses sheep-thyroid, mixing the fresh pulp with pulverized sugar. It is further mixed with carbonate of soda, and with charcoal, in order to hinder decomposition. This preparation has been successfully administered by a number of physicians. In the discussion which followed his brief statement there was a general agreement that the fresh preparations, while more troublesome, were more active than the tablets, extracts and powders. The latter, of course, have the advantages of being always accessible, and fairly constant in their effects. In one case which Dr. Marie treated for six weeks with the natural thyroid, the results were excellent, but after the suspension of treatment for six weeks the symptoms reappeared. In the second course of treatment, preparations of the gland were given, made by evaporation *in vacuo* with the addi-