was thought fair and proper that they should recover. The third case, however, might be considered a test of good management from the beginning. This case, according to statistics, had less than an equal chance. If a mistake had been made by an indefinite treatment, it would have jeopardized the life of the patient at once. But the fourth case, 70 years old, with malaria and alcoholism, would not recover according to statistics. During a course of an indefinite *individual treatment* something might have been done which would prove injurious to the patient; but by the plan here proposed, *founded upon the mechanism* of the disease, it can be safely assumed that nothing injurious would be done; but, with the frequent cuppings to prevent heart failure, the supporting diet and personal hygiene; besides the continued medication of the air, and the inhaling of antiseptics; with the two resolute ammonias and the antifebrin; with the antiseptic resolvents applied hot externally, together with a good moral support, we may leave this desperate case as doubtful, yet not without hope.

No mention has been made in the course of treatment of the pneumonia "*foudroyant*," which kills the patient almost like a thunderbolt, because these cases are scarcely recognized before the overwhelming catastrophe occurs. What should be promptly done in these cases is general venesection if one is sure of his case, but at any rate extensive cupping, to prevent the overwhelming congestion. At the same time medicate the air of the room, and use stimulating inhalants, with a view of reviving the patient, to divert the blood by again and again frequent cupping, so as to preserve the heart; and if these, happily, may stay the progress of the disease long enough, then, to commence the plan of treatment recommended.

CONCLUSIONS.

1. The mechanical interference to the circulation and aeration of the blood in the lungs is the main symptom to be combated in the treatment of pneumonia.
2. It requires prompt, urgent and aggressive treatment in the earlier stages of the disease, and this mechanical relief can only be afforded by decided local depletion.
3. Even after the effusion into the parenchyma has occurred, it requires still further local depletion to preserve the heart, and keep open the circulation in the roots of the lungs.
4. Pneumonia admits of study by mechanical as well as vital principles, and its pathology may be elucidated and its treatment improved by study in these directions.

THE CALIFORNIA BOARD OF EXAMINERS, after March, 1891 will require all who seek to practice medicine in that State to be graduates of a three-year graded medical college.

INTRAPERITONEAL RUPTURE OF BLADDER.

Read in the Section on Surgery at the Thirty-ninth Annual Meeting of the American Medical Association, Cincinnati, Ohio, May 8-11, 1888.

BY H. H. GRANT, M.D.,
OF LOUISVILLE, KY.

Abdominal surgery, though more than fifty years a thrifty identity, has in the last half a decade assumed such a position, through the boldness and success of skilled operators, as to surpass beyond competition all other departments. Even the newly invaded region of the brain, with its brilliant record, is unmentionable in comparison.

Though among the rarer lesions for which abdominal section is appropriate, the still rarer comparative success after the operation lends an interest to the few recorded recoveries after laparotomy for intraperitoneal rupture of the bladder, beyond numerical desert.

In making report of a successful suture of intraperitoneal rent, I venture to call attention to the history of the operation, now about ten years old. Though the classical opinions expressed by Sir William MacCormac,¹ together with the comments and amplifications by Mr. T. Holmes,² cover pretty well both the theoretical features and the practical details of the condition and operation, yet besides the reports of several operations since Mr. Holmes' paper, there remain for elaboration and systematization many salient and important points.

It is a remarkable fact that, though indefinite reference to the propriety of attempting suture of the bladder after laparotomy for rupture has been made by Gross,³ Larrey, Cussach and others, yet, except the suggestions of Holmes in his "Principles and Practice of Surgery" in 1875, nothing like a definite position is mentioned in any textbook on surgery published prior to 1880. Up to these dates the treatment of such condition is dismissed with a few words as to a hopeless expectancy. In a full review of the subject as late as 1886, Mr. Rivington⁴ declares that up to that time intraperitoneal rupture of the bladder, however treated, had been uniformly fatal, excepting only Walter's⁵ case of laparotomy without suture, at Pittsburg, in 1862.

This is perhaps the only laparotomy for the lesion under consideration until Mr. A. Willett,⁶ in 1876, closed by suture of the bladder rent and abdominal wound. It appears that to Mr. Willett is due credit not only for priority in operation, but also in detail of suture of the vesical rent—as he disclaims⁷ knowledge of previous suggestion pointing out the procedure. Interrupted silk su-

¹ London Lancet, December 11, 1886.

² London Lancet, July 23, 1887.

³ MacCormac, London Lancet, December 11, 1886.

⁴ Heath's Surgical Dictionary, 1886.

⁵ Med. and Surg. Reporter, Philadelphia, February, 1862.

⁶ St. Bartholomew Hosp. Rep., 1876, xii.

⁷ Holmes, Lancet, July 23, 1887, p. 154.

tures were used in the bladder twenty-nine hours after injury. Urine escaped, however, through the rent, and death from peritonitis and shock took place in twenty-three hours. In 1878 Mr. Christopher Heath,⁸ forty hours after accident, closed an intraperitoneal rent with continuous catgut suture. Leakage and extravasation occurred, and death on the sixth day.

In 1883 Pilcher,⁹ commenting on these cases, describes with references to Vincent and Stein, the

operation afterward successfully done by MacCormac¹⁰ in two cases in 1886, and now recognized, with slight modification, as the perfect technique. In a table prepared by Sir Wm. MacCormac,¹¹ published in May, 1887, explicit and concise details are given of the history, symptoms, treatment and result in all the cases of intraperitoneal rupture, sixteen in number, reported up to that date. I take the liberty of presenting a new table made up in part of MacCormac's, with fuller de-

No.	Operator and Reference.	Age.	Time after Accident.	Symptoms.	Lesion in the Peritoneum.	Treatment.	Result.	REMARKS.
1	Wm. T. Bull, <i>Annals of Surg.</i> , Vol. 1, No. 1.	46	13 hours.	Characteristic.....	3¾ inch rent in posterior wall.	8 Lembert sutures; catheter tied in bladder; no drainage.	Died in 7 hours.	Fracture of pelvis; careful antisepsis.
2	Jos. M. Fox, <i>Phil. Med. News</i> , Dec. 10, 1887.	38	19 hours.	Characteristic.....	2½ inch triangular rent.	15 Lembert catgut sutures; catheter tied in; no drainage.	Died in 42 hours.	Post-mortem showed sutures intact; careful antisepsis.
3	C. Heath, <i>Med. Chir. Trans.</i> , No. lxii.	Adult	40 hours.	Tense belly, bloody urine by catheter.	Intraperitoneal rent (not described).	Neither catheter or drainage are mentioned; continuous suture of catgut; not further described.	Died on 6th day.	Post mortem showed sutures had given way (probably on 3d day); treatment probably not fully antiseptic.
4	Sir W. MacCormac, <i>London Lancet</i> , Dec. 11, 1886.	33	19 hours.	No shock; 95 oz. of blood by catheter; great pain.	4 inch rent in bladder, far back, not easy of access.	16 Lembert silk sutures, with a few sutures superficial of catgut; catheter every 4 hours; drainage-tube in abdomen.	Recovered.	There was no peritonitis, and patient recovered without a drawback.
5	Sir W. MacCormac, <i>op. cit.</i>	37	27 hours.	Neither shock nor symptoms; fluid in abdomen.	3 inch rent.....	12 Lembert silk sutures.	Recovered.	Neither catheter nor drainage. No peritonitis.
6	A. F. McGill, <i>Lancet</i> , 1886, xxi, 972.	54	68 hours.	Insensible for a time. Peritonitis.	4 inch rent from apex to fundus.	9 chromic catgut sutures.	Died in 17 hours.	Catheter nor drainage mentioned; post-mort. showed bladder wound firm; no fluid in peritoneum.
7	C. J. Symonds, unpublished.	47	7	Collapse, great pain, adhesions over hypogastrium, blood by catheter.	Y-shaped rent, partly intra- and partly extra-peritoneal.	12 Lembert sutures.....	Died on 7th day.	Drain nor catheter mentioned; post-mortem, bladder leaked; pelvic fracture; secondary laparotomy for peritonitis.
8	T. P. Teale, <i>Lancet</i> , June 24, 1887.	25	18 or 20 hours.	Pain, no urine by catheter, dullness in flanks.	1 inch rent in fundus.	Perineal section and drainage; laparotomy; 6 catgut sero-serous sutures.	Died in 19 hours.	Post mortem, bladder water tight, no peritonitis.
9	Mr. Halstrom, <i>Lancet</i> , June 28, '88.	21	13 hours.	No shock.....	Rent 1½ inches.....	Catheter 1st day: 9 Lembert sutures.	Recovered.	No peritonitis, no drainage. Irrigation practiced with boric acid solution.
10	A. Willett, <i>St. Bartholomew Hosp. Report</i> , 1876, p. 209.	48	29 hours.	Shock, pain, bloody urine by catheter. Peritonitis.	Transverse rent 3½ inches.	Interrupted sutures. (Silk?)	Died in 23 hours.	Leakage of bladder wound. No mention of drain or catheter.
11	T. Holmes, <i>Lancet</i> , July 23, 1887.	24	6 hours.	Pain; blood-stained urine on deep catheterization; point of catheter felt in abdominal cavity.	Rent 2 inches.....	8 sero-serous sutures of carbolized silk; perineal section uncompleted.	Recovered.	No abdominal drain; carbolic antisepsis; no peritonitis.
12	E. L. Keyes, <i>N. Y. Med. Record</i> , Dec. 24, 1887.	22	22½ hrs.	Incipient peritonitis; a little bloody urine by catheter; characteristic symptoms.	Rent 1½ inches backward.	Antiseptic care; 9 Lembert sutures of silk; permanent catheter; glass drainage-tube.	Died in 18 hours.	Post-mort., bladder wound intact; death from shock and alcoholic depression.
13	H. O. Hitchcock, <i>Pittsburg Med. Review</i> , March, 1888.	34	15 or 18 hours.	Shock; free fluid in cavity of abdomen; could not urinate; 11 pints by catheter of bloody urine from abdominal cavity; no peritonitis.	Rent 2 inches; peritoneal apposition of edges not entirely accomplished.	Carbolized catgut suture, not defined as to introduction; antiseptic irrigation, but not complete; permanent catheter.	Died on 2d day.	No drainage; vague report of post-mortem.
14	H. H. Grant, unpublished.	19	5 hours.	Shock; bloody urine by catheter; empty bladder later; slight bulging in perineum but no urine on aspiration.	Rent 2½ inches. Fracture of horizontal ramus near symphysis.	11 carbolized silk sutures, Lembert, complete antisepsis, catheterization every two hours; drainage-tube of rubber.	Recovered.	Drainage-tube removed on 6th day; no peritonitis, no result from fracture.

⁸ *Medico-Chirurg. Trans.*, vol. lxii.

⁹ *Treatment of Wounds*, p. 370.

¹⁰ *Op. cit.*

¹¹ *Brit. Med. Jour.*, May 14, 1887.

tails supplied to some of his cases, published since his report of them, and adding four cases treated subsequent to his paper. I have removed from comparison cases Nos. 2, 5, 9, 10, 11 and 15 of his table. In Nos. 2, 11 and 15 (the latter Walter's case at Pittsburg) no sutures were used in the bladder, and in Nos. 5, 9 and 10 the injury was extraperitoneal before abdominal section. To the remaining ten cases, seven of which were fatal, I add 4—one by Mr. Holmes,¹² recovering; one by Dr. E. L. Keyes,¹³ died in eighteen hours; one by H. O. Hitchcock,¹⁴ died on second day; and one by the writer, recovering.

The history of this case is briefly as follows:

Case.—Jesse Minor, white, æt. 19 years, weight 135 lbs., fell from the shaft of a light cart (weight 400 lbs.), the wheel passing over pelvis and hypogastrium—from right pelvic spine—leaving only a slight bruised mark on the skin. Accident at noon April 2, 1888. After lying still about fifteen minutes the boy stood up without assistance, tried to pass water, but failed, though his bladder had not been emptied since 6 A.M. Prof. S. E. Woody saw him in about twenty minutes after the accident, elicited the above history, passed a catheter and drew off about 2 ozs. of bloody urine. A hypodermic needle withdrew no urine from the perineum. There was *intense* burning pain in hypogastrium and severe shock. Dr. Woody recognized the existence of intraperitoneal rupture. I saw the patient with Dr. W. at 1:30. A catheter introduced at this hour—about sixty minutes after previous catheterization—withdrew only a few drops of urine, still bloody. The pain, though somewhat controlled by the large doses of morphia given by Dr. Woody, was still severe on motion; pulse good, but countenance indicated shock. Fluctuation was not made out in abdomen, though some dullness existed on percussion over the flanks. The ensemble of the symptoms indicating explorative section of the abdomen, the boy was removed to his home, a mile away, the family advised of his condition, and preparation made at once for laparotomy.

Operation.—At 5 P.M., after examination under chloroform, with the concurrence and assistance of Dr. A. M. Cartledge, the operation was made, five hours after injury. Dr. S. E. Woody administered chloroform, and Mr. Woody, of the class of the Kentucky School of Medicine, assisted us. The pubes were shaved and washed quickly with warm soap and water, and then with a sublimate solution, 1:1,000; 3 per cent. carbolic solution for instruments and thorough cleansing of our hands and arms to elbows; towels wrung out in sublimate solution 1:1,000 were put about the site of operation, and a gallon of hot Thiersch solution prepared to irrigate the cavity. An incision in the median line from the pubes 3 inches

in length exposed the peritoneum, bruised and discolored, but unruptured save in one or two minute spots, through which the bloody water from the cavity oozed out. The cavity was quickly opened and $\frac{1}{2}$ gallon of bloody urine poured out. The incision was now extended to the umbilicus. The introduction of the finger in front of the bladder discovered a loose fragment of bone about the size of a large chestnut, evidently crushed off the horizontal ramus of the pubes. This was not disturbed. Dr. Cartledge now passed a sound into the bladder, and I could presently find its point emerging through a rent in the fundus, $2\frac{1}{2}$ inches in a transverse direction. The cavity of the bladder could be easily explored by the finger through this rent.

The abdominal cavity was washed out with Thiersch solution and, while the bladder was held up by a stout catgut ligature passed through the peritoneal coat behind the rent, I easily introduced eleven sutures of carbolized silk—rather coarser than would have been preferred—after Lembert's method, beginning a little beyond the end of the rent and drawing together the peritoneal investment of the bladder beyond both extremes of the tear. This part of the operation, done with an ordinary curve-pointed needle without a holder, was accomplished with unexpected facility. The bladder was not tested by injection for want of a proper syringe. The cavity of the peritoneum was again carefully irrigated with hot Thiersch solution, and sponged dry with mops of absorbent cotton. The wound was closed with deep sutures of silk passed through the skin, muscles and peritoneum. Superficial sutures of catgut closed the wound around a drainage-tube of soft rubber, introduced by Dr. Cartledge behind the bladder. Iodoform was dusted over the line of sutures, and bichloride gauze covered with absorbent cotton and bandage completed the dressing. The time occupied in the operation was about one hour.

The patient reacted promptly from the chloroform, and spoke intelligently in a few minutes. A hypodermic injection of $\frac{1}{3}$ -gr. morphia was administered, and he was left in the care of Mr. Woody, with instruction to introduce the catheter every two hours. He spent a fair night, with almost no pain. About 14 ozs. of urine were drawn in eighteen hours. At 8 A.M. his pulse was 115, temperature 99°. He was carefully watched and fed, and the catheter used every two to four hours as time went on, Mr. Woody and Mr. R. C. Telly, my private student, also of the class of the Kentucky School of Medicine, remaining with him alternately night and day. His temperature went to 100° but once—on the ninth day, from constipation—returned to normal after a free evacuation in response to sulphate of magnesia.

The drainage-tube was aspirated with a syringe daily for five days, never yielding over 1 drachm of sweet serum. It was removed on the sixth day.

¹² Op. cit.

¹³ N. Y. Med. Record, December 24, 1887.

¹⁴ Pittsburg Medical Review, March, 1888.

The wound united throughout, except around the tube, by first intention. The patient sat up on the fourteenth day; catheterization was discontinued on tenth day. The fracture of the pelvic ramus gave no trouble, and has occasioned no symptoms. I present the patient to speak for his recovery as complete.

In comparing the various methods of detail in the operation I shall return presently to one or two features in the steps taken in this case. The briefest report of the recorded operations I can make is in a reproduction of the table I have prepared. Fourteen cases are recorded in which the complete steps of laparotomy, intraperitoneal suture of the bladder, and closure of the abdominal wound were practiced. In some the catheter was tied in the bladder, in others repeated catheterization was practiced, and in one or two no steps were taken to drain the bladder. Abdominal drainage was practiced in the majority of cases. Of the total five recovered and nine died.

As a duty of the very highest practical importance bears upon us the obligation of determining, out of a comparison of results, which treatment promises most. Three methods are before us. The plan of inaction, with merely drainage by the catheter, recommended as safest by Heath,¹⁵ after failing in his operation, must, in the light of present knowledge, be condemned as unsurgical and not to be entertained. We will consider laparotomy without suture; laparotomy with suture to external wound; laparotomy with closure of bladder rent.

It will be noticed from the table that no case of recovery presented any peritonitis, which of itself speaks volumes for the safety of the operation under antisepsis.

Notwithstanding the success of Walter's case, treated by the first method, there was severe peritonitis, besides the other reported cases, one by J. Duncan¹⁶ and one by Sonnenberg¹⁷ died on the 3rd and 4th days respectively of intense peritonitis. Moreover, most frequently the rent is so large, or occupies such a site in the bladder, as to render the prevention of seepage of urine into the peritoneal cavity impossible by any form of drainage. Besides, it is difficult to see any advantage to accrue from leaving the wound unsutured after the belly is sewed up as in Walter's case. The presence of carbolyzed silk sutures introduced by the method of Lembert, if not absolutely harmless, are a thousand times safer than extravasation of urine, and, if they occasionally, or even in two-thirds of all cases, fail to accomplish the object of their introduction the condition is no worse than those before. Nay, it is much better, for every hour before the extravasation nature is throwing a breast-work around the parts she desires to protect, and if the stitches do yield on the 3d or 4th

day the damage will be certainly less disastrous.

With reference to the second method referred to favorably by later writers, notably Wyeth,¹⁸ as appropriate for intraperitoneal rupture as well as after extraperitoneal supra-pubic section, a diversity of opinion is likely to prevail. The conditions after supra-pubic lithotomy are more favorable to convenient and successful suture of the bladder than in cases of rupture; yet the results are certainly less favorable in cases so treated. In his "Observations on Supra-pubic Lithotomy"¹⁹ last year, Sir Wm. MacCormac says: "Schmitz has collected fifty-seven cases of suture of the bladder; forty-seven recovered, and eight died. . . . The bladder wound healed by first intention in seventeen cases. The permanent catheter was used twenty-three times; eight times the wound united by first intention. In thirteen cases the catheter was not used; of these only three united by first intention. Twenty-three time catgut was employed, with only four successful results. In ten of the seventeen, where union took place by first intention, the mucous membrane was not included in sutures; in five no statement is made whether or not; in two, it was included; when the sutures gave way, this happened generally on the 4th day. By this time the danger of infiltration is mainly over."

These facts indicate that about one-third of cases of suture hold permanently, in the remaining cases the wound re-opens. Though, in intraperitoneal rupture the conditions are not quite so favorable, and though it is particularly true that the lapse of two or three days is not so much a safeguard against infiltration as under the conditions referred to by Schmitz, yet his conclusions are highly pertinent to the subject. In the same paper MacCormac gives statistics of the suprapubic operation without suture, with mortality of 24.4 per cent.; with suture the mortality is 35 per cent.

In view of these figures, which represent facts, it is perhaps wise to select a medium course. Under such conditions, as considerable period since the injury, with unfavorable local and constitutional indications with irregular lines in the rent, it would be preferable to stitch the tear to the abdominal wound, fill the incision with iodoform gauze, and drain both by catheter and tube. Under most ordinary conditions, and especially when the rent is too far back to admit of reaching the abdominal incision, the third method is appropriate. In a private communication, Prof. J.A. Wyeth mentions a case under his observation in which the bladder rent was closed, and then, for safety, the bladder attached to the abdominal wound. The cut, occurring during ovariectomy, did re-open, but infiltration was prevented by the patient recovering.

¹⁵ Op. Cit.

¹⁶ Lancet, 1886, vol. ii.

¹⁷ Centralblatt für Chirurg. 1885.

¹⁸ Text Book on Surgery, 1888. Page 566.

¹⁹ British Medical Journal, March 12, 1887.

Diagnosis. All experienced observers on abdominal surgery are in accord as to the great advantages of an early operation. The time *par excellence* is in the first eight or ten hours. Shock after such an injury does not subside, but deepens with the growing gravity of the condition. Though case No. 7, of MacCormac's table, operated on twenty-seven hours after accident recovered without peritonitis, yet in Walter's case peritonitis appeared in ten hours, and in most recorded cases indications of inflammation were found at the operation.

Symptoms. Though the symptoms are usually characteristic; clear history, great pain, severe shock, desire to urinate with futile attempts, perhaps little bloody water by catheter, and at times catheter can be felt high up in abdomen—yet occasionally the condition is greatly obscured. In Keyes' case, though the patient was seen in forty minutes after the accident, the nature of the condition was not suspected for fifteen hours. In neither of MacCormac's patients was there shock, and one presented so little distress that he was referred to a dispensary as an out-patient for the first day. Most writers on the subject, refer to the rarity of intraperitoneal rupture. Comparatively, this is true, but I am inclined to believe the infrequency of reports of such condition is due to its being often overlooked. In the *N. Y. Medical Record*, 1888, Dr. M. Singer, of Galveston, reports a case of a man 27 years old, admitted to hospital ten or twelve hours after injury from a kick in the abdomen, catheterization frequently repeated in fifty-four hours preceding death after admission, always found plenty of urine.

Autopsy showed an intraperitoneal rent three inches in length. Attention being drawn to the treatment by recent success, I predict much more frequent reports. Prior to 1886, 3 exploratory laparotomy for rupture, are reported. During 1886, 4 exploratory laparotomy for rupture, are reported. During 1887, 5 exploratory laparotomy for rupture, are reported. Thus far, 1888, 3 exploratory laparotomy for rupture, are reported.

In view of the pretty well established belief that laparotomy of itself, done under careful antiseptic precautions, adds but little to the gravity of any grave condition, it is right to urge that when symptoms strongly suggest rupture of the bladder, and particularly when aspiration of the perineum discloses no infiltration an exploratory operation should be done without delay.

But little is to be added to the conclusions of MacCormac and Holmes as to treatment. Clearly the bladder-wound should be closed after complete cleanly antiseptic irrigation of the abdominal cavity. This irrigation can be most safely done with the Thiersch solution (boracic acid 12 parts, salicylic acid, 2 parts, water 1000 parts), freely and carefully used to remove all contaminating taints. The wound should be approximated with silk

sutures—after Lembert's method. The silk should be carefully carbolized, and of a fairly firm texture, introduced not farther than one-fourth inch apart, cut short and abandoned. Nothing will be lost by washing out the bladder after suture with the Thiersch solution. Both the best of the suture and antiseptic irrigation of the bladder are accomplished in this way. Four ounces is enough to employ. If any leakage appears from such amount of distension, it should be at once looked after. It is true such a step takes a little time, but it is not ill-spent, if it discovers a faulty suture through which a fatal seepage may take place.

The question of abdominal drainage is not decided. It is open to objection on the ground of admission of septic germs, and the weakening of the abdominal walls. The first can be avoided by covering the tube with antiseptic dressings, and carefully, practicing aspiration, and the second by withdrawing the tube in four to six days. By aspiration of the drain all accumulation can be evenly removed, with the risks of absorption of such septic material as may accumulate in the cavity.

Perineal section for the purpose of drainage is condemned, both by MacCormac and Holmes, as not only difficult, upon the empty bladder, but a useless and dangerous complication. When suspicion of urinary infiltration in the prevesical space complicates the diagnosis, aspiration through the perineum, with a long needle, will indicate the condition.

With regard to catheterization, I cannot put the slight risks of the occasional introduction of a soft aseptic catheter, in comparison with the immense advantages of rest of the bladder and prevention of over-accumulation of urine and straining after suture. Though perhaps after the fourth day the patient may be permitted to relieve his bladder in the natural way, even then he should carefully urinate every two or three hours, avoiding all forced contraction of abdominal muscles or of bladder walls.

Refreshing the edges of the rent is wholly unnecessary when the approximation is made by the sero-serous method of suture.

Of antiseptics it is useless to make other mention than that the protection it offers should be secured to every patient by the most skillful and unremitting care.

DR. H. O. MARCY, of Boston: Within the last two years I have had two cases of injury to the bladder which should be put upon record. They were not ruptures from external causes. One was a case in which I found, to my surprise and sorrow, that, while operating upon a large uterine myoma, I had made an incision of from 1 to 2 inches into the bladder. I sutured it together, and union took place. In repairing the rent, I united first the edges of the mucous mem-

brane, and then the overlying structures, with fine tendon sutures. The second case was that of a child of 2 years of age in which there were two openings. I succeeded in closing them, followed by complete cure. All this helps to contribute interest to the subject, and I am sure we are under obligation to Dr. Grant for his demonstration of the subject, for bringing his patient so long a distance to exhibit the result.

GASTROSTOMY; WITH THE REPORT OF A CASE.

BY MILES F. PORTER, M.A., M.D.,
OF FT. WAYNE, IND.

Carl Z., æt. 19 years, German, family history unimportant. Four months before his admission to the Ft. Wayne City Hospital, while yet in Germany, during a drunken spree, he swallowed some "browish-looking fluid" which "burned his throat at the time," since which he has had gradually increasing difficulty in swallowing. He entered the hospital Sept. 9, 1887, when an examination by his physicians, Carl Proegler and Howard McCullough, revealed the existence of an organic stricture of the œsophagus, commencing at the level of the cricoid cartilage, and extending downward about $1\frac{1}{2}$ inches, rendering the passage tortuous and admitting of the passage of no instrument except an elastic urethral bougie; the largest size of this which passed being No. 16, Am.

Repeated efforts to pass olive and acorn ivory-tipped whalebone bougies failed, even with tips of smaller diameter than the elastic bougie passed. Deglutition of liquids and semi-fluids only was possible, but exceedingly slow and difficult. Regurgitation often occurred, and attempts to swallow were accompanied by severe fits of coughing. Notwithstanding the repeated passage of bougies, the stricture grew tighter, deglutition even of fluids became almost impossible, the patient lost strength and weight, and was crying with hunger, when on Dec. 4, 1887, he was advised by his physicians to have the operation of gastrostomy performed. Having gained his consent they requested me to operate, which I did in a room prepared by fumigation with sulphur for 18 hours, thoroughly aired, and supplied with new bed and bedding. I was assisted by Drs. H. McCullough and Carl Proegler, and there were present Drs. Dills and McOscar, a nurse, and the matron of the Hospital.

The patient, having received a bath and clean clothing, was etherized, when his abdomen was washed in a 1:1000 solution of corrosive sublimate, and an incision three inches long was made parallel with and an inch below the costal cartilages of the left side, dividing the muscles—including the outer edge of the rectus—and fascia down to the peritoneum. Forceps were applied to bleeding

points, the peritoneum hooked up and divided on a director the full length of the wound. The greater end of the stomach—which organ was much contracted—together with the lower border of the liver now presented, when the anterior wall of the stomach was drawn out and two threads of black silk were passed parallel and about three-eighths of an inch apart through the peritoneal and muscular coats, by which the organ was held in position by Dr. McCullough while the operation was being completed. Two stitches were passed at each end of the wound, including the parietal peritoneum and all structures of the abdominal wall. Two more were now passed, one above and one below, working toward the centre, including the same structures and also the outer coats of the stomach, after which six sutures were passed on each side of the wound, including the outer coats of the stomach, the parietal peritoneum and skin. All the sutures were now tightened, in the same order as placed, cut short, the wound washed, dried, dusted with iodoform and covered with six layers of sublimated gauze and a generous supply of absorbent cotton secured by a bandage pinned snugly around the body. No ligatures were required. All stitches, 18 in number, together with the guide-threads, were of iron-dyed silk. The guide-threads were of course allowed to remain until the stomach was opened.

The operation occupied less than an hour, and was followed by only slight shock, from which he reacted nicely, the pulse being 76, respiration 26, and temperature 100.5° F. the next morning—18 hours—after the operation.

All went well until the end of 40 hours, when the cough, which had troubled him somewhat ever since his trouble in swallowing commenced, grew worse, respiration became more frequent, the pulse 110, and temperature 102.2° F. An examination revealed a broncho-pneumonia of the lower right side, and showed the wound dry, with no evidence of peritonitis. Up to this time the patient had been taking milk, champagne and brandy *per os*, for, as is usually the case, he had less difficulty in swallowing after the operation; but as efforts at deglutition always excited the cough, enemas were now substituted, and hypodermatics of morphia, grs. $\frac{1}{4}$, given to quiet the cough, which caused pain in the wound. Fomentations were applied to the chest.

The patient gradually grew worse until, at 3 P.M. of the 7th, 48 hours after the operation, the temperature was 103° F., pulse 142 and feeble, respiration 50 with marked cyanosis, and death was looked for by both of my colleagues and myself. We now administered 8 drops of Squibb's fl. ext. of digitalis, hypodermatically, with brandy; ordered fomentations and enemas continued, and 10 drops of digitalis every three hours *per os*, while the morphia was discontinued on account of the cyanosis, though it had produced no pronounced