Annals and Magazine of Natural History

Series 1

ISSN: 0374-5481 (Print) (Online) Journal homepage: <http://www.tandfonline.com/loi/tnah07>

Notice of a black lizard taken in Cheshire

W. Wells Esq.

To cite this article: W. Wells Esq. (1846) Notice of a black lizard taken in Cheshire, Annals and Magazine of Natural History, 17:114, 449-450, DOI: [10.1080/037454809495854](https://doi.org/10.1080/037454809495854)

To link to this article: <http://dx.doi.org/10.1080/037454809495854>

Published online: 16 Dec 2009.

Submit your article to this journal [↗](#)

View related articles [↗](#)

these, and are inspiratory between their costal cartilages. Between the diaphragmatic and intermediate sets of ribs, and between their cartilages they are throughout expiratory. The levatores costarum draw the posterior portion of the lower ribs backwards. In the ass and the dog, the upper fasciculi of the serratus magnus are expiratory, the lower inspiratory, and the intermediate neutral. In man, the greater part of the fasciculi of this muscle is expiratory. In the ass, the lower fibres of the serratus posticus inferior are inspiratory, and the upper fibres expiratory. In the dog and in man, all are throughout expiratory.

MISCELLANEOUS.

Notice of a Black Lizard taken in Cheshire. By W. WELLS, Esq.
To Sir W. Jardine.

Rimrose House, Bootle, Liverpool, April 12th, 1846.

SIR,—When botanizing at Bidston, Cheshire, a few days ago, I met with a lizard a description of which I send you, hoping you will excuse the liberty which I have taken and the imperfections of my remarks, as I am but a tyro in this department of science.

The head, back and throat are a shining black; scales of the belly and tail black, tipped with dark brown. The scales of the head are less numerous than in *Zootoca vivipara*, which it resembles in size and shape except that it is more slender, the tail is longer and more tapering, the head is more bulky, the eyes are perfectly circular and surrounded by a beautiful ring of minute scales, the scales forming the eyebrows do not project over the eye, and those of the back are much smaller than in the common species, which was abundant in the same locality.

DIMENSIONS.

Length of head	$1\frac{5}{8}$ inch.
Body	$1\frac{3}{4}$
Tail	$3\frac{1}{2}$
Total	$5\frac{2}{5}$
Greatest breadth of head	$\frac{1}{2}$
Circumference of body	$1\frac{3}{8}$
Length of longest toe	$\frac{5}{8}$
Shortest	$1\frac{1}{10}$
Scales of the head	20
Joints [scales?] of the tail	62

I am aware that in the 'Mag. Nat. Hist.' vol. i. p. 189, New Series, mention is made of a black lizard by G. W. in a note to his paper on Vipers, but I conceive that the difference between that and the one I have attempted to describe entitles me to consider them different. I have kept a considerable number both of the viviparous and the sand lizard, but none of them evinced the very irritable, untameable disposition of this one; they are in general easily tamed, but the black one seizes everything that is put near it, and never

lets go until forced to do so. I placed a sand lizard within its reach, which it instantly seized, and retained its hold for upwards of two hours, although for more than half that time it was suspended from it.

Before concluding, permit me to mention that the following reptiles have been found by me near Liverpool since August last: *Zootoca vivipara*, *Lacerta agilis*, *Natrix torquata*, *Vipera communis*, *Triton palustris*, *T. aquaticus*, *T. vulgaris*, *Rana temporaria*, *Bufo vulgaris*.

I remain, yours very respectfully,

W. WELLS.

New species of Fossil Bats. By M. H. DE MEYER.

Cheiroptera have but rarely been found fossil in the tertiary formations. Of this family, in the deposits anterior to the diluvian epoch, only the *Vespertilio parisiensis* of the schists of Montmartre was known, a single individual of which is preserved in the Museum of Paris; and two small teeth found in the eocene sand of Kyson, and referred with doubt by Prof. Owen to the genus *Vespertilio*. The species indicated by Karg, at Oeningen, appears to be very uncertain, and the original specimen has not been discovered again. Hermann de Meyer has detected at Weisenau, amongst a considerable mass of fragments of bones, some belonging to two *Cheiroptera*. They consist of one half of the right lower jaw, in which, although the teeth are wanting, the alveoli are sufficiently well preserved to give an idea of the dental system; of three humeri, of which two are left and one right, which show the existence of two species, and prove even probably a generic difference between them; lastly, of one half of a radius which likewise can only belong to a *Cheiroptera*. These two species differ from that of Montmartre, and H. de Meyer has designated them under the name of *Vespertilio præcox* and *V. insignis*, until their generic affinities are definitely fixed.—*Leonhard und Bronn's Jahrbuch*, 1845, p. 798.

Does Magnetism influence the Circulation in Chara?

By M. DUTROCHET.

In some experiments made in the year 1837, M. Dutrochet proved that electric currents only affect the circulation of *Chara* by their action as *exciting* causes. The movement of the fluid is only arrested for a few minutes and then recommences, often with greater rapidity than before, as the equilibrium between the electric and vital forces is restored. This action is exactly analogous to that of hot water or solution of common salt.

M. Dutrochet has recently submitted *Chara* to the influence of a large electro-magnet, capable of supporting a weight of about 4000 pounds. The stem of *Chara* was placed a little in front of a plane passing vertically through the poles of the horse-shoe magnet, but quite within the magnetic influence. Careful observation at the moment of establishing the electric current in the coil, proved that the speed of the circulation was unaltered. Left thus for ten minutes, all remained as before—no influence was manifested. The