not receive official recognition. Possibly an exception should be made in favor of a few old favorites, such as the compound cathartic pill, the Dover's powder, the Brown mixture of ammonium chlorid, etc.

Every definite, simple substance, whether inorganic or organic, which has been shown to be of service as a drug or as an office accessory, should be officinal. If our present patent and trade-mark laws cause a conflict with ethics, the Pharmacopeia should, nevertheless, recognize the drug, and the professions of medicine and pharmacy should then unite to secure a modification of the objectionable laws. In the fight against disease and death, no physician can hesitate to see whether his weapons are labeled "ethical," but must seize that which will be the best protection. If we remember rightly, it has been legally decided that neither the scientific nor the common name of an article can be reserved as a trade-mark. The monopoly of a preparation can be secured in one of two ways, either by patenting a method of preparation, which must be fully described and shown to be distinctly novel, or by the use of a trade-mark and keeping the method of preparation secret. The general processes of a science can not be patented, so that the preparation of a drug is open to anyone who can devise a new method, while a secret nostrum, covered by a trade-mark, is at the mercy of an expert analyst. As a matter of fact, almost all the monopolies of the latter class are compounds of no originality and little value.

The Pharmacopeia should adopt as its standard of strength and quality, the highest attainable, even if the competition of retail pharmacists is thereby excluded and the manufacturer is confined to wholesalers with elaborate and expensive equipments. The petrolata, as standardized by the last edition of the Pharmacopeia, are by no means equal to the clear products of several firms. The homeopathic mother tinctures are said to be, in general, far superior to those of the regular profession. Pill coatings of the crudest kind are still officinal. The manipulations of chocolate, gelatin, keratin and salol coatings, are quite unofficial, nor are the cachets, the gelatin bougies for the urethra, rectum, etc., and many other refinements of pharmacal art, recognized by the authority that ought to stand for the best attainable in pharmacy.

It is still too early to discuss in detail the proper regulation of animal extracts, serums, etc., and the improvements and modifications of opinion which will follow within the next few years would render the wisdom of 1900 crude and imperfect long before the expiration of the next decade. The most that can be expected of the next revision is the formulation of some rules of nomenclature and a general recognition of opotherapy.

At the last revision of the Pharmacopeia, the repudiation of the metric system was greatly feared. At the next revision, there seems to be no probability of the reopening of this issue. So far as the body of the Pharmacopeia is concerned, we believe there is no need of mentioning any unit of measure or weight. As we have already pointed out, the separation of any part of an officinal substance into a pill or other dose, is to be deprecated, hence the Pharmacopeia need only designate the proportional parts in ordering the preparation of galenical. We believe that no one will question the advisability of using convenient percentages, whenever possible, and if a very slight change in relative proportions and active strength is allowed, the percentages may always be stated in convenient numbers. After the pharmacopeial preparations are made, the physician

should be free to use his judgment as to whether he will prescribe in grams, grains or pennyweights, or any other unit on which he may agree with his local druggist. Nevertheless, for the country at large, only the metric and the apothecaries' systems can be officialized.

A PLEA FOR THE AGED*.

BY J. W. BELL, M.D.

PROFESSOR OF PHYSICAL DIAGNOSIS AND CLINICAL MEDICINE,
UNIVERSITY OF MINNESOTA.

MINNEAPOLIS, MINN.

One of the saddest commentaries on the medicine of the past can truthfully be said to be the neglect of the aged, as indicated by the scanty literature on the subject and the general apathy existing in the profession relative to the study of the normal changes as well as the diseases peculiar to advanced life. To the profession of France largely belongs the honor of laying the foundation for whatever we possess, fragmentary though it be, of senile pathology.

Prior to the beginning of this century those who wrote on the subject wrote, like Cicero, from a contemplative literary standpoint with the possible exception of the small treatises of Floyer, Welsted and Fisher. It remained for Pinel, in his elaborate treatise on "Clinical Medicine," published in 1815, the result of his vast experience in the Salpêtrière Hospital, to furnish the nucleus of our present knowledge of senile pathology. In 1839 Constatt published his treatise on the diseases of old age. In this work we behold the first systematic effort to bring together the scattered fragments of senile pathology. Charcot, in referring to this work, says: "In it, imagination occupies an enormous amount of room at the expense of positive and impartial observation." Cruveilhier, a few years later, from material gathered during his stay in the Salpêtrière, shed additional light on the pathology of old age in his "Atlas of Pathological Anatomy."

The works of Rastan and Charcot, both eminent clinicians, having access to the wards of Salpêtrière, have contributed much to our knowledge of the diseases of advanced life. To Charcot, especially, are we indebted for a small but valuable treatise. Geist, Day, McLachlan, Cruveilhier, Beau, Humphrey, Fothergill and Loomis are names worthy of mention as able contributors to this much-neglected subject, and be it remembered that "The Twentieth Century Practice of Medicine" was the first of our many excellent works on practice to add *one* chapter on old age and its diseases. The want of interest, as indicated by the scanty and fragmentary character of the literature on the subject, is largely responsible for the apathy existing to-day in our medical schools. The average medical student passes the college curriculum hearing scarcely a reference made to the changes peculiar to age and having little opportunity to become familiar with the maladies prone to overtake the frame as the years speed onward. He scarcely recalls a reference by one of his teachers to old age, unless suggested in mitigation of the failure of some brilliantly planned but misguided operation or equally ill-timed drug treatment.

Surely if the pathology of childhood demands clinical consideration of a special kind such as it now receives in our medical schools, and it be indispensable that it be known from a practical point of view—and this the

*Presented to the Section on Practice of Medicine, at the Fifth Annual Meeting of the American Medical Association held Columbus, Ohio, June 6-9, 1899.

writer does not question—then, is it not equally true that senile pathology with its associated clinical manifestations calls for special consideration at the hands of medical teachers?

In order to make more clear the necessity for this, let us briefly recall some of the more pronounced changes due to age. From birth onward the increase of growth, though undergoing certain variations, gradually diminishes. Commencing at about 40 years of age, the prime of life declines into old age. The metabolic activity of the body, at first sufficient not only to cover the daily waste, but to add new material, later on is able only to meet the daily wants, and, at last, is too imperfect even to sustain the existing frame. As a result there must follow certain changes in the tissues of the aged, of an atrophic and degenerate character. The atrophic changes involve, especially, the muscles and glands. The cellular elements gradually diminish in size without apparent modification in their structure. The muscular fibers become smaller and more uniform in size; the spleen and lymphatic glands undergo a marked diminution in volume and weight, which increases with advancing years.

In the digestive tract the glandular parts and villi are shrunken. In many instances, however, the glandular elements seem to be those whose powers are longest preserved; and hence the man who, in the prime of life was a martyr to dyspepsia, by virtue of hypersensitive gastric nerves finds in his declining years that his nerves are blunted and his peptic cells are permitted to pursue their chemical labor undisturbed by extrinsic nerve storms, that he can eat and drink with the courage and comfort of a boy.

The eye in its prime in childhood, when its media are clearest and its muscular mechanism most mobile, later in life, when it would be of the greatest service to a still active brain, has reached a clouded and rigid old age.

Senile degenerative changes begin somewhat later than the atrophic. The tissues undergo fatty and calcareous changes. These changes are marked in the blood-vessels, especially in the cerebral arterioles, also in the voluntary and involuntary muscular fibers. Calcareous deposits occur especially in the tendons, ligaments and cartilages. In the central nervous system, the neuroglia tends to predominate over the nervous elements proper—the brain tissue undergoes a chemical change—the fatty constituents are diminished, while the amount of water and phosphorus is increased. As a result, the sensory and motor impulses pass with increasing slowness to and from the central nervous system, and the brain becomes a more and more rigid mass of nervous substance, the molecular lines of which rather mark the history of past actions than serve as indications of present potency.

Everywhere we observe a tendency on the part of the living tissue to fall back on the easier task of forming fat rather than to carry on the more difficult task of manufacturing new material like itself. In no part of the body is this more evident than in the arteries. One of the saddest changes of advancing years, because the most fatal, is the conversion of the supple, elastic, arterial tubes into rigid atheromatous channels, whereby the supply to the various tissues of nutritive material is rendered more and more difficult, and decay hastened. In consideration of a few general changes outlined as the result of age, it is evident that they must very materially modify the manifestations of disease in the aged, as well as its therapy.

For the purpose of illustration and contrast, permit me to briefly recall to your minds some of the more important pathologic, clinical and therapeutic distinctions between senile and adult pneumonia. To study intelligently diseases of the respiratory organs in advanced life it is necessary to bear in mind some of the more important anatomic and physiologic changes resulting from age. The thorax loses much of its mobility, due to ossification of the cartilages, and atrophy of the muscles. The tracheal rings harden; the bronchial mucosa and muscularis atrophy. The lung tissue becomes attenuated, less vascular and elastic. The air-cells become dilated, the capillary vessels in part obliterated. In consequence of these changes the breathing of old persons becomes embarrassed and labored, thus impeding their locomotion and enforcing sedentary life.

The pathologic changes in senile pneumonia involve the apices by preference in the aged, the base in adults. The inflammatory changes are more often central, and the first and second stages shorter in duration, in the aged, than in adults. The terminal bronchi are invariably involved by the inflammatory process; the pulmonary pleura seldom; the reverse being the rule in adults. The hepatized lung is lighter and more friable in the aged.

Clinically, senile pneumonia differs essentially from the pneumonia of adults. It commences insidiously, and usually runs a latent course; especially is the latter true of intercurrent attacks. The disease is usually initiated by a slight chill, as compared with the sharp chill or rigor of adult age. Old persons seldom complain of pleuritic pain, or dyspnea, the reverse being the rule with adults. The systemic disturbance, as evidenced by pulse, temperature and respiration, is never so pronounced as in the adult. Expectoration is more tardy, difficult, scanty and less colored than in adults. The disease being asthenic in the aged, as a rule, is the reverse in adults. The physical signs are modified by the changes in the aged thorax. The percussion note is often one of exaggerated resonance, which slowly gives way to dulness over the affected area.

The auscultatory signs are less positive in advanced life, crepitation being of brief duration or absent, subcrepitant and mucous râles predominating during the first stage. Bronchial breathing, more intense than in adult life, marks the second stage, accompanied by mucous râles, the latter continuing into the third stage. Surely a disease differing so essentially in its pathologic and clinical aspects from that of adult life must require special treatment.

Let us take a second example, a very common affection of the aged, one of special interest to our surgical brethren—hepatic colic. As practitioners we are all familiar with the formidable aspect of hepatic colic in the adult. How different in the aged—slight pain or weight in the right hypochondrium, slight jaundice, perchance nausea, seldom amounting to emesis, accompanied by slight systemic disturbance make up the clinical picture in the aged.

The preceding brief and imperfect sketch tells us that the progress of age establishes a wide difference in pathologic and clinical phenomena by virtue of its physiologic modifications. The writer believes that the time is long past due when there should be an awakening in our medical schools to the importance of this subject; and an honest effort made to furnish systematic instruction, both didactic and practical: 1, in the normal changes, the result of advanced life; 2, in the diseases peculiar to old age.

It would seem criminal to even suggest the addition of another distinct course to the already overcrowded college curriculum, but I would suggest the following method of expansion:

1. That the chairs of anatomy and physiology impart to the student the necessary primary instruction relative to the normal changes peculiar to old age.

2. That the chair of practice, or if deemed best, in order to contrast disease, the chair of pediatrics, enlarge its scope and furnish the necessary pathologic and clinical instruction so essential to fully equip the student for the responsible duty of intelligently advising and treating the aged.

SPUTUM ANALYSIS AND THE EARLY DIAGNOSIS OF PULMONARY TUBERCULOSIS.*

BY S. EDWIN SOLLY, M.D.

COLORADO SPRINGS, COLO.

Seven years ago I called the attention of the profession to the "Neglect of the Early Diagnosis and Treatment of Pulmonary Tuberculosis," in a communication read before the Arapahoe County Medical Association. At that time I used the following words:

"The great importance of the early diagnosis of pulmonary tuberculosis has been forced on my attention by finding so many tuberculous patients seeking the climate of Colorado long after their chances of recovery in that or any other climate have gone by. Considering it a matter largely preventable, by the profession realizing more clearly the causes that lead to this unnecessary and lamentable loss of life, I have analyzed the last 100 cases of pulmonary tuberculosis that have presented themselves for examination. This inquiry has nothing to do with the influence of climate nor the results of treatment, but was undertaken to find out how far the early recognition of the disease and its treatment affected the condition in which the patient first came under my observation.

"After carefully going over the notes of each case, I have made two classes. The first includes those in whom the disease was diagnosed and treated as soon as the symptoms given in the patient's history showed the disease sufficiently developed to be recognized by any ordinarily well-educated practitioner. By treatment, as applied to this class, is meant any means, hygienic, therapeutic or climatic, that was seriously used—not that the treatment was necessarily of the best, but that the cases were taken in hand and the patient advised in a reasonable manner. The broad question of treatment or no treatment of the pulmonary tuberculosis is dealt with, and not its details nor wisdom.

"The second class includes those in whom the history exhibits the fact that the disease was not diagnosed nor treated for a period varying from a few months to several years after the time that the real nature of the malady had clearly revealed itself.

"The 100 cases are not in any way selected, but are taken in the order of their examination, tracing back from the recent time when this inquiry was commenced, and were limited to this number simply on account of time and labor."

I have recently conducted a similar inquiry analyzing my notes of the first examinations of the last 100 cases of pulmonary tuberculosis presenting themselves for examination. I am glad to say that a comparison of the

statistics shows an improvement in the latest investigation. In 1892 I found that diagnosis of the disease was delayed in 52 per cent., and the total average of delay was two years. In the present year I find that the delay was 45 per cent., and the average period of delay nine and a half months. In 1892 sputum examinations had been made in only 42 per cent., some of which were not made until some time after the disease was well advanced, so it may be said that in 65 per cent. this means of diagnosis was neglected. In 1899, 55 per cent. were examined, so that my figures show that in the intervening seven years there is an improvement in diagnosis going on; but the conditions are by no means as yet ideal. For, omitting the question of the use of the more delicate methods of early diagnosis, such as can be made only by experts, there are 45 per cent. of cases which should have been diagnosed by an average practitioner and in which the diagnosis was delayed for nearly a year. In the 55 cases in which there was no delay, 36 arrived in Colorado in the first stage, 7 in the second, and 3 in the third stage. That is, there was 65.5 per cent. in the first stage in those in whom there was no delay. In those in whom there was delay 8 arrived in the first stage, 20 in the second, and 17 in the third; thus but 18 per cent. were in the first stage. An analysis of several thousand treated in high climates shows that the average percentage of cures in the first stage is 65 per cent., and in the second and third stages combined only 15 per cent. It therefore means that 37 per cent. of the phthisical patients coming to Colorado are, in consequence of the delay in diagnosis, only given a 15-per-cent. chance of recovery, as against 65.5 per cent. to which they were entitled. In fact, granting the truth of these statements, 37 out of every 100 consumptives are slaughtered. This large, unnecessary destruction of lives is criminal, yet the authors of this are not themselves deliberate criminals, although guilty of criminal neglect. How can we explain this strange negligence? By investigating the causes and by calling attention to the lamentable facts we shall tend to remove this opprobrium from the profession. Often the delay in the treatment of tuberculosis is caused by the patient's own delay in placing himself in the hands of a physician. Such cases are, however, eliminated from this inquiry.

When a patient appears to be declining in health, without obvious cause, his morning and afternoon temperature should be taken, and if this can not conveniently be done by the physician, at those times, the thermometer can be left with the patient or his friends, with the necessary instructions. Again, the weight is not taken often enough or the pulse-rate noted. In many there is little or no cough or expectoration, or it is often denied or ignored by the patient, whereas some sputum can generally be collected and should by all means be examined. Blood spitting is frequently passed over as of no importance, and ascribed to the throat or nose when it really comes from the chest. If the patient can be examined at the time of the bleeding usually a few tell-tale râles are to be heard, and bloody sputum shows bacilli then and not at other times. A diagnosis of early tuberculosis can rarely be made by auscultation through the clothing. Sweating at night, unless considerable, is often not mentioned by the invalid. Shortness of breath is frequently ignored because the patient does not show it or speak of it when being examined. Both patients and doctors are for the most part too pessimistic in regard to their views of the curability of tuberculosis, and are still living, as it were, in the days when a patient was not considered tuberculous until he had become

*Presented to the Section on Practice of Medicine, at the Fiftyeth Annual Meeting of the American Medical Association, held at Columbus, Ohio, June 6-9, 1899.