A.—All answer this in the negative. This conclusively shows that catheterization is practically harmless in a healthy individual and so may be done for diagnostic reasons or for demonstration and teaching.

Q. 3.—One side being infected, how frequent is infection of healthy side produced by catheterization?

A.—Ten never knew it to occur. One: "Can't answer." One: "Never if bladder is healthy." One: "Most infrequent, no personal experience of occurrence."

This is certainly overwhelming evidence of its usefulness and safety.

Q. 4.—Knowing one side to be infected, can one rely on catheterization of this side alone, leaving the other side to be represented by a catheter in the bladder?

A.—Seven answer: "No." Three: "If bladder is not involved." One: "In some cases." One: "If urine from catheter in bladder is normal." One: "If catheter is inserted high in ureter and if bladder is healthy."

From these answers one must conclude that it is necessary to catheterize both sides in the majority of cases. Particularly since we know the bladder or its contents to be infected in most cases.

Q. 5.—After nephrectomy without resection of the ureter or injection of same toward the bladder with pure carbolic acid, is it not common to have pus in the urine coming from ureter so left?

A.—Six say: "Yes." One: "Yes if pyelitis existed." Two: "Yes in tuberculous cases." One: "Majority for short time, persists in few." One: "Yes for few months." One: "No." One: "No. Observed, however, in two cases pyonephrosis."

This being answered by almost all in the affirmative, it seems best surgery to resect the ureter and inject any portion remaining with carbolic.

Q. 6.—If infection of healthy side does occur should one expect marked symptoms and should these follow soon after the instrumentation?

A.—Nine: "No such experience." Four: "If it occurs, marked symptoms soon."

These answers imply that such an accident must be very rare and that should it occur marked and rapid symptoms should be expected.

These six questions cover, as I intended they should, every possible point on which a question of justification for this procedure can be raised, and I am most grateful for the responses received.

In order to have experts available in this field it is essential that all work in the smaller cities be turned over to one or two men who are willing to give it their earnest attention. No man who has not had good surgical training should attempt this work.

Having established the usefulness of this procedure and almost conclusively proven its safety when done by one trained for the work, we are ready to look into the bladder and see what can be learned from the use of the ureteral catheter.

We will spend no time in determining or discussing which medium, water or air, or direct or indirect view, is best. It matters not what instrument is used so long as the operator who uses it can get results. For a long time I used air only, but of late have found many instances in which I prefer to use water to distend the bladder.

An examination of the bladder is in a way unavoidable while catheterizing the ureters. The condition of the mucous membrane, the presence of cystitis, ulcer, polyp, tumor or stone, the existence of diverticuli, or sinuses connecting with another viscus may be recognized. The jet of urine as it spurts from the ureteral opening without the entrance of the catheter often makes clear a vague pathologic kidney condition. I have twice seen the expulsion of a calculus through the

dilated ureteral opening. A tuberculous kidney can often be diagnosed by the appearance of the mucous membrane about the ureteral opening in the bladder. Recently I saw a picture on looking into the bladder that showed such an amount of venous stasis from pressure of a large renal growth of the left side that the veins honeycombed the surface of the mucosa so extensively that the bladder seemed trabeculated. On dilating still further the pressure from within entirely overcame the condition. Stricture of the ureter at its opening or along its tract gives symptoms long before any marked destruction of the kidney occurs, and when relieved naturally avoids the loss of the kidney and often the life of the patient. Obstruction of the ureter by calculus, kink or pressure from extrinsic tumors are diagnosed. Sacculations of the ureter with pus pockets are made out by the flow of pus in good quantity before the catheter has reached the pelvis with a cessation of flow on passing beyond the pocket. Pyelitis, pyonephrosis, hydronephrosis, pyonephritis and calculus are easily distinguished, and in some instances even before the urine is microscopically examined. The microscopic examination of the urine so drawn in the hands of experts will to-day differentiate the various pathologic kidney lesions.

Bacteriologic examination even more than the above gives the greatest amount of information concerning the various infections with which the human genitourinary tract abound. The distention of the kidney pelvis with a small quantity of sterile water or antiseptic solution will often reproduce exactly the symptoms that exist during an attack. Leaving the catheters in place for several hours gives a very good comparative test of the functioning power of each kidney, and if the diseased organ is not functioning all the more reason for its removal. The first urine escaping from the catheter is often stained with blood and should not be retained for examination unless the bleeding persists. A few blood cells found microscopically and no other renal elements should not be taken to signify a pathologic condition.

Lastly, the introduction of the ureteral catheter prior to performing complicated pelvic surgery or hysterectomy is a safe guide in avoiding injury to the ureter. I have once seen the ureter cut in a Kraske operation. Even though it was successfully transplanted it would not have happened had there been the catheter as a guide.

Till some better method is at our disposal let us take the optimistic view of this procedure which, I feel, is to stay supreme for a long time.

83 St. Paul Street.

SYSTEMATIZED TECHNIC OF SUPRAPUBIC PROSTATECTOMY

G. KOLISCHER, M.D.
CHICAGO

Since the inauguration by Belfield of the suprapubic operation for relief of prostatic hypertrophy, and its further development by McGill, Freyer, Guiteras, Burckhardt and others, a certain standstill seems to have taken place in the further elaboration of its technic, while perineal prostatectomy has become a competitor, especially since the Zuckerkandl, Rydygier, and the French technics were adapted and amplified by Young.

The definite success of suprapubic prostatectomy will depend, as that of other surgical methods, on the reduction of primary mortality and the securing of good and permanent functional results. An analysis of the basic factors on which these two issues depend will show the way to their improvement.

Primary mortality is essentially influenced by the immediate and remote dangers of anesthesia, by hemorrhage and by infection, especially of the peritoneum and the prostatic venous plexus. The primary mortality is, furthermore, influenced by the length of time patients are kept in bed after operation; old individuals, whose hearts and excretory organs are, if not already materially injured, certainly impaired in resistance, do not very well stand a prolonged confinement in bed. The functional results depend on the extent and nature of the surgical injuries inflicted on the tissues by the operation and on the conditions resulting therefrom. Bruising and tearing the tissues will not only interfere with quick healing, but will also tend to result in cicatricial distortions which will impede the urinary flow. If favorable conditions for the obliteration of the bed of the prostate are not secured, that cavity will remain a granulating basin in which stagnation of the urine, with all its undesirable consequences, will again occur.

The working points, from a consideration of which the evolution of a successful technic could be developed, might be enumerated as follows: In order to reduce to a minimum the primary and remote dangers of anesthesia, nitrous oxid and oxygen narcosis should be employed. The absence of pronounced or prolonged malaise after this anesthetic enables the patient to take food and drink very soon after operation. The absence of prostration and stupor, so common after ether or chloroform, does away with paresis of the intestines and renders it possible to start out a few hours after operation with the regular breathing exercises so important in old patients. Since nitrous oxid does not attack the heart muscle, the circulation is left unimpaired and hypostatic pneumonia and embolism do not occur. All these conditions make it possible to let the patient sit up twenty-four hours after operation and to begin taking him out of bed on the third day.

In order to prevent involvement of the peritoneum by an infection which might already have existed in the bladder or which might appear during the course of the healing process, the peritoneal duplication should, before the bladder is opened, be removed from the field of manipulations and should be entirely and securely secluded from the operative wound. In order to prevent tearing and subsequent thrombosis and infection of the prostatic venous plexus, the shelling out of the prostatic tumor should be done in such a way that this plexus is never touched. In order to prevent superficial necrosis of the denuded surfaces and to foster their quick healing, all traditional drainage of the perivesical spaces should be abolished; such drainage, furthermore, leads quite often to infection of the concerned cavity from without. In order to create in the operative field conditions most favorable for obtaining good functional results, the various operative steps should be adapted to the specific character of the prostatic hypertrophy encountered. The recognition of such conditions can be obtained only if the prevailing formations can be not merely palpated, but also seen, for faultless technic depends on the possibility of doing

all the operative steps under the guidance of the eye. This means that the inside of the bladder must be freely exposed through a liberal incision; the incision leading down to the prostate and the subsequent enucleation of the tumor can then be modified according to the conditions peculiar to the case.

A comparison of the technic of various surgeons as gleaned from personal observation and from publications, together with a study of the different types of prostatic tumefaction, as well as various attempts to adapt the different steps of the operation to the above-mentioned demands and points, led me to the development of a certain technic in my cases which has so far proved very satisfactory. It is my purpose, first, to describe the steps of this operation as they are common to all the cases, after which the modifications rendered necessary by the peculiarities of various types will be specified.

Previous to the opening of the abdomen, the bladder is filled with a lukewarm 2 per cent. collargol solution; the amount of fluid injected depends on the condition and size of the bladder. The injection of this fluid should be kept up until the bladder is moderately distended; in other words, until the muscular resistance of the bladder against further distention is perceived; it is, therefore, best, before the patient is anesthetized, to make this injection with a smoothly running hand-syringe. The catheter used for injection is left in place and, in order to prevent its slipping out, a gauze strip is tied around the penis just tightly enough to prevent any dislocation of the catheter. Care should be taken not to strangle the member by this manipulation; otherwise extensive subcutaneous hemorrhage or edema might follow. If a soft rubber catheter is used, the escape of the fluid is prevented by tying a loop in its distal end; in case an elastic Mercier is employed, it is closed by a little wooden stopper. Even in cases that do not permit of any distention of the bladder on account of certain inflammatory conditions, a catheter is inserted and left in place so as to mark the internal urethral opening after the bladder is once incised. The patient is then anesthetized and afterward placed in Trendelenburg's position.

A median incision over the region of the bladder is now made through the skin; this incision goes well down to the symphysis and is carried upward above the border of the bladder dulness. The incision must be of this extent, because its upper end, as will be shown later, is closed up again before the bladder is opened, and sufficient space must be reserved for the ensuing manipulations. The subcutaneous fat and fascia are also split with the knife and all bleeding points carefully attended to. The muscles appearing in the bottom of the wound are separated by blunt dissection, exposing the perivesical and preperitoneal fat and the peritoneal duplication. Both the recti are now drawn aside by retractors and the perivesical fat and the peritoneal fold are stripped together from the bladder and then pushed upward by means of a gauze sponge; this detaching is carried on until the whole anterior aspect of the bladder is freely exposed. The perivesical fat and the peritoneal fold are then caught in a suture running transversely from the uppermost part of the edge of one rectus, through the fat and the fold, to the uppermost part of the edge of the other rectus. An assistant pushes this entire flap underneath the muscles while the suture is tied. Two more

sutures reunite the fascia and bury the flap completely, so that from now on the peritoneum is kept absolutely out of the field of operation and out of the reach of any existing or subsequent infection. Two sutures are now inserted at either side of the intended bladder incision pretty close to the symphysis; these sutures are later on used as guy-ropes for lifting up the edges and the base of the bladder after the viscus has been opened.

Fig. 1.—The prostate exposed through the mucosa incision and seized by vulsellum; a, edge of severed mucosa; b, prostate.

Fig. 2.—Partial enucleation of the prostate and lifting up of the loosened part of the gland above the incision; c, prostate.

The nature of the prostatic tumor is now determined by inspection and palpation; this information, together with the above-mentioned landmarks, enables us to

Fig. 4.—Prostate delivered in front of the mucosa incision.

The bladder is then incised in a vertical line by means of a pointed bistoury. The fluid pouring out of the bladder is taken up by gauze pads until everything inside and outside of the bladder is perfectly dry. The

Fig. 3.—Further progress of enucleation; a, vesical mucosa; b, prostate.

incision in the bladder wall is then enlarged until the base of the bladder and the internal urethral orifice are brought into plain view.

A Simons or Martin vaginal retractor is now inserted in the upper angle of the bladder wall while the lower angle is stretched and pushed back by the insertion of a

bayonet-shaped vaginal lever; in this way the field of the endovesical operation is made satisfactorily accessible. The tip of the catheter plainly marks the internal urethral orifice, while the intraureteral ligament and the ureteral openings can also be readily seen. This free exposure of the inside of the bladder also makes it possible to avoid incising the atheromatous arteries which quite frequently run over the prostatic region.

decide whether a partial exsection is sufficient, as is sometimes the case, and just where the incision into the vesical mucosa can be made to the best advantage.

The introduction of one or two fingers of the operator into the rectum, in order to push the prostate upward, I have abolished entirely as non-aseptic. Only in rare cases of very soft prostates it might seem to be to advantage to mark the prostate better by such a manipulation, and then the fingers of an assistant should be made use of for this purpose. The prostate is fixed and lifted up into the field of interference by grasping its most prominent part by means of a vulsellum, after which an incision is made with a knife into the mucosa of the bladder at the selected place. The use of the finger-nail, instead, in digging down through the mucosa to the prostate is clearly a surgical impropriety. Bruised and torn tissue does not heal readily, and the statement that it is easier and safer to find the right cleavage between the mucosa and the prostate by means of this blunt digging does not hold good if one operates under the guidance of the eye. I think, furthermore, that at the present time few surgeons, at least no American surgeons, would operate without rubber gloves, which fact would certainly make this digging more difficult and time-wasting. The incision should not be made too short, for if the tumor is forced through a small opening tears in the mucosa might result, although the elasticity of the mucosa insures a certain amount of stretching.

The incision into the mucosa is deepened until a white layer appears. There are now two possibilities: If the incision has been made over a portion of the tumor that protruded prominently into the viscus, the true capsule of the prostate will be immediately recognized from the fact that the palpating finger does not find a movable layer over the prostate, whereas, if the incision has been made near the base of a prominence, the first white layer to appear is the so-called sheath

of the prostate, recognized by its mobility over the underlying prostate and its true capsule. In case this sheath is encountered, it is also split by the knife, and the prostate, covered by its adherent true capsule, appears in the incision. This exposed portion of the prostate is now seized with a vulsellum and dragged upward under moderate tension (Fig. 1).

A poker or the tip of the index-finger is then introduced back of the vulsellum, between the prostate and the mucosa and sheath, and the covering layers are stripped from the gland (Figs. 2 and 3). Two principles should be kept in mind during this proceeding: The prostate with its capsule should be enucleated from the sheath formed by the rectovesical fascia, and the enucleation should proceed from the posterior surface of the tumor toward the symphysis. While this enucleation is being carried on, the gradually appearing parts of the prostate are seized progressively by vulsella and lifted out in the general direction of the enucleation while the index-finger, by slitting motions, detaches it from its surroundings.

Fig. 5.—Solitary pedunculated excrescence of the prostate; 1, incision in vesical mucosa; 3, urethral orifice; 4, 4, ureteral openings. Half-schematic drawing.

Thus the enucleation is done under the constant guidance of the eye. One should be careful to watch for tissue strands running from the capsule of the prostate to its sheath; these strands should not be torn, but should be severed with a knife or curved scissors. In this way injury to the sheath and breaking into the prostatic venous plexus are avoided. After the entire prostate is freed and lifted up, the finger detaches it from the triangular ligament. This is the only step of the enucleation that has to be done in the dark and by palpation alone; no difficulty is encountered, however, because by this time the entire prostate has been developed into the cavity of the bladder and no possible harm can be done by rolling it off from the ligament (Fig. 4). This enucleation almost always produces the organ *in toto*. Should such not be the case the remaining portion is again seized by a vulsellum and removed

in the same manner. The index-finger is then reintroduced into the cavity and a careful search is made for small nodules which are occasionally left in the depth. These also are enucleated with the index-finger in order to avoid a recurrence of the tumefaction. After all this is done the bladder and the prostatic side are flushed out with a hot 2 per cent. collargol solution until the fluid returns clear. Everything is then dried by means of large gauze sponges and the catheter is removed.

The mucosa flap covering the prostatic cavity, which by this time has shrunk considerably, is now pressed down toward the underlying tissue and kept in this position by packing the bladder with a wide strip of very weak iodoform or of similar gauze. The end of this strip is led to the surface and the bladder closed with catgut sutures, save for a little gap in the vertex which permits the emanation of the gauze. These sutures are inserted in such a way as to bring the denuded surfaces of the muscularis together and to avoid any impaction of the mucosa. After the bladders of

Fig. 6.—Pedunculated median lobe protruding into the bladder coincident with general hypertrophy of the prostate; 1, first incision into the vesical mucosa; 2, occasionally necessary incision; 3, urethral orifice; 4, 4, ureteral openings. Half-schematic drawing.

old people are opened the mucosa will as a rule detach itself spontaneously from the muscular coat, appearing as a fine border; if not, it is detached to a small extent with a knife-handle or small poker in such a manner that when the sutures are inserted the mucosa lips will fold themselves together and protrude like an edge into the cavity of the bladder. After this the fascia of the muscles is reunited by catgut sutures up to the drainage openings; the skin is then closed in the same manner.

The gauze strip just mentioned also serves the purpose of leading the urine out of the bladder until natural urination takes place. It is not only unnecessary but also disadvantageous to insert a rubber tube deep into the bladder, as this necessitates leaving a larger opening; moreover, the pressure of such an extraneous

body often leads to necrosis. One should not forget that the bladder, being a contractile organ, drains itself, as it were.

I have entirely abandoned both suspension of the bladder to the abdominal wall and drainage of the perivesical space. The suspension of the bladder tends to excite spasmodic contractions of that organ which interfere with the knitting together of the tissues and cause pain and discomfort to the patient. I was induced to do away with drainage of the perivesical space by the fact that in my experience this space is entirely obliterated by natural pressure after the patient is brought back into a recumbent position, and that the abolition of drainage leads to quicker healing. This observation was also made and published by several German surgeons.

Fig. 7.—Ring-shaped hypertrophy of the prostate surrounding the whole prostatic urethra; 1, incision through vesical mucosa; 3, urethral orifice; 4, 4, ureteral openings. Half-schematic drawing.

An extremely vital step in the operation is the determination of the location of the incision through the vesical mucosa, for the proper selection of this location ensures a greater facility in enucleating the tumor and the most favorable conditions of the cavity and wound after removal has been completed. The importance of a definite diagnosis after opening the bladder is exemplified by the type of prostatic obstruction illustrated in Figure 5, which shows the protrusion into the bladder of the median lobe of an otherwise untumefied gland in such a way as to act like a ball-and-socket valve. In a case like this it would be absolutely unnecessary to remove the entire gland, the exsection of the pedicle of this lobe, as indicated in Figure 5, being sufficient. The pedicle is circumscribed by an oval incision and a

wedge-shaped piece of the underlying tissue is exsected in continuity with the pedicle. The resulting wound is then firmly closed by a few deep catgut sutures and the bladder sewed up completely. As a rule patients urinate naturally a few hours after operation; in the rare cases in which they are unable to do so the bladder is emptied at proper intervals by means of a soft catheter. A permanent catheter should never be employed.

Another type is seen in the development of a more or less distinctly pedunculated median lobe in addition to hypertrophy of other parts of the gland, and is marked by the pushing of the base of the bladder into the cavity (Fig. 6). In such an event the protruding lobe is seized at its top by a vulsellum and dragged upward until the basal mucosa becomes tense. A crescent-shaped incision into the mucosa is then made somewhat distant from the base of the pedicle, as indicated by the heavy line in Figure 6. After the prostatic surface is exposed the gland is enucleated in the manner de-

Fig. 8.—V-shaped hypertrophy of both lateral lobes; 1, summit of lateral lobe; a, b, incision through vesical mucosa; 3, urethral orifice; 4, 4, ureteral openings. Half-schematic drawing.

scribed above. The enucleation often runs along quite smoothly until the junction of the bladder mucosa with the tumor in front of the pedunculated lobe is reached. If the detaching of this part of the mucosa seems difficult, or if the resulting anterior flap appears a little too large, the mucosa is severed by an incision indicated by the dotted line in Figure 6.

In those cases in which the anterior portion of the prostate also participates in the enlargement, so that a hypertrophied ring encircles the whole urethra, the incision through the mucosa is best made in such a way as to half encircle the urethral opening; this semicircular incision is made either to the right or to the left, according to the side of greater prominence (Fig. 7). Here, too, the enucleation of the prostate is done under

Fig. 9.—Hypertrophy of prostate with protrusion into the bladder of the median lobe carrying along the prostatic urethra; 1. line of incision through vesical mucosa; 3. urethral orifice; 4. 4. ureteral openings; 5. median lobe of prostate. Half-schematic drawing.

Fig. 10.—Median lobe of the prostate growing into the bladder as a massive tumor; 1. anterior circumference of mucosa incision; 2. posterior circumference of mucosa incision; 3. urethral orifice; 4. 4. ureteral openings; 5. massive tumor of prostate. Half-schematic drawing.

the guidance of the eye and by gradually lifting the tumor out by means of progressing vulsella. On this occasion I wish to mention that the partial or total removal of the prostatic urethra does not in the least interfere with a smooth healing and a perfect final result, provided that the after-treatment be carried out in a manner adapted to this particular incident.

In instances where the hypertrophy is confined mainly to both lateral lobes, so that what is commonly called a "prostatic thor" results, the incision is made over the most prominent contour of one of the lobes, as indicated in Figure 8, and the enucleation is performed in the above described manner.

Another type of prostatic hypertrophy, as illustrated in Figure 9, demonstrates again how important it is to make a correct final diagnosis after the bladder is opened. It will occur in certain cases that the median lobe, by growing into the viscus, carries along the cen-

Fig. 11.—Collar-shaped hypertrophied prostate; 1. line of incision through mucosa; 2. collar-shaped protrusion of prostate into the bladder; 3. urethral orifice. Half-schematic drawing.

tral end of the urethra so that the internal urethral orifice appears at or very near the top of the tumor. Were this incision to be made indiscriminately—for instance, were it to cross the anterior surface of the tumor—severe injury to the urethral end, with all its complications such as hemorrhage or traumatic stricture, could not be avoided. In such instances the mucosa incision should encircle the posterior base of the tumor, as indicated by the dotted line, and the enucleation thus begun will proceed without difficulty and without dangerous lesion to the urethra.

In other cases the median lobe will grow into the bladder as a massive tumor of such extent that it practically divides that organ into two halves (Fig. 10). It is entirely erroneous to make the incision, as is so often done, over the top of a tumor of this description, for such an incision means only unnecessary hemorrhage

and loss of time. It is impossible to find the right cleavage at the top of such tumors because there is no sheath and because the true capsule of the prostate and the vesical mucosa are thinned out to such an extent in this location that they scarcely can be isolated. In such instances the base of the tumor is encircled by a complete circumcision as indicated in Figure 10. This incision should be made in such a way as not to interfere with the urethral openings which, according to the nature of the case, may lie in front of or behind the tumor. The enucleation is then started from the posterior incision.

In cases of collar-shaped prostate it is important to grasp the prostate with a vulsellum in the proper manner. It is a mistake to apply the forceps to the protruding edge at right angles to the elevation, for if this be done the forceps or vulsellum will tear out and unnecessary injury and hemorrhage will be produced. The vulsellum should be applied at the posterior aspect of the collar, somewhat above its junction with the base of the bladder (Fig. 11). This incision through the mucosa is made parallel to the collar and posteriorly from it, somewhat distant from the insertion of the vulsellum. The enucleation of the prostate is then carried on from behind forward. Thus the mucosa adjacent to the internal urethral orifice rolls off easily from the tumor and any injury to the urethral mouth as well as any tearing of the edges of the vesical mucosa is avoided.

92 State Street.

IGNORANCE OR MALPRACTICE?

FINANCIAL AND CLIMATIC CONDITIONS IN THE TREATMENT OF TUBERCULOSIS

W. WARNER WATKINS, M.D.

PHOENIX, ARIZ.

The object of this communication is to call attention to a phase of the tuberculosis question which is being treated with reprehensible neglect by far too large a percentage of general practitioners. It must be understood that these remarks apply solely to a particular class of patients and to a particular custom of which they are the unfortunate subjects; the word "custom" is used advisedly since the practice does not merit the term "treatment."

The patients are those who have reached an advanced stage of phthisis and in whom the disease is making steady progress, insidious or rapid as the case may be; whose financial means are small and who can not, under the most favorable surroundings, provide themselves with more than the necessities of life or who perhaps are the actual wards of charity.

The custom is that, in the face of all known teachings concerning the disease and in the face of common sense and humanity, physicians will persist in sending this class of patients away from home, family and friends to die in a strange land, by holding out to them the fatuous hope of recovery in a "change of climate."

There is no intention here to disparage the value of a suitable climate in the treatment of pulmonary diseases; but every intelligent physician knows that any climate, however favorable, is valuable only as a subsidiary adjunct to one of the three essentials for the constitutional upbuilding required to check the progress of a pulmonary tuberculosis, these essentials being rest, proper food and fresh air. The Salt River Valley may have the

most equable climate and the finest atmosphere in the world—and we who live here firmly believe that it has—yet any physician who advises a patient to sacrifice the first two essentials for the sake of the third is guilty of something more than mere ignorance, for in this day of free knowledge ignorance on this point is sin. Not even 360 days of sunshine each year can equip our climate with any curative value in tuberculosis when the system is deprived of food and the patient of rest or, even when these are granted, when the patient is hopelessly diseased. I am as firm an advocate as any one of this and similar climates as an adjunct to outdoor life (fresh air) when the patient is financially able to provide himself with the other more vital essentials of rest and proper food. But if he has not the means to place himself under the direction of a physician, if he is not able to pay for the necessary care when he is required to rest in bed or otherwise, if he is not able to provide himself with the very substantial food which is always required, then he can hope for absolutely no more improvement in Arizona or any other resort than will come to him at his own home—wherever that may be—with the advantage of the solicitous care of family and friends. The physician who advises him otherwise is guilty either of ignorance or malpractice; and even with all these advantages possible, moribund patients can hope for nothing more here than at home and should be so advised.

It seems inconceivable that physicians can dwell in enlightened communities and remain ignorant of the rudiments of pathology and therapeutics as applied to tuberculous processes when the entire medical world is surcharged with the purpose of educating even the public in these facts. But that such ignorance does exist is only too evident to those of us who practice in communities like this, where are congregated the victims of such professional ignorance, or, if not ignorance, then malpractice, pure, simple and unadulterated. It is deception as cruel and heartless as are the methods of the ubiquitous fakers against whom such a relentless warfare is being waged.

CASE 1.—A boy, aged 16, worked on an elevator in Pittsburgh, Pa., until forced by a hemorrhage to stop; his mother being dead, his father a loafer, the boy was dependent on his own efforts. Some business man gave him \$150 and on the advice of a prominent physician of Pittsburgh he came to Arizona with these instructions: "Live outdoors, stay away from doctors and you will get well." A charitable institution found him and referred the case for admission to St. Luke's Home, to whose staff I belong. The patient was found to be in the advanced stages of pulmonary tuberculosis with consolidation, cavities and visceral involvement. He was not admissible to St. Luke's and was advised to return by the first train to his home town and friends, which he did.

CASE 2.—A young married woman (two months pregnant) with acute miliary tuberculosis was sent to Phoenix from one of the most enlightened cities of southern California; she spent her last dollar for traveling expenses and arrived in Phoenix absolutely destitute, not having means to provide even a night's lodging. Charitable people learned her plight, collected a sum of money and placed her in a tent colony for two weeks, from which she passed first into the County Hospital and then under the care of the Associated Charities, which provided for her until she died. Her physician in California, in whom she had placed implicit confidence and who knew her financial condition, had told her that Arizona was her only hope for recovery.

CASE 3.—A young man was sent here from Oregon, with an advanced tuberculosis of three years' standing which had finally reached that stage of morbidity where Arizona was the "only hope" in the judgment of his physician. The patient learned from the same source that all he had to do was to