the deeper air passages and sets up a chronic irritation. The question of the relation between chronic catarrh of the upper and of the deeper air passages has not received the attention it deserves; it is barely mentioned in the most general terms in connection with bronchitis, and the possibility of emphysema; bronchiectasis or fetid bronchitis being due to such causes is usually ignored.

The peculiar connection, often definite, again but suggested, between nasal catarrh and rheumatism may perhaps in part be explained by what has been referred to in this paper, namely, the secondary involvement of the throat in nose affections and the later development of dyspepsia and liver derangement followed by symptoms of general malnutrition, and this gradually gliding into rheumatic disease. Later on in life, even though the catarrhal disease be cured or modified, the rheumatism or malnutrition, or both, may remain and be a constant threat to comfort and safety.

When early in life obstruction or inflammatory disease of the nose is present it often, and in adenoid disease practically always, exerts a marked influence on the nutrition, formation and position of the teeth and the development of the dental arches. Thus are caused inequality in the size and structure of the teeth and the formation of secondary, irregular and contracted arches. These effects are especially observable when the child has been affected in early infancy. Unfortunately this influence unless corrected will very soon leave its impress so deep as to follow the child into adult life, its permanent teeth needing much attention and being the cause of faulty mastication and digestion, as well as of difficult respiration.

In a pamphlet³ published some ten years ago I have referred at more or less length to a number of diseases to which nasal disease gives origin. In several other papers⁴ I have called special attention to the symptoms occasioned by such abnormal growths. In a later pamphlet⁵ I called attention to the influence of nasal disease on the curvature of the cornea through a probable action on the muscles of the eyeballs.

I hope I have been able to give you a few thoughts on the subject which may stimulate to further observations which will, I think, aid in proving the proposition that nasal disease, especially in children, is a not infrequent factor in the causation of disease, and that it modifies many diseases, interferes much with the cure of various affections, and that when carefully handled it is promptly controlled by proper local and general treatment.

TREATMENT.

The first indication for treatment is to avoid the causes of nasal diseases, such as colds, exposures, exhaustion, local mechanical irritation, and anything that tends to keep the general vitality below normal.

All obstructions to the proper inspiration of air must be removed, hyperemia and congestion must be modified and reduced by local treatment and internal medicine. Deviated septi must be straightened, enlarged turbinates must be reduced, hypertrophied tissue must be brought to normal size by operative measures, or other treatment or both. All nose diseases must be treated in

connection with the other diseases of which they may have been the cause. In all cases one should persist in the most perfect and antiseptic cleansing possible. Stimulation of the various tissues must be used when necessary, soothing preparations, however, are more often required. Albolon, cosmolen, glycerin and alkalin solutions are in constant demand. Acids, iodine, nitrate of silver, the electric needle, the snare, even the saw and knife have their uses at times. What to use and when to use it, is a matter of judgment.

Internal Treatment.—We must never neglect thorough antiseptic cleansing of the throat and nose, with a careful preparation and covering of the parts by oils or salves after our local nose work is completed. In addition to this, every case of nose disease demands internal treatment. It may be iron or phosphorus, or even iodine, or it may be a digestant or a nerve tonic or an alkali or a laxative. But the medicine which I believe is needed more than any other is cod liver oil.

Hygiene and Diet.—In addition to the avoidance of the cause, the child should be made to live, so far as possible, a perfect hygienic life. He should retire early, get plenty of sleep; take a sponge bath with friction in the morning on arising, if his vitality admits of it. He should wear proper and sufficient underwear for the season; the underwear should be of pure wool, usually very light weight. I believe linen underwear in our climate is unsuitable for the child. He should not be dressed in too heavy clothes. He should wear stout shoes with rubber overshoes whenever it is wet under foot or where there is snow on the ground. He should not use coffee nor tea; he should drink milk or water, with an occasional change to chocolate or cocoa or lemonade, as necessary. His food should not be predigested, but it should be digestible. He should not drink during the course of the meal. He should go out of doors every day, rain or shine, unless he is sick. He should have a proper variety of work, study and play. He should take his meals with regularity and not be fed too often. He should not be denied sweets and candies on all occasions. A child needs sweet foods, and as long as the stomach does not rebel they are agreeing with him. He should run and race and play with other children.

He should, as a rule, take his hot baths at night, on retiring. Beyond all else, perhaps, he should go to bed early. Children need a great deal of sleep; they can absorb all that is allowed them. By sleeping ten or twelve or more hours they are fitting their bodies not only for the present, but for the strain of adult life, as well. They are storing up energy and nerve force for future emergencies.

TREATMENT OF CHOREA BY PROLONGED WARM BATHS.*

W. C. HOLLOPETER, M.D.
PHILADELPHIA.

In this brief contribution I wish to emphasize the virtues of the prolonged warm bath in shortening the attack of chorea.

With the great increase in the number of adults suffering from nervous diseases, it would be natural to expect an alarming increase of chorea among school children. This, however, is fortunately not corroborated by statistics. We have less chorea to-day than we had

3. The Nose as a Frequently Unrecognized Cause of Disease; *Codex Medicus*, Phila., vol. 1, No. 1, p. 17.

4. The Treatment of Nasopharyngeal Adenoids; *Proceedings Amer. Med. Assn.*, June, 1899. The Easiest, Quickest, Safest and Most Thorough Method of Removing the Third Tonsil; *The Penn. Med. Jour.*, April, 1901. The Recognition and Prompt Removal of Postnasal Adenoids in Children; *THE JOURNAL A. M. A.*, March 28, 1903.

5. The Influence of Nasal Disease in Modifying the Quantity and Axis of Corneal Astigmatism; *Ophthal. Rec.*, March, 1897.

* Read at the Fifty fifth Annual Session of the American Medical Association, in the Section on Diseases of Children, and approved for publication by the Executive Committee: Drs. S. W. Kelley, H. M. McClanahan and John C. Cook.

fifty years ago; Watson, writing at that time, analyzed 1,029 cases, against Holt's 146 of to-day. With the population of our cities rapidly increasing, and with this growth the crowding and struggling for higher educational and social preferment, we find the army of neurasthenics markedly augmented among the adult population; and while we have neurotic children, true chorea does not seem to be increasing in the same ratio.

Heredity and environment, the twin jailers of childhood, do not hold their victims securely now that domiciliary hygiene is better appreciated and more widely taught. While we know that few American children of city rearing escape the entanglements of neurotic heredity, we may also state that the very satisfactory advancement made in domiciliary and personal hygiene has neutralized in a great measure the benumbing influence of heredity. The great attention given to our children's comfort and their personal hygiene, in the better classes of society, has done much toward reducing the number of choreic children. The care of the child's teeth; food, and how it is taken; sleep, and its environments; study—all of these influence the development of the child, and all have been greatly improved during the last few years. Much more advanced study, considering the child's age has been demanded—more than is consistent with growing health. The child's intellectual advancement to-day is phenomenal, and this advancement has only been accomplished because of a wider knowledge of hygiene, both personal and domiciliary. Motley and Bancroft graduated from Harvard at 17, but the boys and girls of to-day are competent to cover more ground for their entrance into college than did either Motley or Bancroft at graduation.

Much has been said about pressure in school as a factor in producing nervous conditions. This is also a relative factor. I do not believe that we have so much pressure in our schools as we have poor personal hygiene and home irregularities. The boy who eats rapidly and irregularly of improper food, disregards his teeth and stomach, and the many other personal attentions which are so essential in attaining and maintaining a high standard of health in a growing child, soon lowers his vital resistance, and, in this way makes a bid for chorea, as well as the whole train of acute infectious diseases.

The cramming system in existence in most of our schools has been a favorite subject with many educators. Much can be said on the subject both for and against it, but I think that the cramming takes place more frequently in the stomach than in the brain. Irregular and improper feeding, with careless, exciting home life have more to do with making a nervous child than school life, although the studies be somewhat difficult.

THE RELATION OF RHEUMATISM TO CHOREA.

The relation of chorea to rheumatism is always interesting, because never satisfactorily answered; an equal force of authority being claimed both for and against it. Possibly chorea is a selective action of the rheumatic diplococcus on a nervous system with special predisposition to chorea. It is possible to find rheumatic histories for all our cases of chorea—what parents do not have rheumatism in some form after rearing a family?

REVIEW OF LITERATURE.

The professional drift is undoubtedly in the current leading us to believe that chorea has some youthful resemblance to mature rheumatism. An outline of the literature of the etiology and treatment of chorea will

show that my ideas of hydrotherapy in connection with chorea are not original, but have not been strongly held, therefore no favorable results were secured.

The very interesting discussion on the pathology and treatment of chorea which took place July, 1903, in the British Medical Association, is undoubtedly the most important recent contribution to our knowledge of this very important and interesting disease of childhood. The essence of this discussion was (and the majority of the members were of this opinion) that chorea was a disease of microbic origin—a diplococcus—some relationship existing between it and some previous microbic disease giving us a mixed infection. Drs. Poynton and Pain have isolated a diplococcus from rheumatic cases associated with chorea. A large number of observers have verified in part the results of these investigations.

Dr. E. P. Lees believes that chorea is a disease of the brain. He says concerning the symptomatology of chorea that its most obvious characteristics are disorderly muscular movements which are spasmodic, chronic, irregular, involuntary, and exhibit imperfect control and co-ordination. The muscles which are most affected in chorea are particularly those which are under the most delicate voluntary control, and in view of this fact and of the other attending symptoms, he believes that chorea is unmistakably a disorder of the brain, and particularly of the motor centers. If the cord is involved, it is but rarely, and only occasionally it is observed that the peripheral nerves are also affected.

The value of the routine treatment of rheumatism by the salicylates, in Dr. Lees' opinion, is strong corroborative evidence that the micro-organism is closely related to that of acute articular rheumatism—an English opinion, strictly speaking.

Two very interesting cases which prove but little of what is now taught and believed, and yet suggesting much that is new and interesting, will be sufficient to illustrate my theory. A brief history of each case will tell its own story.

CASE 1.—E., aged 12; the sixth child of German parentage.

Family History.—Father and mother both living and well. Father aged 48, mother 42. Father is a baker, health good; he is a hard worker. He has poor teeth, eats rapidly, has poor digestion, and occasional rheumatic and gouty pains. He has not lost a day from work by reason of sickness for 10 years. Mother is overburdened from childbearing and domestic toil with inefficient help, and may very properly be denominated a neurasthenic, yet performing her usual exacting domestic duties.

History.—The child was sent to school when 6 years old, and after being in school for several weeks commenced to manifest symptoms of chorea; the irregular movements increasing rapidly until every superficial muscle was twitching and jerking. The child could neither stand, walk nor rest. Every joint was sore and inflamed by reason of friction.

Treatment and Result.—She was encased in cotton and bandaged to protect the skin and limit friction. In 12 weeks the violence of the movements subsided and convalescence was established. The child has had six annual attacks, all commencing in the fall (September) and lasting from 3 to 4 months.

At no time did we find fever. This child has no heart lesion, and contrary to the most eminent authority has not been, nor is she at present, anemic.

Remarks.—The child's history is interesting because of the annual attacks and their violence; the duration, the absence of anemia and heart murmur. The last three attacks were materially shortened by prolonged warm baths.

The other case is that of a child whose family have for a long time been my patients.

CASE 2.—The oldest daughter died from heart and kidney sequelæ of scarlet fever. The father is inclined to have rheumatic manifestations, but the mother has unusually good health; even the grandparents possess to a slight degree the stigma of the rheumatic tendency.

History.—F., the second daughter, is the same age as the child mentioned previously. She has had five annual attacks of chorea, commencing each time at the end of her school year, thus showing nerve fatigue and loss of control. The attacks of the two earliest years lasted from three to four months; those of the last three years terminating within six weeks by means of my hydropathic treatment.

Treatment.—This treatment I have found to be eminently satisfactory. The usual detail as to personal wants, food and sleep having been fully observed, I found by accident, that the warm general bath was unusually soothing to the exhausted, twitching muscles. I noticed that after the bath, the child had a longer and more refreshing sleep; a sleep that strengthened and fortified her for the day; so rendering each day more comfortable than the preceding.

TECHNIC OF THE BATH.

The technic of the bath, which constitutes the entire and most satisfactory treatment in my hands, is to place the child in a bath at a temperature that will prove no shock nor surprise to it; a temperature of from 90 to 98 is sufficient, according to individual peculiarities, and cool down so as to be soothing to the whole cutaneous surface. Washing the face before the child is put into the bath is oftentimes advantageous.

The child, when once placed in the bath, is entertained by the nurse or mother by placing some plaything or floating toy in the water, and while the child's attention is so occupied the time will pass rapidly and the attendant's duty will consist in avoiding draughts and in maintaining the temperature of the water sufficiently high (from 90 to 96) so as not to produce any immediate or secondary shock to the child. In this way, the child may be entertained for a long time in the water; the whole body being immersed except the head and neck for at least one or two hours, twice a day. The last ten minutes of the time can be properly devoted to a gentle superficial massage of the arms, legs and trunk; after which the child will be sufficiently weary to take a nap.

I have treated probably 40 or 50 cases by means of the prolonged bath, following this method of procedure with the result of shortening the duration of the attacks from 3 months to 6 weeks.

DISCUSSION.

DR. C. F. WAHRER, Fort Madison, Iowa, said that he was certain that many cases of chorea, so called and so treated, are not chorea at all, in spite of the fact that many physicians think it is easy to diagnose chorea. He mentioned a boy about 8 years old who developed certain anomalous movements of the head; these would be frequently repeated, and terminated usually in a slow rotation of the head, also a slow shrugging of one shoulder. Within a few years these movements ceased, and instead of them he gave vent occasionally to a peculiar expression sounding like "ha-ha." This in turn ceased and gave way to an ear-splitting whistle, of which he was apparently entirely unconscious. Subsequently he developed other peculiar phenomena, such as licking the ends of his fingers, before answering when spoken to. He was treated in the usual manner with arsenic and antirheumatic remedies without any appreciable benefit. He was a very bright boy, distinguishing himself at his studies, and subsequently graduated in medicine and took a postgraduate course. He is now in active practice, and although an able

thinker and fine scholar, he still develops these little peculiarities. At present he frequently gives vent to a dry cough, which is particularly noticeable when his attention is relaxed. He has no heart lesion. Dr. Wahrer never regarded this as a case of chorea, but rather one of tic or habit spasm. Such cases are comparatively common, and the usual treatment for chorea has no effect on them. When the patient's attention is called to these movements he is able to stop them; this is not so in true chorea.

DR. W. C. HOLLOPETER said that by chorea he did not mean the cases of habit spasm to which Dr. Wahrer referred. Some children constantly bite their fingernails or have other forms of habit spasm, but those are not cases of true chorea. Dr. Hollopeter gave small doses of arsenic or iron to some of his patients who were anemic, but in the last series of fifteen cases he withdrew drugs entirely. Every child was examined carefully, and if there was any nasal obstruction, or stomach or intestinal disorder, the defect was corrected so far as possible. Dr. Hollopeter tried the hydropathic treatment as an auxiliary to the rest treatment and found it beneficial, but exceedingly irksome to the patients. Many would not remain in bed, but they had no objection to remaining in the bath, playing with boats and various water toys. He believes also that the water acts as a better support than the bed and is much more comfortable than the bandage sometimes applied to the extremities and exposed parts to prevent friction. In those cases in which a heart lesion existed it was not aggravated by the water treatment.

DOUBLE RADIAL RUPTURE OF THE IRIS.*

S. C. AYRES, M.D.

CINCINNATI.

This case of double radial rupture of the iris is very remarkable and, so far as I have consulted the literature on the subject, there is only one which corresponds to it. In that case, which occurred in the practice of Dr. Büniger,¹ in 1821, the left eye was struck by a chain, causing an artificial pupil which could be distinguished from one made by an operation, only by a band of iris remaining at the ciliary margin. At first, sight was very much impaired, probably from intraocular hemorrhage, but perfect vision was finally regained.

My case is as follows:

History.—G. H., age 10, was first examined Nov. 12, 1902, and gave the following account of the accident: He was struck by a BB shot from an airgun at a distance of about five feet. The shot struck the edge of the lower lid of the left eye at about its anterior third. It embedded itself in the bulbs of the ciliæ and almost passed through. It was removed at home by squeezing the lid, when it dropped out. I saw the boy five days later and found a well-defined coloboma of the iris as clean and sharply cut as if it had been done by an expert. It was about 4 mm. in width. There was no blood in the anterior chamber, and only a moderate injection of the conjunctiva. Lying in the coloboma was a mass of iris tissue. It was entirely separated from the iris on both sides as well as from its ciliary attachment (Fig. 1). The lens was opaque and vision was reduced to perception of light, probably on account of intraocular hemorrhage. Tension was normal and the eye not painful. In my judgment the posterior lens capsule was ruptured, as there was no evidence of soft lens substance in the anterior chamber at any time. Atropin was ordered.

In the course of time the edges of the coloboma gradually approached each other and the upper and inner portions of

* Read at the Fifty-fifth Annual Session of the American Medical Association, in the Section on Ophthalmology, and approved for publication by the Executive Committee: Drs. Frank Allport, John E. Weeks and R. L. Randolph.

1. Graefe and Walther: *Jour. d. Chir. und Augenheilk.*, vol. ii, p. 669.