magnet from one limb to another on the opposite side, just as may be the developed contracture. As an illustration of the method of obtaining these results an instance is described in detail. The patient, having been hypnotized, is in a state of lethargy, with muscular hyper-excitability; but it would not do to apply Esmarch's bandage while she is in that condition, because the mechanical stimulation would probably cause a contracture of the whole limb. The patient's eyelids are therefore raised. She at once becomes cataleptic, the limbs retain any position in which they may be placed, but the muscles have ceased to be sensitive to mechanical excitation. Advantage is taken of this loss of excitability to apply the bandage. The eyelids are then pressed down. The catalepsy at once ceases, the patient resumes the condition of lethargy, and the muscles recover their excitability. Then, without touching the limb elsewhere, a small button is pressed several times against the ulnar nerve behind the elbow. No effect follows until the bandage is removed, and then, as blood returns to the limb, the hand slowly assumes the position due to contracture of the muscles supplied by the ulnar nerve, especially the interossei. In this experiment the excitation of the ulnar nerve has impressed the nerve centre in a peculiar way, but only in the parts corresponding to the ulnar nerve. The nerve action for the contracture in the muscles supplied by the nerve takes place in the nerve centre, but not in the periphery until the return of blood to the muscles restores their contractility. Again, the patient being asleep, the arm is anæmiated. The ulnar nerve is then stimulated, of course without effect. Nevertheless, the spinal cord has been influenced, and if the circulation were to be restored the contracture would come on. But if, instead of this, a magnet be applied to the other arm, the contracture is produced in this in exactly the same way as if it were actually transferred from the first side. The lateral contracture has been transferred. These phenomena, in the opinion of M. Charcot and M. Richet, are evidence of the reflex nature of the neuro-muscular hyper-excitability, whether produced by the mechanical stimulation of a nerve, a muscle, or a tendon.—*Lancet*, May 7, 1881.

Treatment of Diphtheria.

WEISE states (*Berl. Klin. Wochen.* No. 4), as the result of his experience of Guttman's treatment of diphtheria by pilocarpin, that it produces an excellent and rapid effect in many cases. His own treatment, under which he has had fifty-four cases without a death, is with salicylic acid and benzoate of soda. Every hour and a half the patient inhales, or has his throat painted with, a solution of salicylic acid (one part, by weight, to fifty of glycerine and rectified spirit in equal parts), and at the same time takes benzoate of soda internally, and stimulants. The inhalation is given with an instrument constructed by Dr. Weise, consisting of a small spray-apparatus combined with a tongue depressor.—*London Med. Record*, April 15, 1881.

Treatment of Pharyngeal Diphtheria.

Dr. OERTEL directs his treatment (*Archives of Laryngol.*, Jan. 1, 1881), first, to the destruction of the cause of the disease; and second, to the removal of the products from the affected parts. For the former, he thinks carbolic acid a most efficient remedy. When the results obtained from its use have been unsatisfactory, the solution has not been sufficiently strong. Of twenty-seven of the graver cases which have been treated by Dr. Oertel, he believes that three-fourths would have ended fatally but for the use of carbolic acid. He employs a five per cent. solution nebulized by means of a steam-atomizer, and avoids the use of a brush or sponge, as he thinks these only irritate the parts. He considers the absorption of the carbolic acid