SELECTED ABSTRACTS.

The Subcutaneous Injection of Paraffin for the Correction of Deformities, with Report of Cases—JUNIOUS F. LYNCH, M. D.
Norfolk, Va.—*Virginia Medical Semi-Monthly*. November, 1902.

The author records his experience with this subcutaneous injection of paraffine for the correction of "saddle nose," and illustrates four cases "before" and "after," in which there were satisfactory results. He employed soft paraffin, or white vaseline, and not the hard substance from which candles are made. It should have a melting point above 99° and below 104° F., for if it is too soft it will be taken up by the lymphatics and if too hard necrosis will result. The preparation used has a melting point of 102° F., and is prepared by Mr. W. B. Martin, of Norfolk. The technique is simple. The field of operation is prepared as for any other surgical procedure, and the instrument and paraffin sterilized. The paraffin injection is preceded by that of a few drops of a four per cent. solution of cocain for painlessness. The melted paraffin is then drawn into the syringe and allowed to cool until it emerges from the needle a worm-like mass; the needle is then inserted above the site of the depression and the injection is made, the nose at the same time being molded to the shape desired. After the removal of the needle an antiseptic collodion dressing over the puncture is all that is needed. The reaction is very slight and usually confined to a feeling of fullness and tension and tenderness of the nose lasting for a day or two. No unsightly bandages are necessary, no time is lost from business and all of the author's cases left the operating room to go to work. After a time the paraffine becomes encapsulated and of cartilaginous consistence. As a rule only one injection is necessary, and there is no danger.

F. C. E.

Bony Cysts of the Middle Turbinate Body—GEORGE E. SHAMBAUGH, M. D.—*Annals of Surgery*. July, 1902.

The author gives interesting histories of three cases of bony cysts of the concha bullosa and advances the theory that the concha bullosa should be looked upon not as an inflammatory product, but as an anatomical variation, the result of a developmental anomaly.

E. D. LEDERMAN.