The Boston Medical and Surgical Journal

TABLE OF CONTENTS

April 25, 1918

ORIGINAL ARTICLES

- WOUNDS OF THE JOINTS. *By L. Eloesser, M.D., San Francisco* 555
TRENCH SANITATION. *By Captain C. E. Burt, M.R.C., Camp Grant, Ill.* 567
PREVENTION IN PUBLIC HEALTH. *By D. M. Lewis, M.D., New Haven, Conn.* 571

MEDICAL PROGRESS

- PROGRESS IN GYNECOLOGY. *By Stephen Rushmore, M.D., Boston* 574

BOOK REVIEWS

- Pocket Formulary. *By E. Quin Thornton, M.D.* 578
Lectures on Medicine—A Handbook for Nurses. *By Chalmers Watson, M.D.* 578

EDITORIALS

- MEDICAL RESERVE CORPS. 579

- ANNUAL REPORT OF THE CANCER COMMISSION OF HARVARD UNIVERSITY. 580
METHODS FOR FIELD STUDY OF INDUSTRIAL FATIGUE. 580
THE PRESENT SMALLPOX SITUATION. 581
THIRD ANNUAL REPORT OF THE FORSYTH DENTAL INFIRMARY. 592
SCHOOL OF INSTRUCTION ON PUBLIC HEALTH. 583
MEDICAL NOTES. 583

THE MASSACHUSETTS MEDICAL SOCIETY

- NOTES FROM THE DISTRICT SOCIETIES. 585

MISCELLANY

- SUGGESTIONS FOR STATE BOARD OF HEALTH REGULATIONS. 586
MUNICIPAL CIVIL SERVICE COMMISSION, NEW YORK. 588
CENSORS' EXAMINATIONS. 588

Original Articles.

WOUNDS OF THE JOINTS.

BY L. ELOESSER, M.D., SAN FRANCISCO.

[From the Division of Surgery, Medical Department, Stanford University.]

THE mass of material that has accumulated during the present war has served to change many of our notions as to the treatment of injuries to the joints.

The difficulties of treatment are largely mechanical and due to the complex anatomy. We speak of a joint cavity; more accurately, each joint is a series of cavities with more or less open communication—a series of cavities with many interdependent pouches and pockets. Even the main cavity of a joint is not an entirely free space; in hinged joints it is divided by the bones and the intra-articular ligaments into two or more or less independent front and rear parts. The cavities are lined by a single layer of flat cells, which secrete a slimy synovia and are differentiated from connective tissue by the demands of use. Under the synovial layer lies an intricate network of lymph spaces, which, however, seems to have no direct connection with the joint cavity. As long as the synovial endothelium is intact, one may fill the

joint to bursting without injecting the peri-articular lymph channels.

When injured, the synovial endothelia tend to revert to a primary form; they turn into granulation tissue, lose their specific character, and end as scar, as adhesions or fibrous bands. Years of experience have taught us to respect the endothelium of the belly, to depend upon its power of combating bacteria when uninjured, to try to preserve its specific nature, and not to cause it to degenerate by rubbing it with gauze or destroying it with chemicals. Our experience with joints has not been sufficiently general to make these principles as widely known in the treatment of infectious arthritides as they are in peritonitis.

Payr¹ has made a classification of the sources of joint infection that is of much practical value. He says that bacteria may reach the joint from four sources: (1) directly from without, being carried in by a bullet or entering through an open wound; (2) through a fissure in an infected bone or through a communicating bursa; (3)—in exceedingly rare instances—through the encroachment of an infectious process from the neighboring tissues; and finally (4) through the blood stream, as in the metastatic joints of pyemia.

It is important to keep apart each of these four sources of infection,—the direct, by means

of a missile; the indirect, through a fissure in a bone; the secondary, by spread of a neighboring suppuration, and the metastatic. It is important, not only theoretically, but practically.

Each kind of infection has its rational mode of treatment. If the bacteria are carried in by a foreign body, obviously the thing to do is to get the offending invader out as soon as possible. If it is removed early, before the lining of the joint and the capsule have been irreparably damaged, recovery with complete restoration of function may be expected.

If the invasion has its source in an osteomyelitis which pours pus into the joint through an infected fissure in the bone, or if the pus arises in an infected bursa, it is useless to hope for improvement as long as the primary focus is active. If the original infection lies in a bursa, merely opening it may allow the joint to heal without attacking the main cavity at all. If, however, the pus comes from a fissure in the bone, if there is necrotic marrow, or sequestra that act like a foreign body and keep up a continually active bacterial focus, it is hopeless to strive to preserve the joint. The best we can do is to remove the dead bone, or if a manifestly infected joint fracture comes into our hands early, to forestall a certain sepsis by a thorough primary revision, removing loose fragments of bone and foreign bodies, and beginning antiseptic treatment before sepsis and deeper invasion of the tissues have begun.

The third source of infection,—from a neighboring focus of suppuration, is exceedingly rare. There are pus joints, it is true, where it seems as though a secondary indirect infection must have taken place; where a man shot through the thigh, three or four inches above the knee, for instance, develops a purulent joint. Careful scrutiny of the wound channel, however, will usually show that the connection is a direct one—will reveal a fragment of shell lodged in the suprapatellar bursa or some other communicating cavity. Opposite observations are certainly frequent and astonishing enough. One may see enormous streptococcic abscesses reaching from the ankle half way up the thigh, or have the leg rotten through and through with a gas phlegmon and yet see the knee uninvolved, with no exudate and with perfectly clean joint surfaces and synovia. Osteomyelitic foci in the epiphyses of the tibia may slowly reach the knee or ankle, pieces of disintegrating cartilage

may even come out of a cavity in the end of the bone marrow and thus prove the involvement of the joint, and yet the suppuration may not spread further and the man may remain fever-free. Such joints are usually the seat of a subacute arthritis with much atrophy; the capsule is more or less swollen, boggy and contracted, and the joints remain stiff; they may finally ankylose completely, but active joint suppuration and sepsis scarcely occur. Active suppuration and sepsis scarcely occur *if* the joint is left alone. The greatest danger to the joint lies, not in the nearby focus, but in the hands of the surgeon who opens it under a mistaken diagnosis. If the men are seen early, and if one surgeon has them in hand from the beginning, it is usually not hard to say whether there is pus in the joint or not. But if, as in military practice, the wounded have passed through three or four hands, it may be impossible to say, when they come to you, whether the pus lies in the joint or outside. The only thing to do is to wait and see. One certainly runs much less risk of spoiling a joint that lies near a suppurating wound by waiting and watching it for three or four days, than by opening it too soon and finding nothing but the pus that one's own hands have wiped into it. One cannot emphasize too strongly the necessity for distinguishing between pus *in* a joint and pus *around* it. To open it unless the pus is *in* it is to ruin it forever.

The last form of infected joint is the metastatically infected one. If there is pus, incise and let it out; if in doubt, the aspirating needle will tell.

The four sources of infection and their treatment, then, are: (1) Direct infection from without; remove the offending material as soon as possible. (2) Indirect infection from communicating bursae or joint fractures; treat the primary focus, resect if necessary, resect prophylactically if manifestly infected. (3) Secondary indirect infection from neighboring abscesses—rare; leave the joint alone unless sure it is involved. (4) Metastatic infection; evacuate the pus.

All of these infections, whatever their origin, may at one time or another lead to similar pathological and clinical lesions. Of these, there are two distinct groups,—distinct in their course, in their severity, and in their after-effects. The first is the suppurative form, without deep involvement of the joint capsule; the

second is the drier form, the capsule phlegmon. It is important to keep them apart. The whole future of the joint, whether it is to be movable or permanently stiff and crippled, depends upon whether the capsule is deeply involved or not,—whether the bacteria are merely in the joint cavity or whether they have wandered out of the cavity and infiltrated the capsule. If the deep capsule is unaffected, the nature of the exudate is not of decisive moment—whether serous, bloody, fibro-serous or frankly purulent, the damage is capable of complete repair. If, however, the synovial lining is destroyed, if the capsule is swollen and inflamed, the joint is gone. The swelling will close the fine vessels that pass from the capsule into the ends of the bone, the bone will atrophy, granulations will sprout from the synovia, cover and erode the cartilage, adhesions will form, and the joint, if it does not ankylose altogether, will be permanently roughened and stiff. These two forms, then—the exudative and the phlegmonous—should be recognized and kept apart.

The exudate may consist of turbid or bloody serum and fibrin or of frank pus—empyema. It first distends the loose para-articular bursae. The joint is swollen, not cylindrically, but irregularly. The contours of the distended and fluctuating bursae stand out. The adjacent soft parts are but little inflamed—neither hot, tender nor painful. The joint may even be moved to some degree without pain. The temperature is usually high, but there may be remissions in the morning. There may be sweats, sometimes even chills. The white count is increased. The swelling alone may be the distinguishing feature,—the other symptoms moderate or absent. The irregular, fluctuating distention of the para-articular bursae and the aspirating needle make diagnosis certain. One must be on the lookout for these joints. Septic patients may not complain of them nor speak of them spontaneously—they are not painful enough.

If pus is found, the cause—a foreign body—should be removed and the pus drained off. The joint, however, should not be left open. We have learned that if the cause of a peritonitis is removed, the belly will usually take care of itself; that to leave it wide open in the hope of draining it is not only unnecessary but harmful, and that the best way to keep the peritoneal endothelium intact, to keep it from forming adhesions, is to let it alone. So with a joint. If it is kept open, exposed to the air or covered with

gauze, both synovia and cartilage are lost. The synovia turns into granulation tissue and the joint obliterates. Therefore, let the pus out through a small stab, just large enough for a drain, keep the cavity intact and remove the drain early, as soon as the pus disappears. If the joint is opened wide to extract a foreign body, the synovia should be sewn up loosely, and a smooth, non-irritating drain, preferably a glass tube, introduced. It is interesting to watch the drain. The day after operation, or perhaps in 48 hours, it will contain, not pus, but clear serum or a transparent yellow coagulum. This is the time to take the drain out. The man may still have fever and the joint may be full of fluid again the next day. The fluid does not matter; it does no harm but good. Distention will keep the capsule from shrinking and adhering. In the course of three or four days the temperature will begin to fall and the swelling recede. If it continues, the joint should be aspirated to see whether it does not contain serum instead of pus before it is reopened. One or two further aspirations, or perhaps another 24 hours of drainage through a stab wound may be necessary for the exudate to disappear permanently.

The joint may be irrigated through the drainage tube. Payr recommends leaving 10 cc. of camphor-phenol in the cavity; others² have gotten much benefit from Carrel-Dakin's solution. This is probably the best antiseptic to use; it not only sterilizes the joint, but dissolves fibrin and coagula. If but an isolated part of the cavity is involved, irrigations or injections should not be practised; in any case they should be gentle. Pressure may force bacteria into bursae or pouches originally uninfected.

Several days of fever need cause no alarm. Temperatures of 102-103° quite commonly follow arthrotomy.

Blood in an uninfected joint remains fluid a long time, for weeks,—an infected hemarthros contains small shreds and clots. A smear will show an undue proportion of leucocytes, but a culture may be necessary to determine finally the presence of bacteria. If the exudate is bloody, antiseptics like camphor-phenol, which clot blood, should not be used.

We stated that a joint consisted of a series of communicating cavities rather than of a single large one, and that even the large main cavity was divided into separate chambers. Some of these may be infected, while others are not. A

knee may contain pus under the quadriceps and in front of the crucial ligaments and clear serum behind. The elbow may have pus in the olecranon fossa and be unaffected in front. In these cases of isolated empyema it is important to know where the pus lies, to see where the swelling is, and which part of the joint fluctuates. If the pus lies in front, drain the front. If behind, drain the back. If it lies in a bursa, drain it. If in several, drain each one. But do not drain the unaffected portion of the joint. Let it alone and, above all, do not pull a drain through from the purulent into the uninfected part. The usual text-book incisions are rarely indicated. Through-and-through drainage across the knee above the patella, for instance, is quite useless. Early cases had better not be treated with traction—the distention of the joint may let pus flow from an infected pocket into healthy parts.

Until the swelling has gone and the temperature fallen, the joint should be kept quiet in a splint and treated by some method of hyperemia, preferably baking. Passive congestion is not so good. As soon as temperature and fluid have disappeared the splint should be removed and the man encouraged to move his limb; actively at first, then with semi-active exercises in some form of swinging apparatus.

All energy should be bent upon prompt treatment. In this exudative form of arthritis we can save not only life and limb, but the joint, provided we get at it before the capsule is invaded.

In the second form of infectious arthritis, in the capsular phlegmon, we must content ourselves with more modest aims. We cannot hope for the joint. We must strive for life and limb. A phlegmon presents a picture quite different to that of an empyema. Tenderness and pain are marked, the joint is held anxiously stiff. The slightest movement is painful, but the swelling is less than in empyema. The contours of the joint can no longer be made out; the spindle-shaped swelling reaches all the way around the limb. There is periarticular edema above and below the joint, but the borders of the capsule may sometimes be felt through it as a hard ridge. The skin is thin and hot, red or cyanotic. The muscles are much wasted. If the phlegmon is of long standing, the region may be riddled with sinuses that pour out a thin watery pus and flakes of fibrin. From their mouths protrude thick, glairy granulations, covered with a firm

membrane that makes them bleed when it is pulled off. The membrane is characteristic. A sinus that discharges much thin pus and whose granulations are covered with membrane is almost pathognomonic to lead into a joint. The joint itself is usually shrunken and empty rather than distended with fluid. The pus comes mainly from neighboring bursae or intermuscular abscesses. The capsule is swollen, boggy, reddened, and bleeds easily. The joint cavity is obliterated by organized fibrinous deposits. The cartilage is eroded; its edges are overgrown by synovial granulations. It may be undermined by them in places and exfoliate in green or yellow flakes, exposing the granulations that cover the end of the bone. If the phlegmon is the result of an osteomyelitis that has found its way into the joint through a fracture, the epiphyseal marrow may be full of pus, the ends of the bone black or green, dry and necrotic, and the blood sinuses plugged with dark clots or pus. The x-ray will show decalcified bone; the corners of the epiphyses will be irregular and hazy.

As time goes on the bursae and weakest parts of the capsule give way, and the pus pours out into certain typical spaces between the muscles. It may appear at so great a distance from the wound that one may never think of its articular origin unless one has learned to recognize these long tubular pus sacs. A suprapatellar bursa, for instance, breaks through into the loose space under the quadriceps, makes a huge abscess along the front of the thigh, and appears at the inside of the groin, or below and behind the trochanter. The bursae below the heads of the gastrocnemius break through into the deep muscular planes of the leg and reach the skin at the side of the Achilles tendon, above the ankle. The shoulder joint breaks through along the biceps tendon and appears at the inner side of the biceps, above the elbow. Waxy edema or brawny infiltration of a limb, without much tenderness, should excite suspicion of this form of abscess.

The temperature in capsular phlegmon is continuous, but it need not be high. Remissions alternate with exacerbations as new bursae are involved and fresh abscesses appear. The white count is moderately increased—10,000 to 18,000. Loss of sleep and sepsis cause the men to fall away rapidly; drenching sweats, septic diarrheas, amyloid, apathy and coma signalize the danger of death. In the hip joint the temperature may be continuously high, as in typhoid, or intermit-

FIG. 1.—Obliteration of tibio-tarsal joint after osteomyelitis of tibial epiphysis. Joint remained free of pus.

tent, as in malaria; the depth of the joint conceals it from examination so that mistakes for one of these diseases are by no means impossible.

As the man recovers, the joints stiffen and contract in typical positions of flexion. The swelling recedes, the sinuses close, but muscular atrophy remains, and all attempts at restoring motion by passive or active exercises are unavailing.

Treatment offers great difficulty. One will never cease to have patients who, whether they recover or succumb, one would wish to have treated otherwise.

A certain number of phlegmons may be prevented; all of them, perhaps, if the men were treated immediately after injury by sacrificing numerous joints. The indications in the two extremes are simple; empyemas—the lighter cases, may be prevented from going on to a phlegmonous course by proper early treatment as outlined above; in the graver cases—obviously infected joint fractures—the joint should be sacrificed from the beginning. It should be opened widely as soon as practicable, the

path of the projectile, shreds of fascia and blood-soaked tissues excised and foreign bodies extracted. The ends of the bone should be inspected; loose fragments and those attached by slight shreds of periosteum, whose life appears improbable, should be removed. The joints will be stiff at best. If we leave bone that is likely to die we stand a good chance of having a supuration that will cost life or limb. The joint should be filled with an antiseptic or irrigated with the Carrel-Dakin solution. If this solution is not used multiple drains should be led out of the dependent parts, *i.e.*, the back, and the capsule loosely closed in front. The limb should be immobilized in a splint or fenestrated cast and kept quiet until danger of sepsis has passed. It should not be disturbed for unnecessary dressings—too many are as bad or worse than too few.

The cases which are hard to know how to treat are those lying between these extremes, but where the phlegmon is well established. We are beset by difficulties at every turn. If we wait too long we lose the man. If we amputate too early we lose a limb which might, perhaps, have been saved. If we immobilize it is difficult to watch the joint. If we do not, sepsis continues. We are between many fires; certain general principles of treatment, however, apply everywhere.

A phlegmonous joint should be securely fixed in slight flexion by a well-fitting splint, or by a circular plaster cast fenestrated widely enough

FIG. 2.—Changing cast for injury of knee.

for the limb to be watched and kept immovable during dressings. An interrupted bridge cast, which leaves the joint free all around, is not as good as a circular one. The joint sags down between the two halves of the cast. If the plaster or splint has to be removed, the joint should be kept quiet by slings suspended from a frame or slipped over the shoulders of an assistant, so that the patient may not move it while under the

Treatment varies with different joints. Sepsis in the upper extremity is not so dangerous to life or limb as sepsis in the hip, knee or ankle; on the other hand, resection in the arm is not nearly so crippling as in the leg, and usually results in mobile joints.

The shoulder, unless the bone is splintered, mostly heals without resection, often with a good movable joint. It should be treated in ab-

FIG. 3.—Gunshot wound of elbow; a, b, x-rays five weeks after injury; c, extension and supination; d, flexion and pronation after conservative treatment.

anesthetic. A glance at Fig. 2 will explain. If provision is not made for keeping the joint quiet during operation, the patient's struggles may so spread infection that the operation may do more harm than good. Every movement is followed by an exacerbation of fever and local symptoms. Unnecessary transportation to and from bed should be avoided. Small abscesses may be incised under local anesthesia without removing the splint or taking the man from bed.

duction so that pus may not sink down along the biceps, and so that if the joint stiffens, the man may still abduct by using his scapula.

The elbow is so complex a joint, and suppuration in it is so tedious that it is best to resect moderately early, say in two to four weeks, if fever and phlegmon have not disappeared by then. If the radius is broken at the neck the head becomes necrotic and sequesters, but ordinarily the classical resection of both radius and

FIG. 4.—Late resection of elbow. "A", head of radius sequestered and eroded.

ulna is better than a hemi-resection of one or the other bone, even though only one of them be affected. In joint fractures the elbow almost always stiffens, so that a later resection or some form of arthroplasty is necessary. An early resection will save this.

Sepsis in the wrist and carpus is obstinate. The hand long remains swollen and tender, and abscesses appear above and below the wrist, especially along the deep fascial planes of the forearm. Suppuration in the carpal joints is often followed by pyemia. As long as the man's gen-

eral condition seems good one may wait, keep the wrist in dorsal extension in a splint, bathe the hand in hot brine, bake it and see whether the carpals become necrotic before excising them. The wrist stiffens, but the men have a fair grip if it stiffens in dorsal extension; even excision of the carpals leaves a useful hand. If excision is done early enough, exarticulation at the wrist will rarely be necessary.

Capsular phlegmons in the lower extremity and their sequels give some of the gravest cases in military practice. Septic hips make long and high fever. One recognizes them mainly by tenderness over the head of the femur at Poupert's

FIG. 5b.—Gunshot wound. Fracture of head of femur. Rifle bullet found in head of femur. Septic hip joint. Necrotic and putrid head removed 11 days after injury. Recovery.

FIG. 5a.—X-ray 11 days after injury.

ligament. Mobility may be not much impaired. If recognized, they need not be so much dreaded as suppuration in the knee. A broken head of the femur should be removed early; it will die and rot and have to come out anyway. The joints should be left well open after resection and drained to the rear and outside. Jones' long frame and adhesive plaster traction will hold the leg quiet. A plaster spica is almost impossible to keep from getting foul and the men develop bed-sores under it. The joint may contain very little pus. In one case I suspected suppuration from the continued fever and joint tenderness. I opened the hip, but could find nothing abnormal. I closed the joint. The man

FIG. 6.—Suspended and fenestrated cast for wound of knee.

got well and kept a perfectly movable hip. The joint must have been infected in spite of the apparent absence of pus.

The most difficult joint phlegmons to treat are those of the knee. I think one of two courses is best: The first is to treat them conservatively, immobilizing with Thomas splints or widely fenestrated plaster casts in a few degrees of flexion, suspending the whole splint from a frame, baking the joints and watching them daily—not only the knees but the men—incising abscesses as they appear, irrigating with Dakin's solution, and continuing this treatment until the knee is quiescent. If, however, the knee gets worse and the general condition becomes alarming, one will have to exarticulate or do a Gritti operation. In the Gritti the patella should be left in the anterior flap and wired into place later, after the wound is clean. The disadvantage of this conservative method is that by the time one decides to exarticulate, multiple incisions of abscesses have so spoiled the soft parts that one gets but a poor stump. The other plan is to do an early resection with posterior drainage or with the Carrel-Dakin irrigations. This procedure is the one to follow in shell wounds complicated by fracture. To be of use, the operation should be done before para-articular abscesses have burrowed their way up and down the limb. Once the pus has traveled away from the joint, resecting the ends of the bone will not save the leg.

I can heartily warn you against the other method of treatment—the method of opening

the joint wide, cutting across the crucial ligaments and splitting open all the bursae. It looks beautiful when you have done operating; the joint is wide open, everything is exposed, and if you put on a few pounds of traction you may separate the femur and the tibia for an inch or more, but—the men all lose their legs. The back of the tibia becomes rough and erodes the popliteal artery, and it bleeds, or abscesses burrow up and down, and you have to amputate any way.

There remain, of course, early amputations and exarticulations. Military exigencies may force one to them, but they should not be done if there is a chance of watching the man while his limb is in danger.

Suppuration in the ankle and the tarsals is as tedious and painful as it is in the hand, and more dangerous. Suppuration in the foot seems prone to make all sorts of unforeseen complications—thrombophlebitis and encephalitis and pulmonary infarcts, and other rare sources of anxiety. Apart from these complications, if one

FIG. 7.—Loose bone removed at prophylactic resection of knee four days after injury.

FIG. 8.—Capsular phlegmon of knee with intermuscular abscesses; joint opened widely. Open wound treatment. Subsequently erosion of popliteal artery, fulminating hemorrhage, exarticulation, finally recovery.

keeps the foot quiet in a splint or cast, bakes it or soaks it in hot brine, it tends to get well. Para-articular abscesses appear, or rather, conceal themselves in the sole, where the thick skin and fascia hide them, unless one watches closely. They should be incised from the inner side of the foot, so that the man may not tread on the scar afterwards. If suppuration does not cease, the astragalus may be resected with benefit and without doing much harm to the foot. One certainly should not amputate without trying both this and all other kinds of atypical tarsal resections. Any kind of a misshapen elephantine foot, if it is sound and has good skin and fat on the sole, is better to walk on than no foot at all.

Now we have not considered wounds, but traumatic infections of the joints. Are there no joint wounds that are not infected? Yes, some

perforating gunshot wounds heal perfectly aseptically under conservative treatment. Some even with foreign bodies, especially rifle bullets, may be treated as though they were aseptic if one gets them early. As soon as circumstances permit of aseptic operating, the wound of the soft parts should be excised, the joint opened, the foreign body extracted, and the joint closed. If asepsis is doubtful the synovia should be closed, but the skin and other soft parts left open or sewed together quite loosely; or the joint itself may be drained with a glass tube for 24 hours. A few days of fever, 102° or so, either before or after operation, do not spell disaster. All these men have temperatures which often drop if the limb is immobilized and the patient left alone.

Aseptic cases, like empyemas, should be allowed

FIG. 9.—Shell wound of knee; fracture of patella. Synovia closed by suture. Recovery of function four weeks after injury

FIG. 10.—Piece of steel imbedded in posterior part of external semilunar for three years. Chronic arthritis; lipping of upper edge of patella and upper edge of rear of femoral condyles.

active motion as soon after operations as fever, pain and tenderness permit. Capsular phlegmons should be kept quiet. Endeavors to re-establish motion by manipulation are quite futile and cause the patients needless pain.

Foreign bodies should be removed, even if they heal into a joint aseptically. It is true that, take it all in all, more harm than good has been done in the present war by hunting for bullets, and that extraction operations have caused much needless mutilation and suffering. A joint, however, is no place for a foreign body. An aseptic piece of steel will wear off the cartilage when the joint is moved, cause roughening and incongruity of the articular surfaces, and end in a chronic deforming arthritis. A case in point: A man who had injured his knee by a blow from a steel wedge three years previously had had recurrent attacks of inflammation and hydrops. Flexion and extension were considerably impaired. An x-ray showed a chip of steel in the posterior capsule, although the man had no knowledge that anything had entered the knee at the time of injury. On opening the knee a spot bare of cartilage was seen on the inner femoral condyle. All about it were signs of deforming arthritis. In places the cartilage was peeling off; in others it had peeled off and flakes lay free in the joint. The steel chip was

FIG. 11a.—Old gunshot wound of knee. Subsynovial deposits of lead. Severe deforming arthritis.

FIG. 11b.—Same case as 11a. X-ray of resected femoral and tibial condyles. Lead in synovial tissues. Erosion of bone by synovial granulations.

FIG. 11c.—Same case as 11a. X-rays of resected femoral and tibial condyles. Lead in synovial tissues. Erosion of bone by synovial granulations.

FIG. 11*d*.—Same case as above. Sections of subsynovia, showing lead deposits.

FIG. 11*e*.—Same case as above. Sections of subsynovia, showing lead deposits.

FIG. 12.—Localization of foreign body by needle method of fluoroscopy. (Holzknecht) *a.* Shell fragment in semilunar cartilage. *b.* First needle has missed fragment. *c.* Second needle has missed fragment. *d.* X-ray perpendicular to axis of needle. *e.* Shell fragment removed. Joint space outlined.

found imbedded posteriorly in the external semilunar and removed. The joint, however, was so changed that it will never regain free motion.

Lead (from pistol bullets or shrapnel) is particularly dangerous. It seems partly to dissolve in the synovial exudate and to settle as a grayish-white deposit under the capsule. It may be seen in sections as black grains lying in the subsynovial tissues. Several men who had carried aseptic pistol bullets in their joints for a number of years have developed an uncommonly severe hypertrophic arthritis.

Extraction operations should not be entered upon lightly. They should be done with competent assistance, under good asepsis and with a good x-ray apparatus. The foreign body may be very difficult to find. It often traverses the joint and is stopped by hard, thick capsule or cartilage. If it lies in tissue of this consistency the finger cannot feel it even when a millimeter or two from it. One may approach very closely with the scalpel and yet miss finding it. Radiographic plates by the MacKenzie-Davidson, or one or another localizing method, do not help nearly as much as the fluoroscopic screen. If the hospital is equipped for aseptic fluoroscopic operations the body may be extracted directly under the screen; if not, a needle may be introduced down onto it under the fluoroscope after the method of Holzkecht and the projectile removed by cutting down on the needle, which is left *in situ* as a guide.

Removal may be particularly difficult in the knee, where foreign bodies are quite typically found buried in the posterior attachment of the semilunar cartilage. If a lateral x-ray plate (like Figs. 10 and 12a) shows the body just above the rear edge of the top of the tibia, removal should not be attempted without the aid of a screen. A body imbedded in the cartilage may be impossible to locate without it.

The best time for removal is early—within the first twelve hours after injury. Later, if the joint enters an acute inflammatory stage, it should, if the body is small, be immobilized and treated conservatively with baking and compresses until the inflammation has subsided. The trauma of extraction will set up a strong reaction during the acute stage. Acute generalized inflammation will often quiet down after immobilization; four or five days after the fever falls, the foreign body may be removed and the joint closed loosely over a glass drain.

CONCLUSIONS.

1. Sources of joint infection are:
 - (a). Direct, from without.
 - (b). Indirect, from a communicating suppuration.
 - (c). Secondary, by extension of a neighboring process.
 - (d). Metastatic.
2. Infections may be:
 - (a). Empyematous—prompt treatment will save the joint.
 - (b). Phlegmonous—the joint will stiffen.
3. Leaving a joint wide open is harmful. Treatment should strive to preserve the synovia.
4. Foreign bodies, especially lead, induce deforming arthritis, and should be removed.

REFERENCES.

- ¹ Münch. med. Wehst., 1915, Nos. 37, 38, 39, pp. 1241, 1282, 1321.
- ² Campbell and Wolfenden: Lancet, Vol. xciii, p. 185.

TRENCH SANITATION.*

BY CAPTAIN C. E. BURT, M.R.C.,

First Lieutenant, Medical Corps, U.S.A.; Surgeon,
183rd Infantry Brigade, Camp Grant, Illinois.

IN the consideration of a subject so important, it is to be regretted that more literature is not available. Physicians who have the subject from first-hand information are all more or less engaged in the work of the war, and have little time to write. The Advisory Committee of National Defense is taking up these subjects most essential to us, and is having manuals printed, revised from the English and French, which will soon be in the hands of the members of the Medical Corps. This paper will deal with the most important subjects, with a short description of the most particular methods that the peculiar environment abroad has developed in modern trench warfare, which I have collected from those who have written, and conditions which I have personally observed as battalion surgeon of the trenches in the training camp. I assure you, however, that I make no claim for personal originality, but give due credit to those who have been able to write.

You are all much better acquainted with trench conditions from your readings and from the lectures by the Allied officers than I am, so that you can easily draw upon your vivid imaginations to realize the difficulties encountered in dealing with this most important subject.

* Read at a meeting of brigade officers at the request of the commanding general.