fact that he found the vessels of the cord rather overfilled with blood, he recommends the trial in such cases of some preparation of ergot.
ALLEN.

129. ZUR LEHRE DER SPINALEN NEURITISCHEN MUSKELATROPHIE (ATROPHIA MUSCULARIS PROGRESSIVA SPINALIS NEURITICA, BERNHARDT) (PROGRESSIVEN NEUROTISCHEN ODER NEURALEN MUSKELATROPHIE HOFMANN) (Spinal Neurotic Muscular Atrophy). E. Siemerling (Archiv. f. Pyschiatric, 31, 1898, 1899, p. 105).

Siemerling contributes a complete history, with report of microscopical findings, in a case of this interesting and rare form of muscular atrophy. The history is of a young man, 20 years of age, with no especial family hereditary influences. Up to the age of 5 years he developed normally. At this time there commenced an atrophy of the small muscles of the lower extremities, and two years later the hands commenced to be involved. The atrophy progressed until at the age of 13 he was unable to walk. He later developed hypochondriasis. At the time of examination his intelligence was fair; there was some immobility of the pupils to light, no changes in the fundus, and extreme atrophy of the muscles of all four extremities. Knee-jerks were absent. The right leg was in contracture at the knee-joint. No club foot. Complete paralysis in the lower extremities, less in the upper. Fibrillary twitchings of the intercostal muscles. Loss of electrical contractility for both galvanic and faradic currents. Psychically there was marked hypochondriacal depression, with delusions. The patient died in collapse.

The microscopical analysis showed degeneration of the posterior and lateral columns, especially in the dorsal and lumbar regions. Atrophy of the anterior horn cells, Clarke's columns and the anterior roots. The posterior roots were intact in their extramedullary portion, and in some parts of their intramedullary course. Simple degeneration of the sensory and mixed nerves, with atrophy of the musculature. The résumé of the microscopical findings was as follows: Disease of posterior columns, most marked in the lower dorsal regions. In the upper dorsal and cervical regions almost total degeneration of the central part of Goll's column. In the columns of Burdach the degeneration becomes markedly less as the upper cervical region is reached. There is degeneration of the lateral columns posteriorly and partly of the anterolateral columns, the lesions being most marked in the lower dorsal and lumbar regions and decreasing upwards. The cerebral ganglion cells showed no alterations. The muscles of the lower extremities were in a condition of extreme degeneration, the gastrocnemius showing advanced fatty changes. All of the peripheral nerves examined, brachial, radial, median, ulnar, sciatic, crural, peroneal, and saphenous nerves, were markedly degenerated.

The author further gives a critical digest of the various theories accounting for the disease, and, basing his opinion on this one case, believes that the disease is one of intramedullary origin, the posterior roots being affected after their entrance into the spinal cord. He does not exclude, however, a peripheral origin for the disease.

J. L. LIPP.

- 130.—PARALYSIE OBSTETRICALE CONSECUTIVE A DES TRACIONS ASYNCLI-
TIQUES SUR LA TÊTE (Obstetrical Paralysis from Manual Traction
on the Head). Planchu (Lyon médical, 88, 1898, p. 545).

Plexus paralysis due to difficult parturition generally occurs in breech presentations, or when severe traction is made with forceps. In the case reported the presentation was by the vertex and the head

was born spontaneously, but the shoulders were arrested. The nurse seized the head with her hands and with all her strength exerted traction upward—toward the right shoulder of the child. Immediately after birth paralysis of the left arm was noticed, the superior (first) cord of the plexus being the most severely injured. PATRICK.

131. CASUISTISCHE BEITRÄGE ZUR HIRNCHIRURGIE UND HIRNLOCALISATION, etc. (Clinical Contributions to Cerebral Surgery and Cerebral Localization). Ludwig Mann (*Monatsschrift für Psychiatrie und Neurologie*, Vol. 4, No. 5, p. 369).

Mann reports a case in which two focal symptoms were observed after a cortical injury, viz., loss of sensation in the left thumb and index finger, and loss of power to sing. At the necropsy a cyst was found involving the second right frontal gyrus, and a small portion of the anterior central gyrus in the lower part of its middle third. It was not certain whether the cyst was due to the original injury or to operation. The loss of sensation in the thumb and index finger was explained by the involvement of the lower part of the arm center, but the absence of paresis in thumb and finger was striking. This is said to be the only case with necropsy in which amusia was not combined with aphasia. Mann believes that possibly a lesion of the second frontal gyrus may cause amusia. In his case the lesion was on the right side of the brain, and the patient was right-handed. SPILLER.

132. UN CAS DE SCLÉROSE EN PLAQUES A TREMBLEMENT UNILATÉRAL (A Case of Multiple Sclerosis with One-sided Tremor). P. Remlinger (*Rev. de Médic.*, Vol. 19, 1899, p. 244).

The case reported showed the following cardinal symptoms: There was an intention tremor of the upper and lower extremities, limited to the right side; exaggeration of the tendon reflexes, more marked on the right side; gait markedly spastic; tremor of the lower lips; diminution in sight, due to beginning optic nerve atrophy; slow speech, with monotonous and scanning cadence. These symptoms had been developing for the past six years, during which time the limitation of the tremor to the right side was remarkable, being so constant.

JELLIFFE.

133. UN CAS DE MENINGITE CEREBRO-SPINALE SIMULANT LE TÉTANOS (A Case of Cerebro-Spinal Meningitis Simulating Tetanus). La Presse Médicale, 24 Décembre, 1898 (Leroux & Viollet).

The patient, a man of 40, had noticed some stiffness in the lumbar region three days before admission, followed quickly by paralysis of the legs and difficulty in taking nourishment. When admitted to the hospital he was rigidly extended, the back being some distance above the surface of the bed. The neck was very stiff, and there was moderate trismus. From time to time he had paroxysms of pain, causing him to cry out. The temperature was 40° C., the pulse 120, and the respirations 40. The latter were almost exclusively diaphragmatic. The pupils were perfectly normal. A diagnosis of tetanus was made. Roux and Martin, of the Pasteur Institute, saw the case in consultation and pronounced it meningitis. The patient continued to grow worse and died on the second day. At the autopsy small collections of pus were found just back of the frontal convolutions, and there was intense congestion of the arachnoid plexus to the plexus of the fourth ventricle. Cultures made by Martin showed the presence of the diplococcus of Friedländer. The authors also report a case with very similar symptoms, the patient having the slight contraction of the extensors of the vertebral column, slight rigidity of the lower extrem-