tinuance of a mucus discharge after diphtheria or the exanthemata. Physicians, on the other hand, should not pass these complaints lightly by ignoring them, or directing in an indifferent manner some simple and likely inefficient remedy.

The management of hypertrophies at this early stage is as simple as the results are happy. After a careful examination the nares and upper pharynx should be cleansed of mucus by an antiseptic and detergent spray, *e. g.*, Dobell's solution, Seiler's tablet, Listerin in diluted solution, etc., and then applications made to the affected mucous membrane of an astringent or alterant, depending on the condition of the parts. Should we find simply an increased secretion with no appreciable alteration in the turbinates, tonsils, and surrounding tissues, it is likely we will accomplish all that could be desired by directing spraying of the nares thoroughly by one or other of the solutions mentioned twice daily and continued faithfully until the membrane presents a normal appearance and the secretion ceases.

In cases still farther advanced in which we find the turbinates quite red and somewhat swollen; the pharyngeal tonsil secreting more mucus than normal, and rather large; the faucial tonsil red and somewhat hypertrophied, we will find it advantageous to use some application to diminish the inflammatory condition. These applications may be made quite painless by spraying the parts with a 5 per cent. solution of cocain or introducing a cotton tampon saturated in the same solution. In the nares a solution which has gained favor in the various clinics devoted to these diseases is that of iodine. The strength of the solution may be graded by the severity or obstinacy of the case. It is well to begin with a weak solution, increasing the strength as the necessities of the case demand. In some cases which resist this plan of treatment a touch of the fused bead of chromic acid along the turbinates will often accomplish the desired object; this latter plan should be immediately followed by the antiseptic spray to dilute and limit the destructive power of the escharotic. It is merely necessary in these early cases to touch the mucous membrane lightly to produce a superficial eschar; the resulting slough and subsequent contraction will so squeeze the blood vessels as to bring about the desired diminution in the size of the tissues.

The pharyngeal tonsil also in many cases will respond satisfactorily to applications of iodine on a pledget of cotton carried by an applicator so curved as to pass readily behind the soft palate and up to the vault. Glycerole of tannin has been found to act very happily, though perhaps a little more unpleasant in its effects. The application of chromic acid (fused) to a pharyngeal tonsil should not be advised, though undoubtedly as useful as in the nose, unless the operator is steady and familiar with making applications by the aid of the rhinoscopic mirror.

It would be hardly necessary to refer to the treatment of catarrhal and slight hypertrophic conditions of the faucial tonsils the result of the causes named, were it not that here, in spite of the ready accessibility of the part, they are subject to neglect. The same process of cleansing is as necessary as in the other hypertrophies, after which an application of a solution of 60 gr. to the drachm of nitrate of silver, glycerole of tannin, or in stubborn cases producing a superficial slough by fused chromic acid, will probably answer every purpose.

In the more advanced cases or in those which fail to respond to the above-outlined plan of treatment and a steady increase of the hypertrophied tissues occurs, there is but one plan which should be carried out at as early a date as possible, namely a destruction of the hypertrophied tissue. The various plans to accomplish this object will not be touched upon, but can be readily learned by consulting any of the modern treatises devoted to the treatment of diseases of the nose and throat.

In recapitulation the object of this paper has been to call attention to the following points:

1. The importance and desirability of early attention to diseases of the upper respiratory tract consisting of the nose and naso-pharynx.

2. The production of alteration in nutrition and possibly pulmonary and bronchial troubles by hypertrophic conditions of the upper respiratory tract.

3. The etiology of these states referring to the influence of environment and especial mention of diphtheria and the exanthemata as prominent factors in this category.

4. The prevalence of these conditions.

5. Carelessness and indifference in both physician and patient as to the various catarrhal states which are believed to be the precursors of the diseases under consideration.

6. The necessity of early treatment and the power for good exercised by the family physician in these troublesome conditions.

7. The clinical picture presented by these unfortunates, laying especial stress on the alteration in facial expression and the distressing and annoying condition of the child during sleep.

TREATMENT OF DIPHTHERIA.

Read before the Wayne County Medical Society, 1894.

BY OSCAR S. ARMSTRONG, M.D.

DETROIT, MICH.

There is no disease with which we have to contend that requires better surroundings or greater care in its treatment to insure a successful outcome, than diphtheria. It is rightly dreaded as one of the most deadly foes of childhood, and even adults do not escape its attacks. It is a treacherous disease, and one or two members of a family may have it in a mild form, while in the remaining children it may present its most malignant characteristics. Nor does its treacherous character end here, for often when our patient seems to be doing nicely, with membrane entirely gone, pulse and temperature normal and everything progressing favorably, some of its dangerous sequelæ make their appearance.

The first record we have of diphtheria in America was the epidemic in Boston in the sixteenth century, since which time it has become completely domiciled. Originally the disease appeared in epidemic form only, but at the present time it has become essentially endemic in nearly all our large centers of population. Its ravages are not confined to the homes of the poor in overcrowded and filthy tenement houses, but extend to those of the wealthy in country and city. Until the last few years little was understood of its true nature, but pathologists by careful and patient investigation have discovered its cause. Klebs was the first to demonstrate the existence of a specific bacillus of diphtheria, and his conclusions were afterwards confirmed by Löffler. Experiments have proven that

the bacillus of Klebs-Löffler, when brought in contact with the mucous membrane or inoculated therein, will produce a true diphtheritic membrane with its accompanying constitutional symptoms. We are thus prepared at the present time to treat diphtheria from a more intelligent standpoint than ever before. Yet notwithstanding this fact, from statistics which I have been able to collect, the mortality is simply frightful, and if an epidemic prevailed in any of our cities in which the death rate was correspondingly so great, it would, I venture to assert, create consternation not only among the laity but the profession as well. It seems to me that we have become so accustomed to its pestilential presence that we accept its frightful mortality as something unavoidable, and it has ceased to alarm us.

To quote from an interesting paper in the last JOURNAL OF THE AMERICAN MEDICAL ASSOCIATION, by Dr. E. L. B. Godfrey, of Camden, N. J., on the "Treatment of Diphtheria," there were 2,624 cases of diphtheria reported to the Board of Health of Philadelphia from January 1 to October 28, 1893, with 750 deaths, making a death rate of 28.5 per cent. During the same period in Camden 250 cases were reported with 56 deaths, making a death rate of 25 per cent. During the year of 1892 there were reported to the Board of Health of Detroit 1,156 cases, with a death rate of about 26.5 per cent. For the present year I have been unable to secure the statistics, but Dr. McLeod, the Health Officer, assures me that the number of cases has decreased about 40 per cent. as compared with 1892.

It would seem apparent from the diverse opinions and methods of treatment that no specific has yet been discovered. The symptoms in the early stages are so closely allied to those following the contraction of a severe cold that many times parents are misled, and do not call their physician until the disease is thoroughly entrenched. Every family physician ought to instruct father and mother how to examine their child's throat, and whenever he (or she) complains of sore throat or gives evidence of it by symptoms such as languor, headache, fever, etc., it ought to be inspected and if inflamed or white spots show, the physician summoned at once to make a diagnosis and care for the case. In the autumn of 1891 I had under my care three cases of diphtheria in a family of five children all making favorable progress. One day when making my morning call I was asked by the mother to see a fourth child, a little girl 3 years of age, whom she said was not well. On examining the child's throat I found it literally full of diphtheritic deposits of a purple hue. Fauces, tonsils, uvula and soft palate were covered and the tissues necrosed, the offensive odor being almost unbearable. The countenance was suffused, eyes dull, stupor manifest, temperature 106 degrees F. and pulse so rapid it could not be counted. It was a case of diphtheria of the most malignant type. I gave a very unfavorable prognosis and asked for a consultation. During the afternoon I saw her again with my friend, Dr. Mulheron, a gentleman of experience in this class of cases. All we could do for our little patient availed nothing and she died the same evening, about twelve hours after I first saw her. Since that time I make it an invariable rule to see every child in the house in which I am treating a case of diphtheria, at each visit.

The period of incubation in diphtheria varies from

two or three days to a week. If the attack is mild the symptoms are at first hardly perceptible. There is usually slight chilliness followed by fever, thirst, headache, anorexia, furred tongue, more or less muscular soreness and a certain amount of rigidity of the muscles of the neck and throat. There are undoubtedly cases of the disease so mild that they are not recognized, and the little one associates with his playmates and conveys to them the virulent germ. In this manner many children acquire diphtheria, leaving us ignorant of its method of introduction. If the attack is malignant in character its advent will be marked by correspondingly severe symptoms. It does not follow, however, even in the worst forms that the patients will complain of sore throat. This one diagnostic point will often enable us to differentiate between it and tonsillitis, for in the latter the pain on deglutition is generally much greater.

A large and well-ventilated room should be chosen, and the carpet or rugs, upholstered furniture and curtains removed. The most favorable temperature for my patients has been 68 degrees F. The disease being at first local and afterwards constitutional requires both forms of treatment. During the past two years I have adopted a plan of treatment that has given me excellent results in *every case*. The first and most important thing for the physician to do is to get the good will and confidence of his little patient. When this is done half the battle is already won. When the little one is willing and anxious to do as you direct, you are enabled to conduct a case as satisfactorily with a child of 4 or 5 years of age as though he were an adult. On the contrary, if a spray or gargle has to be forced on the unwilling patient with nerves wrought up to the highest tension, medication, no matter how well considered, is of no benefit, and I believe is positively harmful. The patient should be put to bed and made to understand that the recumbent position is a necessity and that this rule must not be violated by any exertion of his own. When it is necessary to use the spray, gargle or swab, he should be supported in a half reclining position by the nurse. The rule applies with equal force when nourishment or medicine is given.

As a local application I have found nothing that approaches in value the peroxid of hydrogen. It may be used as a gargle or spray or held for a few moments in contact with the membranous deposit by means of a swab. I use it varying from 30 per cent to full strength, the latter only when it is applied directly to the exudate. Children use a 30 to 50 per cent. solution as a gargle or spray, without any discomfort or objection. When the tonsils are enlarged and painful, the throat is covered by a piece of soft flannel moistened with a hot mixture of equal parts of lard and turpentine, enveloped by a thick fold of wool cotton held in place by a bandage. If the throat is not painful the turpentine and lard are omitted.

The medicines most in vogue in past years for internal administration have been the tincture of the chlorid of iron, sulphate of quinin and chlorate of potassium, all of which are more or less disagreeable to the taste and require a great deal of persuasion to get the little sufferer to take them. During the past two years I have used the bichlorid of mercury internally, in all my cases of diphtheria, as a routine plan of treatment and with unvarying good results. I combine it with alcohol, glycerin, eucalyptol and

minute doses of arsenic, in a strength varying according to the age of the patient. It may be given in doses of one-fiftieth to one-sixtieth of a grain to a child of 3 years, every two hours. For the past year I have been using it as prepared by the Detroit Chemical Company, containing one-sixtieth of a grain to the drachm. The preparation is known as diphtherol, and is combined with pleasant aromatics so that children take it without the slightest difficulty. Due credit must be given to the alcohol which this remedy contains, for in a disease so depressing to the vital centers and in which the toxic symptoms are so pronounced, supporting and stimulating treatment ought to be pursued from its commencement. The effect of the treatment with the bichlorid is truly remarkable, and I have seen cases that I, at my first visit, deemed hopeless, improve so rapidly that in twenty-four hours the crisis was passed. It ought to be given freely until its constitutional symptoms appear, when the frequency of administration should be lessened. In very bad cases I have given 1 drachm of the diphtherol every hour for the first day, or until the constitutional effects in the way of uneasiness and pain in the abdomen were manifest. When the nasal passages are affected, I use the spray of peroxid of hydrogen solution with great comfort and relief to the patient. It ought to be used as often as every two hours.

Complications may arise requiring the use of quinin, digitalis, strychnin, etc., but they can in the majority of cases be administered in the form of a suppository. Since I have been using the plan of treatment here outlined, I have been fortunate in having laryngeal invasion in but one case. A possible intubation or tracheotomy was averted by forming a tent under which was introduced hot steam medicated with eucalyptol and the application of the hot pack to the neck.

I have thus briefly sought to direct your attention to the use of a remedy that as nearly approaches a specific for the disease as possible. Hoping and believing that you will find as much satisfaction in its use as I have done, I commend it to your favorable consideration.

A NEW APPARATUS FOR NEBULIZING NON-VOLATILE MEDICAMENTS

AND FOR ADMINISTERING THE SAME THROUGH IMPACTED AIR, AS A MEDIUM, BY THE PROCESS OF FORCED DILATATION, INSTEAD OF BY INHALATION, SPRAY, OR DOUCHE.

BY GEORGE F. HAWLEY, M.D.

Formerly House Surgeon at the Throat and Chest Hospital, Golden Square, London; Assistant Surgeon to the Ear Department of the London Hospital; late Assistant Surgeon to the Illinois Eye and Ear Infirmary, Chicago; Attending Physician Railway Brotherhood Hospital; Member Chicago Medical Society; Professor of the Hygiene and Anatomy of the Throat, Chicago College of Vocal and Instrumental Art.

The object of this paper is to describe an apparatus for preparing and administering nebulous medicaments by a process of forced dilatation, the basis of which is the discovery that the lips will act as a safety valve, and that, into any tube, cavity or air cell into which air can be forced by a high pressure, the nebulous medicament prepared by this apparatus can be forced with absolute safety, thereby enabling the physician to use agents that are capable of preventing or controlling diseases.

My attention was first called to this apparatus and process by an expert in the chemical and mechanical phenomena of pneumatics. I am using the apparatus as auxiliary to the regular medical and surgical treatment of the ears, nose, throat, and chest, with decided advantage to myself, and with most gratifying results to my patients.

The apparatus as shown in the illustration consists of a number of strong flint glass bottles, one for each medicament; adapted to be used in connection with any compressed air receiver. The valve located at the top of each bottle is provided with inlet and outlet ports in multiple, both ports being simultaneously opened or closed by turning the faucet. The inlet port of each bottle is connected with the same source of supply, namely, a compressed air receiver, and also with the vulcanite tube that descends into the medicated solution in the bottle. On turning the faucet the impacted air passes from the receiver through the inlet port down the vulcanite tube, through an ingeniously constructed device, and by eduction draws the medicated solution up into the tube and forces it out against the interior wall of the bottle, thus producing an impacted, nebulous medi-

cament. To the outlet port is attached a silk covered, flexible tube, the free end of which terminates in a vulcanite mouthpiece adapted to become sealed when the lips are tightly compressed around it, also an adjustable nozzle which becomes sealed when pressed into the nostril. Through these tubes the impacted-medicinal-nebula is forced into all the air passages, cavities and cells through and from the bottle selected; by a process of forcible dilatation, —a process that is rational, and capable of producing results that can not be effected by the use of any apparatus adapted to administer by inhalation, spray or douche.

The process of administering consists of: 1, compressing the air or gas into a suitable receiver to a pressure of from 20 to 100 pounds to the square inch or to a density sufficiently great to rapidly fill and forcibly dilate the air cells, the bronchi, the nasal cavities and the Eustachian canals by the expansive force of its own elasticity; 2, forming a communication through suitable tubing to be placed in the mouth or nose in such a manner as to allow the excess of pressure to escape from the mouth. The bottles containing the medicated solutions, being located intermediate and through which the compressed air passes at a uniform pressure with that in