The Journal of the American Medical Association

VOL. XXVIII.

CHICAGO, ILL., MAY 22, 1897.

No. 21.

LECTURE.

NERVOUS DYSPEPSIA.

A clinical lecture delivered in Rush Medical College.

BY HENRY M. LYMAN, A.M., M.D.

PROFESSOR OF THE PRINCIPLES AND PRACTICE OF MEDICINE.

Gentlemen:—You have recently had opportunities for witnessing the effects of inflammation of the mucous coat of the stomach in several patients who have exhibited, with varying degrees of severity, the symptoms of acute and chronic gastritis. This morning I wish to call your attention to a case that illustrates another form of gastric disturbance—nervomuscular dyspepsia. It has been taught by the dominant school of pathologists that such disturbances of digestion are always due to catarrhal gastroenteritis. But the opinion has long been held, and is gaining ground, that in certain cases the disorder of digestion is dependent upon alterations in the nervous system, and that inflammation takes little or no part in its causation. This is apparently true of the gastric crises of *tabes dorsalis*; though, in default of an autopsy, it is always possible to imagine the existence of a latent inflammation that is only prominent during a paroxysm of gastric disturbance. But, on the whole, it seems more likely that the relation in such, and in similar cases, is analogous to what is sometimes observed in the laryngeal cavity when its muscular apparatus is attacked by muscular rheumatism, producing hoarseness or complete aphonia without actual inflammation of the overlying mucous membrane until a day or two after the beginning of recovery from the muscular disorder. In like manner severe pain and disordered secretion may occur in the gastrointestinal canal without manifestation of the ordinary symptoms of inflammation until after the beginning of digestive disturbance.

Nervomuscular dyspepsia may therefore be defined as a complex of symptoms, dependent chiefly upon a disordered condition of the nervous system. The history of the patient before you will amply illustrate this proposition. He is a thin and angular person, about 45 years of age, who from boyhood has been engaged in sedentary and intellectual pursuits. He was an unusually good scholar while a student in the public schools. He was subsequently employed as a general utility man in connection with a country newspaper, to which he occasionally contributed an effusion of his own. Then he studied law in a night-school, and tried to get practice in a large city, but broke down in the attempt. After this experience he was for a number of years employed as a copyist in a government office. He could never afford to take a wife, but admits occasional excursions on the street, and probably rather frequent masturbatory excesses. He, however, has thus far escaped syphilitic infection, and denies having had gonorrhea. He is almost

totally abstinent from alcohol, but is an almost constant smoker, latterly using a pipe instead of cigars, which he can no longer afford to purchase. For three or four years he has been “down on his luck”—out of employment part of the time, and continually harassed by debt and its accompanying trials. Naturally gifted with good health, his elastic temperament has long preserved him from serious disease, but during the past year he has lost vigor, is depressed in spirits and suffers with frequent pain in the epigastric region. This pain, however, is not aggravated by taking food. On the contrary, it is for a time quite abolished after each meal, and recurs about three hours after eating, when the stomach is nearly or quite empty. A glass of milk and one or two crackers will then restore comfort again. Sometimes he is awakened in the night with similar sensations of distress, so that he has learned to carry a crust of bread to bed with him, for use before morning. On examination of the gastric juice, free hydrochloric acid is found to be present in more than the ordinary proportion. Other acids are insignificant in quantity, and cause no considerable difficulty. The gastric contents that are obtained with the aid of the syphon after a test meal are simply the remains of that meal alone, showing that the stomach is capable of emptying itself into the duodenum within four or five hours. Muscular atrophy of the stomach can be therefore excluded.

But here is another patient who, if subjected to this test, furnishes the debris of one or two previous repasts in addition to the substances derived from the test meal. He presents the usual symptoms of chronic catarrhal gastritis, and on rising in the morning splashing sounds can be heard in his stomach if he be vigorously shaken. Inflation of his stomach places its greater curvature a little below the umbilicus. His bowels are constipated, and he suffers frequently with intestinal flatulence. In his gastric juice there is deficiency, but not complete absence, of free hydrochloric acid. In short, without going farther into particulars, we may say that he is a victim of gastric dilatation not dependent upon malignant disease, but accompanied by frequent attacks of pain and gastrointestinal uneasiness, which are only less severe than the paroxysms of *gastralgia* that frequently occur as a consequence of chronic gastritis.

Between these types of neuro-muscular distress many shades of disorder may be discovered, but they can be all grouped into a few classes. An attempt has been made to form a classification based upon the chemic qualities of the gastric juice with reference to the relative quantity of hydrochloric acid that may be present. But the amount of acid thus present in the juice is quite variable, while the nervous phenomena are always predominant, and should be principally considered both in the diagnosis and in the treatment of nervous dyspepsia. The relative acidity and

catarrhal condition of the gastric contents are subordinate factors that must receive due consideration, but may not be made preëminent. It is, accordingly, advisable to regard nervous dyspepsia as comprising those digestive disturbances that are chiefly dependent upon disorder of the nervous system, and are characterized by symptoms of nervous derangement. Like all other nervous phenomena, therefore, there may be either exaltation, depressions or incoördination of function, giving rise to increase, or reduction, or perversion of sensation, muscular movement, circulation and secretion in the stomach. Indigestion may be, therefore, either painful or painless; it may be hurried beyond measure, or it may be retarded by atonic conditions of the muscular fibers in the gastro-intestinal wall; there may be excessive hyperemia and inordinate secretion in the mucous membrane, or the contrary condition may obtain; and finally, as a result of incoördination of nervous functions, the quality of the gastric juice, in the matter of digestive ferments and hydrochloric acid, may exhibit considerable variation from the normal standard.

Obviously, a chemic examination of the gastric juice is needful for the determination of these different qualities of the digestive fluids. Dyspepsias have been, therefore, subjected to a chemic classification based upon the relative quantities of hydrochloric acid that can be demonstrated by analysis. In certain cases the acid is present in excess (hyperchlorhydria), while in others it is deficient (hypochlorhydria), or entirely absent. But in a considerable number of instances clinical observation discovers evidence of disordered digestion without any notable deviation from the normal acidity of the gastric juice. So complicated is the web of symptoms that it is impossible to arrange all cases under a few simple species, like animals in a menagerie. Each patient exhibits individual symptomatic peculiarities, like the numerous varieties presented by certain plants whose species shade into each other after a fashion that renders exact classification almost impossible. But, for the sake of convenience, we may recognize three principal groups into which nervous dyspepsias may be divided—always, however, remembering their susceptibility to variation, and their consequent liability to pass from one group into another, or even to merge themselves among the host of inflammatory disorders in which nervous phenomena occupy a comparatively subordinate position. These three groups may be conveniently considered in succession.

SIMPLE NERVOUS DYSPEPSIA.

This form of digestive disorder is most frequently encountered among neuropathic subjects with rickety antecedents, or among members of the arthritic family. Retarded dentition, profuse sweating during sleep, local spasms, laryngismus stridulus, and intellectual precocity are of common occurrence in the early life of these patients. As the years advance, many of them become victims of eczema, dandruff, early whitening of the hair, premature baldness, bronchitis, asthma, palpitation and irregularity of the heart, neuralgia, gallstones, renal calculi, varicose veins, hemorrhoids, diabetes, polyuria, obesity, rheumatism and gout. Catarrhal conditions of the alimentary canal are often experienced, especially in the childhood and old age of such persons. Dyspepsia thus inaugurated frequently is associated with or replaces the phenomena of muscular and nervous rheumatism. It is easily

induced by fatigue of mind and body and is one of the most common causes of persistent neurasthenia.

The general health and appearance of the patient exhibit little alteration in this form of dyspepsia, unless the function of the small intestine be seriously hindered. There is frequent complaint of epigastric uneasiness after eating, and this is accompanied by a feeling of distension in the stomach, and by eructations of tasteless and odorless gas. Sometimes there is a sensation of impediment in the act of respiration "as if the breath would not go down to the bottom of the lungs." The head aches; the countenance is flushed; the bowels are constipated, and hemorrhoids often exist. Shortly after eating, considerable pain may be experienced; it is a dull, vague sensation of distress that is not definitely located, but is felt with varying severity throughout an ill-defined region beneath the diaphragm. Sometimes the patient experiences vertigo, or feels as if a band compressed the cranium above the eyebrows; occasionally the temporal arteries beat violently, and the corresponding veins are distended with blood. In certain cases it is impossible to lean forward without exciting severe pain in one or both sides of the head. There may be a feeling of weight or pressure in the occipital region, and it is not an uncommon event for the patient to hear a cracking sound in the back of the neck when the head is rotated upon the axis, especially on turning from side to side in bed. Usually there is great depression of spirits, and a tormenting fear of impending evil. The feelings exhibit great instability, passing rapidly from an extreme of vivacity to the profoundest depths of despair.

If the abdomen be examined an hour or two after a meal, considerable gaseous distension of the stomach and intestines will be discovered. Sometimes a succussion sound is audible in the stomach if the body be violently shaken. The differential diagnosis between simple distension of the stomach caused by nervous dyspepsia and permanent dilatation of the organ can be established by examination before breakfast, at which time succussion sounds can not be evoked in cases of temporary distension; whereas they can be always demonstrated when there is persistent dilatation of the stomach.

Palpation of the abdominal wall, in nervous dyspepsia, seldom indicates any extensive tenderness or special uneasiness under pressure. Frequently, however, a moderate degree of tenderness can be detected just below the tip of the ensiform cartilage, and in rheumatic subjects the upper portion of the recti muscles is often sensitive under manipulation.

Provided that intestinal digestion be not seriously hindered, the amount of free hydrochloric acid in the gastric juice may vary considerably without materially affecting the nutrition of the patient. Mere chemic examination of the gastric juice, therefore, affords little information regarding the totality of the digestive process. Of far greater importance is the condition of the nervous system. So long as actual dilatation of the stomach does not exist there will be no extraordinary delay in the passage of food into the intestine, there will be little if any fermentation, and no remarkable failure of health, even though the amount of free hydrochloric acid in the gastric juice should fall below the normal standard.

Such disturbances of digestion as are present are not confined to the stomach alone; they also involve the intestinal canal. In many instances there is some

distension of the colon; constipation and hemorrhoids afford evidence of intestinal and hepatic disorder. Occasionally the phenomena of pseudo-membranous enteritis make their appearance.

NERVOUS DYSPEPSIA WITH EXCESSIVE SECRETION OF HYDROCHLORIC ACID.

In certain cases of nervous dyspepsia accompanying neurasthenia the secretory nerves that should regulate the formation of gastric juice do not exert their normal coördinative influence, and there is an excessive production of hydrochloric acid. Sometimes the acid thus furnished is sufficient to be a veritable cause of inflammation. Such excessive secretion is not unusual during the paroxysms of vomiting that constitute the gastric crises of *tabes dorsalis*.

In typical cases of this variety of dyspepsia great pain, beginning two or three hours after a meal, is experienced in the epigastric region. Considerable relief may be procured by swallowing milk, or an aqueous solution of an alkaline salt. It is completely removed by the ordinary meals, which, virtually, are equivalent to the administration of a full dose of alkaline salts; but the cessation of pain is only temporary, and suffering is renewed as soon as the stomach is empty again. In this way sleep is often disturbed after midnight by a burning sensation, or by vague feelings of uneasiness in the epigastrium, sometimes radiating into the interscapular region. Vomiting may occur, and it is sometimes voluntarily excited by the patient, who has learned how much relief may be obtained through evacuation of the stomach. Similar relief is afforded by irrigation of the gastric cavity. In many cases, however, there is absence of nausea, but the acid contents of the stomach are eructated, causing an intensely burning sensation in the esophagus and fauces. Such patients are thirsty; they often have a craving appetite that impels them to its frequent gratification; and they instinctively prefer animal food, because it is digested more easily than vegetable food. Usually, there is considerable emaciation; the skin assumes an earthy hue; nervous exhaustion is indicated by irritability, fatigue, restlessness and inability to sleep. Many of the sufferers are despondent and hypochondriacal to the last degree.

Milder forms of the disease are less painful. There is merely a sensation of epigastric uneasiness beginning within several hours of eating and arrested by the following repast. These patients desire food at short intervals, and if the usual mealtime is passed without taking nourishment they experience a feeling of exhaustion with distress at the pit of the stomach, often accompanied by a throbbing headache and vertigo. They are generally nervous, irritable, neuralgic and predisposed to arthritic disorders. The bowels are constipated, though occasionally there is obstinate diarrhea—especially in the night or early in the morning—that does not yield to astringents and opiates. Sometimes tenderness on pressure is felt over the pyloric region, and percussion indicates distension of the stomach. Actual dilatation is not to be presumed unless a succussion sound can be obtained in the morning before food has been taken. Catheterization of the stomach yields a liquid that is strongly acid, ropy and of a yellowish color. By washing out the stomach at bedtime, and on the following morning removing the liquid contents of the organ before food has been taken, it is easy to deter-

mine the fact of hypersecretion and gastric dilatation if hydrochloric acid be present.

In a certain proportion of these cases the patient is tormented by repeated attacks of vomiting, which may continue for several days together. Such attacks are frequently induced by intellectual fatigue, and are an expression of nervous disorder accompanied by headache, pain in the stomach and nausea; all of which may be relieved by vomiting the excessively acid contents of the stomach or by their neutralization with an alkaline solution.

NERVOUS DYSPEPSIA WITH DIMINISHED SECRETION OF HYDROCHLORIC ACID.

In the third variety of nervous dyspepsia the symptoms do not materially differ from those which are experienced in the previous form; but the amount of hydrochloric acid, both free and combined, is considerably reduced, or may be temporarily absent. Fermentation consequently occurs in the contents of the stomach, liberating various organic acids and acid substances. In such cases there is loss of appetite, pain in the epigastric region, distressing eructation and frequent vomiting. The stomach is distended with gas, and succussion sounds are distinctly audible; pain is felt soon after eating, but it is usually less severe than the suffering that is caused by an excess of hydrochloric acid. In many cases there is regurgitation of bile into the stomach, hindering gastric digestion and adding to the discomfort of the patient who loses flesh and acquires a cachectic hue suggestive of cancer of the stomach.

Evidently, these forms of dyspepsia can be differentiated only by the aid of chemical analysis. The mere presence or absence of free hydrochloric acid can be readily determined by the ordinary color tests; but for an accurate diagnosis a quantitative analysis is required. The prognosis depends upon the duration of the disease and its severity. Mild and recent cases occurring in young subjects, usually recover, but dilatation of the stomach, and excessive secretion of hydrochloric acid are unfavorable symptoms. In the severer forms there is reason to fear the passage from simple nervous excitement to actual inflammation, ulceration, or malignant disease.

The indications for treatment are three-fold: 1, to relieve pain; 2, to correct nervo-muscular disorder, and 3, to rectify the secretions.

1. Uneasy sensations occurring when the stomach is empty, or when it contains an excessive quantity of hydrochloric acid, may be temporarily relieved by the administration of milk, or water containing bicarbonate of sodium. But when there is intense gastralgia nothing less than hypodermic injections of morphin and atropin will give relief. Moderate pain may be suppressed by the use of Hoffmann's anodyne, cannabis indica, cocain, hyoscyamus, or a saturated aqueous solution of chloroform. This last remedy is the most prompt and efficient of all, except morphin. In every case of severe pain the stomach should be thoroughly evacuated by an hypodermic injection of one-tenth of a grain of apomorphia. A large poultice, into which have been stirred a drachm each of chloroform and tincture of opium, should be laid over the whole abdomen and lower half of the thorax. A cathartic should be administered as early as possible; it may consist of ten grains of calomel with five grains of bicarbonate of sodium, to be followed four hours later by an ounce of castor oil. Sometimes considerable relief

may be derived from copious draughts of hot water. Solid food should not be taken until the acute symptoms have yielded to treatment and rest.

2. Inefficient muscular contraction in the walls of the stomach and intestines demands the use of nuxvomica, ipecac, massage, electricity, injections of hot water into the colon, and stimulant laxatives. Nuxvomica is a better tonic than strychnin; it may be given in doses of five to ten drops of the tincture, a quarter of a grain of the extract, or half a grain of the powdered nut, after each meal. Ipecac stimulates the secretion of bile, and is especially useful when there is gastro-intestinal flatulence after eating; it should be given in doses of one-tenth of a grain, associated with nuxvomica, piperin, ginger, and other aromatic stimulants, as in the well-known pill of Fothergill, which may be thus modified:

R. Pulv. ipecac. . . . . gr. 1-10
Ext. nuc. vom. . . . . gr. ¼
Pulv. piper. nig. . . . .
Ext. gentian . . . . . āā gr. i
Fiat pil. No. 1. Sig. One such pill after each meal.

Alain in appropriate quantity may be added when there is constipation; and an occasional dose of blue mass and colocynth is often useful.

3. For the chemic rectification of the secretions, when there is excessive production of hydrochloric acid, the patient must abstain from alcoholic beverages and highly seasoned food. Mastication must be slowly and thoroughly performed. Starchy and saccharine substances should not be eaten; milk, eggs, peptonized meat powders, minced meat, tender steak, and other nitrogenous articles of diet should constitute the bulk of the food. When the excess of acid is present during the period of digestion alone, causing pain as the stomach becomes empty, sodium bicarbonate may be given in doses of five to twenty grains about three hours after each meal. If the bowels be constipated an equal quantity of Hubbard's magnesia may be added to each dose. When the stomach is continually charged with hydrochloric acid it will be necessary to increase the dose of sodium bicarbonate to four or six drachms a day. Vichy water, lime water, lithia water, and Murray's fluid magnesia may also be used as antacids. When there is evident dilatation of the stomach with stagnation of its contents, the syphon should be employed, in order to remove the irritating liquids. In default of such apparatus, the older physicians gave emetics of lobelia with good results.

When, without dilatation of the stomach or stagnation of its contents, there is a deficiency of hydrochloric acid in the gastric juice, ten or fifteen drops of the dilute acid may be given in a teacup of hot water after each meal. An equally good result follows the similar use of the other mineral acids; hence the extensive popularity of Horsford's Acid Phosphate, and kindred acid preparations. But it must be noted that acids and pepsin and other digestives so often administered are not curative of dyspepsia. They are merely palliative, and give only partial and temporary relief. The disease can be overcome only by the removal of its cause, whatever that may be. In the vast majority of cases it is an exhausted state of the nervous system that underlies the disorder, and the patient must reform his whole course of life before he can be cured.

Dilatation of the stomach, and an excessive acidity due to fermentation and the production of the organic

acids in its contents require daily irrigation of the stomach, and the administration of salol, salicylic acid or beta-naphthol, before meals. In mild cases much benefit is derived from the long continued administration of small doses of the alkaline salts. Potassium citrate or acetate or carbonate, with sodium bicarbonate, associated with a bitter tonic like nuxvomica, compound tincture of gentian, or the elixir of calisaya bark, electricity, massage and hydrotherapy are indispensable in this class of cases. Change of occupation, change of habits, the attainment of prosperity, long vacations, mountain climbing, sea-voyaging, and cool baths are far more effectual than the most elaborate medication.

ORIGINAL ARTICLES.

PRIMARY PAPILLOMA AND PRIMARY CARCINOMA OF THE FALLOPIAN TUBE.

Read at the Meeting of the Chicago Pathological Society, April 12, 1897.

BY EMIL RIES, M.D.

PROFESSOR OF GYNECOLOGY, POST-GRADUATE MEDICAL SCHOOL, CHICAGO.

The two cases which I present here were operated on by Dr. T. J. Watkins, who kindly offered them to me for pathologic examination. They are of great clinical as well as pathologic interest, one case indeed being unique in its kind. Only six cases of primary papilloma of the tube are contained in the literature, the present being the seventh case. Of primary carcinoma of the tube the tables of Doran¹ and of Saenger² contain nineteen cases, the present being the twentieth.

I. Primary papilloma of the tube. Secondary papilloma of both ovaries. Repeated vaginal drainage for purulent salpingitis, at last salpingo-oophorectomy by laparotomy. Recovery. Good health fifteen months after the operation.

Clinical history.—Patient, 34 years old, gives a history of inflammatory disease of the pelvic organs after a miscarriage in her twenty-ninth year, from which she recovered perfectly. In her thirtieth year she suffered some violent injury to her left inguinal region. This was followed by "inflammation of the ovary," which remained painful until operation. The diagnosis was pyosalpinx and in June, 1894, the swelling in the pelvis was opened through the vagina. The tumor evacuated a quantity of pus. In November, 1894, the tumor had to be opened again and patient wore a drainage tube for eight months. In August, 1895, third operation. Patient wore the drainage tube for two months after she left the hospital. At last in January, 1896, laparotomy was performed and the appendages were removed. The patient recovered uneventfully. She was seen a few days ago by Dr. Watkins and was found to be in perfect health. She does not menstruate and has no leucorrhea. The uterus is atrophied and freely movable.

Macroscopic description of specimens.—Left side: Ovary not enlarged (35 x 26 x 20 mm.). Its surface is covered in several places by wart-like excrescences, which in some places sit on the ovary with a broad base, while in others they are distinctly pedunculated. The tube is normal externally with the exception of a slight thickening of its isthmus part. The fimbriated end is open. Cross sections through the tube show nothing abnormal. The ovary contains on sec-